(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(84) Designated Contracting States:
DE ES FR GB IT

(30) Priority: 14.02.1995 JP 25398/95
14.02.1995 JP 25402/95
12.09.1995 JP 234236/95
20.10.1995 JP 272432/95

(71) Applicant:
MITSUBISHI CHEMICAL CORPORATION
Chiyoda-ku, Tokyo 100 (JP)

(72) Inventors:
• ANDO, Ryoichi
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• SAKAKI, Toshiro
Mitsubishi Chemical Corporation
2-chome Chiyoda-ku Tokyo 100 (JP)
• MASUDA, Hirokazu
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• INAKOSHI, Naoto
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)

• JIKIHARA, Tetsuo
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• FUJIMURA, Yoshiyuki
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• NIWA, Takuro
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• YOSHII, Narihiko
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• TABATA, Reiko
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• SAITO, Ken-Ichi
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)
• ARITOMO, Keiichi
Mitsubishi Chemical Corporation
Aoba-ku, Yokohama-shi Kanagawa 227 (JP)

(74) Representative:
Godemeyer, Thomas, Dr.
Hauptstrasse 58
51491 Overath (DE)

(54) **OXYGEN-CONTAINING HETEROCYCLIC DERIVATIVES**

(57) An oxygen-containing heterocyclic derivative represented by the following formula (I) and its salt, and a solvate and a hydrate thereof:

EP 0 810 221 A1

$$R^1 \left(\underset{\underset{R^2}{\overset{R^3}{|}}}{\overset{}{N}} \underset{\underset{O}{\parallel}}{C} \right)_n \overset{R^4}{\underset{}{N}} \underset{\underset{R^5}{|}}{CH} \underset{O}{\overset{\parallel}{C}} \overset{}{\underset{\underset{R^6}{|}}{N}} CH \overset{A}{\underset{OR^7}{}} O$$

(I)

wherein:

R^1:	hydrogen atom, R^8-CO-, R^8-O-CO- or other wherein R^8 represents C_1-C_{20} alkyl, aryl or other,
R^2, R^4 and, R^6:	a hydrogen atom, C_1-C_5 alkyl or other,
R^3 and R^5:	a hydrogen atom, C_1-C_{20} alkyl or other,
R^7:	a hydrogen atom, R^9-CO- or other wherein R^9 represents C_1-C_{10} alkyl group or other,
A:	C_1-C_3 alkylene group
n:	0 or 1.

The oxygen-containing heterocyclic derivatives of the present invention have potent inhibitory activity against cysteine proteases and are excellent in oral absorbability, tissue distribution, and cellular membrane permeability. Therefore, they are useful as therapeutic medicament for diseases such as cerebral apoplexy, Alzheimer's disease and the like.

Description

Technical Field

5   This invention relates to novel oxygen-containing heterocyclic derivatives.

Background Art

Functions of cysteine proteases such as papain, cathepsin B, cathepsin H, cathepsin L, calpain, and interleukin-1
10   β converting enzyme in living bodies have been elucidated, and as the progress, it becomes revealed that abnormal
accentuation of these substances is a cause of various types of diseases. There also are increasing numbers of litera-
tures reporting that a cysteine protease inhibitor was found to be effective when applied to an animal model of such dis-
ease.
      Cysteine proteases such as calpain or cathepsin B are considered to be involved in an early stage of elimination of
15   Z-filaments or other with degradation of muscle fiber proteins in a process of skeletal muscular decay which can be
observed in myopathy such as muscular dystrophy and amyotrophia (Metabolism, Vol. 25, Extra Edition, "Highlights of
Metabolic Disease," p. 183, 1988). In addition, E-64-d, a cysteine protease inhibitor, was reported to have life prolonga-
tion effect on muscular dystrophy hamsters (Journal of Pharmacobio Dynamics, Vol. 10, p. 678, 1987). Accordingly,
cysteine protease inhibitors are considered as potential therapeutic medicaments for muscular dystrophy, amyotrophia
20   and the like.
      In ischemic diseases such as myocardial infarct and cerebral apoplexy, a major cause of cellular dysfunction after
ischemia is active oxygen produced by xanthine oxidase. Some articles suggested that calpain, being activated by an
increased concentration of Ca^{2+} during ischemia, partially degrades xanthine dehydrogenase as a precursor of xan-
thine oxidase to convert it into the oxidase (New England Journal of Medicine, Vol. 312, p. 159, 1985). It is also sug-
25   gested that the activation of calpain may directly trigger myocardial cellular death and cerebral nerve cellular death
(Latest Medicine, Vol. 43, p. 783, 1988). NCO-700, an inhibitor of calpain, was reported to be effective in an animal
model of myocardial infarct (Arzneimittel Forschung/Drug Research Vol. 36, p. 190, p. 671, 1986) and E-64-c sup-
pressed the degradation of microtubule-binding proteins after cerebral ischemia (Brain Research, Vol. 526, p. 177,
1990). Therefore, calpain inhibitors are considered as potential therapeutic medicament for ischemic diseases such as
30   myocardial infarct and cerebral apoplexy.
      A protein called amyloid deposits in senile plaques that can be uniquely observed in brains of patients suffered from
Alzheimer's disease. It is known that the amyloid is synthesized by the degradation of amyloid protein precursor (APP).
Some articles suggest that amyloid is produced by abnormal metabolism due to an abnormally accentuated protease
to form senile plaques, whilst amyloid is not produced in a normal metabolic process of APP (Scientific American,
35   November, 1991, p. 40). Accordingly, protease inhibitors are expected to be used as therapeutic medicament for Alzhe-
imer's disease.
      There is reported that calpain is activated in a head injury model using a rabbit (Neurochemical Research, Vol. 16,
p. 483, 1991). A protective effect on axon was observed by the administration of leupeptin as a calpain inhibitor to a
head injury model using a rat (Journal of Neurosurgery, Vol. 65, p. 92, 1986). Therefore, inhibitors of calpain are con-
40   sidered to have an improving effect on disturbance of consciousness and movement disorder caused by a head injury.
      Myelin binding proteins that exist in dendrites of nerve cells were found to be degraded by calpain (Journal of Neu-
rochemistry, Vol. 47, p. 1007, 1986). Therefore, inhibitors of calpain are considered as effective on diseases caused by
demyelination of nerve cells such as multiple sclerosis and neuropathy of peripheral nerve.
      In most cases of cataract, it is suggested that crystallin, a water-soluble protein in crystalline lens, is hydrolyzed by
45   the action of a protease, which results in the cloudiness of crystalline lens. In experimental models of cataract and cer-
tain types of human cataract, calcium concentrations in crystalline lens are increased (Investigative Ophthalmology &
Visual Science, Vol. 28, p. 1702, 1987; Experimental Eye Research, Vol. 34, p. 413, 1982), and calpain exists in major
abundance among proteases contained in crystalline lens (Lens and Eye Toxicity Research, Vol. 6, p. 725, 1989).
Therefore, abnormal accentuation of calpain is considered to be one of causes of cataract. There is also reported that
50   an inhibitor of calpain, i.e., E-64, was effective in an experimental model of cataract (Investigative Ophthalmology & Vis-
ual Science, Vol. 32, p. 533, 1991. Accordingly, inhibitors of calpain are considered as potential therapeutic medicament
for cataract.
      It is known that neutrophils, which are intimately involved in inflammation, respond to stimulations with chemotactic
factors or phorbol esters by causing degranulation or producing superoxide, and this process is considered to be medi-
55   ated by protein kinase C (PKC). There is reported that calpain has a function to activate PKC, thereby exhibits promot-
ing activity on the degranulation and suppressing activity against the production of superoxide (Journal of Biological
Chemistry, Vol. 263, p. 1915, 1988). It has also been reported that concentration of cathepsin B in rat macrophages is
30 to 40-fold higher than that of leucocytes or neutrophils, and moreover, th enzyme concentration of inflammatory
macrophages is 6-fold higher than that of ordinary macrophag s (Journal of Biochemistry, Vol. 98, p. 87, 1985). Fur-

thermore, it has recently been revealed that the enzyme that catalyzes th conversion of pre-interleukin-1 β into interleukin-1 β (interleukin-1 β converting enzym) is a cysteine protease (Nature, Vol. 356, p. 768, 1992), which clarifies that the activation process of cysteine protease plays an important role in the formation of inflammation. From these findings, it is considered that inhibitors of cysteine protease can be used as anti-inflammatory agents.

I-type allergic reaction progresses with the mediation of immunoglobulin E (IgE) that is produced by immunization of a living body with an antigen. It has been reported that Estatin A, a cysteine protease inhibitor, specifically suppresses the production of IgE, whereas it does not affect on the production of IgG (The Journal of Antibiotics, Vol. 42, p. 1362, 1989). Accordingly, cysteine protease inhibitors are considered to be usable as anti-allergic agents.

When necrosis of hepatocytes occurs, it is suggested that Ca^{2+} permeability of cell membranes increases due to disturbance of cellular membranes and then intracellular Ca^{2+} concentration is elevated to activate calpain, which leads to degradation of structural proteins or other as substrates of calpain and results in cellular death. Therefore, inhibitors of calpain can be used as therapeutic medicaments for fulminant hepatitis.

Cathepsins such as cathepsin B and cathepsin L participate in degradation of bone collagen in osteoclasts. There is reported that serum calcium concentration and hydroxyproline concentration were lowered when B-64 or Estatin A being an inhibitor of cathepsins was administered to rats whose osteoclasia was accentuated by administering par-ath rmone (Biochemical and Biophysical Research Communication Vol. 125, p. 441, 1984; the Japanese Patent Unexamined Publication (KOKAI) No. (Hei) 2-218610/1990). Therefore, inhibitors of cathepsins are considered as potential therapeutic medicaments for osteoporosis, hypercalcemia and the like.

Substrates of calpain include a class of sexual hormone receptors such as estrogen receptor and androgen receptor. Calpain is known to activate these receptors, and it is suggested that abnormal accentuation of calpain causes diseases that are positively caused by abnormal activation of sexual hormone receptors such as, for example, breast cancer, prostatic cancer, and prostatic hypertrophy. Accordingly, inhibitors of calpain are considered to be useful as therapeutic medicaments for the aforementioned diseases.

It is suggested that receptors of epidermal growth factor (EGF) are activated as cellular tumorigenic transformation proceeds, and it is known that calpain activates the EGF receptors as its substrates. There is also reported that calpain was activated in cells infected with human adult T-cell leukemia virus (ATLV/HTLV-1) (Biochemistry, Vol. 57, p. 1202, 1985). On the other hand, it is suggested that cathepsin B is intimately involved in processes of cancerous metastasis, because cathepsin B promotes collagen degradation, which is an important step of cancerous metastasis, or directly degrades collagen, and it also has close relationship with plasma membranes of neoplastic cells (Tumor Progression and Markers, p. 47, 1982; Journal of Biological Chemistry, Vol. 256, p. 8536, 1984). From these teachings, inhibitors of cysteine protease are considered to be effective in inhibition of cancerous growth and prevention of cancerous metastasis.

Activation of blood platelets triggers agglomeration, which may lead to thrombus formation. There is reported that E-64-d as being an inhibitor of calpain suppressed platelet aggregation caused by thrombin (Thrombosis Research, Vol. 57, p. 847, 1990). Accordingly, inhibitors of calpain can be used as platelet aggregation inhibitors.

As explained above, abnormal accentuation of cysteine protease may become the causes of various kinds of diseases, and some cysteine protease inhibitors are reported to have efficacy in animal models or the like.

However, almost all of known inhibitors are irreversible inhibitors, for example, E-64 (Agricultural and Biological Chemistry, Vol. 42, p. 529, 1978), E-64-d (Journal of Biochemistry, Vol. 93, p. 1305, 1983), NCO-700 (the Japanese Patent Unexamined Publication (KOKAI) No. (Sho) 58-126879/1983), epoxysuccinic acid derivatives such as Estatins A and B (The Journal of Antibiotics, Vol. 42, p. 1362, 1989), and α -substituted ketones of peptides whose typical examples include chloromethyl ketones of peptides (Journal of Biochemistry, Vol. 99, p. 173, 1986) and acyloxymethyl ketones (Biochemistry, Vol. 30, p. 4678, 1991). Irreversible inhibitors are generally considered to have strong toxicity because they are likely to non-specifically react with biogenic components including the target enzymes, and therefore, only a few compounds have been used clinically. As reversible inhibitors, peptidyl aldehydes such as leupeptin (The Journal of Antibiotics, Vol. 22, p. 283, 1969) and calpeptin (Journal of Enzyme Inhibition, Vol. 3, p. 195, 1990) are known. However, they are considered to have problems relating to chemical stability, stability in a living body, cell membrane permeability and the like.

Disclosure of the Invention

From these reasons, the present inventors conducted researches to obtain cysteine protease inhibitors having excellent oral absorbability, tissue distribution, cell membrane permeability and other. As a result, they achieved the present invention.

The gist of the present invention is oxygen-containing heterocyclic derivatives represented by the following formula (I) and their pharmaceutically acceptable salts, and hydrates and solvates thereof:

(I)

wherein R^1 represents a hydrogen atom, R^8-CO-, R^8-O-CO-, R^8-NH-CO-, or R^8-SO$_2$- (R^8 represents a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of a C_3-C_8 cycloalkyl group, a C_3-C_8 cycloalkyloxy group, fluorenyl group, a C_1-C_6 alkoxy group, a C_6-C_{14} aryl group which may optionally be substituted, a C_6-C_{14} aryloxy group which may optionally be substituted, a C_6-C_{14} arylthio group which may optionally be substituted, a C_6-C_{14} arylsulfonyl group which may optionally be substituted, and a residue of a heterocyclic compound which may optionally be substituted; a C_3-C_8 cycloalkyl group; a C_6-C_{14} aryl group which may optionally be substituted; a C_2-C_5 alkenyl group which may optionally be substituted with an optionally substituted C_6-C_{14} aryl group; or a residue of a heterocyclic compound which may optionally be substituted); R^2, R^4, and R^6 independently represent a hydrogen atom, a C_1-C_5 alkyl group, or a C_2-C_6 alkanoyl group; R^3 and R^5 independently represent a hydrogen atom, a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of a C_6-C_{14} aryl group which may optionally be substituted, hydroxyl group, a C_1-C_5 alkoxy group, a C_1-C_5 alkylthio group, and a C_7-C_{12} aralkyloxy group, or they independently represents a C_6-C_{14} aryl group which may optionally be substituted; R^7 represents a hydrogen atom, a C_1-C_5 alkyl group, or R^9-CO- (R^9 represents a C_1-C_{10} alkyl group or a C_6-C_{12} aryl group which may optionally be substituted); symbol "A" represents a C_1-C_3 alkylene group which may optionally be substituted with a C_1-C_3 alkyl group; and symbol "n" represents 0 or 1.

The present invention will be detailed below.

In the aforementioned general formula (I), examples of the C_1-C_{20} alkyl group defined by R^8 include methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, isopentyl group, neopentyl group, tert-pentyl group, hexyl group, isohexyl group, heptyl group, octyl group, nonyl group, decyl group, undecyl group, dodecyl group, tridecyl group, tetradecyl group, pentadecyl group, and octadecyl group. These alkyl groups may have one or more substituents selected from the group consisting of a C_3-C_8 cycloalkyl group such as cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group, and cyclooctyl group; a C_3-C_8 cycloalkyloxy group such as cyclopropyloxy group, cyclobutyloxy group, cyclopentyloxy group, cyclohexyloxy group, cycloheptyloxy group, and cyclooctyloxy group; fluorenyl group; a C_1-C_5 alkoxy group such as methoxy group, ethoxy group, propoxy group, isopropoxy group, butoxy group, isobutoxy group, tert-butoxy group, pentyloxy group, and isopentyloxy group; a C_6-C_{14} aryl group such as phenyl group, naphthyl group, and anthryl group; a C_6-C_{14} aryloxy group such as phenoxy group and naphthoxy group; a C_6-C_{14} arylthio group such as phenylthio group and naphthylthio group; a C_6-C_{14} arylsulfonyl group such as phenylsulfonyl group and naphthylsulfonyl group; and a residue of a heterocyclic compound which contains from 1 to 4 heteroatoms selected from oxygen atom, sulfur atom, or nitrogen atom and has 5 to 10 ring-constituting atoms, and whose examples include furan ring, dihydrofuran ring, tetrahydrofuran ring, pyran ring, dihydropyran ring, tetrahydropyran ring, benzofuran ring, isobenzofuran ring, chromene ring, chroman ring, isochroman ring, thiophene ring, benzothiophene ring, pyrrole ring, pyrroline ring, pyrrolidine ring, imidazole ring, imidazoline ring, imidazolidine ring, pyrazole ring, pyrazoline ring, pyrazolidine ring, triazole ring, tetrazole ring, pyridine ring, pyridine oxide ring, piperidine ring, pyrazine ring, piperazine ring, pyrimidine ring, pyridazine ring, indolizine ring, indole ring, indoline ring, isoindole ring, isoindoline ring, indazole ring, benzimidazole ring, purine ring, quinolizine ring, quinoline ring, phthalazine ring, naphtylidine ring, quinoxaline ring, quinazoline ring, cinnoline ring, pteridine ring, oxazole ring, oxazolidine ring, isoxazole ring, isoxazolidine ring, thiazole ring, thiazolidine ring, isothiazole ring, isothiazolidine ring, dioxane ring, dithian ring, morpholine ring, and thiomorpholine ring. Examples of the C_3-C_8 cycloalkyl group and C_6-C_{14} aryl group defined by R^8 include the same groups as those mentioned as the substituents of the aforementioned C_1-C_{20} alkyl group. Examples of the C_2-C_5 alkenyl group include vinyl group, 1-propenyl group, 2-propenyl group, 1-butenyl group, or 1-pentenyl group. The alkenyl groups may be substituted with a C_6-C_{14} aryl group such as those explained above as the substituents of the aforementioned C_1-C_{20} alkyl group. Examples of the residue of a heterocyclic compound include the same groups as those explained as the substituent of the C_1-C_{20} alkyl group.

The C_1-C_5 alkyl group defined by R^2, R^4 and R^6 may independently be, for example, methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, pentyl group, isopentyl group or the like. Examples of the C_2-C_6 alkanoyl group include acetyl group, propionyl group, butyryl group, valeryl group or the like.

Examples of the C_1-C_{20} alkyl group defined by R^3 and R^5 include independently th same groups as those defined as to R^8. These alkyl groups may have on or more substituents selected from the group consisting of a C_6-C_{14} aryl group such as those explained as to R^8; hydroxyl group; a C_1-C_5 alkoxy group such as those explained abov as to R^8;

a C_1-C_5 alkylthio group such as methylthio group, ethylthio group, propylthio group, isopropylthio group, butylthio group, isobutylthio group, tert-butylthio group, pentylthio group, and isopentylthio group; and a C_7-C_{12} aralkyloxy group such as b nzyloxy group, phenylmethoxy group, and naphthylmethoxy group. Examples of the C_6-C_{14} aryl group includ th same groups as those mentioned above.

Examples of the C_1-C_5 alkyl group defined by R^7 include methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, isopentyl group, neopentyl group, tert-pentyl group or the like.

Examples of the C_1-C_{10} alkyl group defined by R^9 include methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, isopentyl group, neopentyl group, tert-pentyl group, hexyl group, isohexyl group, heptyl group, octyl group, nonyl group, decyl group or the like. Examples of the C_6-C_{12} aryl group include phenyl group, naphthyl group or the like.

Examples of the C_1-C_3 alkylene group defined by the symbol "A" include methylene group, ethylene group, propylene group or the like, and these alkylene groups may have one or two C_1-C_3 alkyl groups such as methyl group, ethyl group and propyl group.

In the aforementioned definition, the aryl group and the heterocyclic residue may further have one or more substituents when they are attached at the end of each functional group, and examples of said substituents include a halogen atom such as fluorine atom, chlorine atom, and bromine atom; a C_1-C_5 alkyl group such as methyl group, ethyl group, propyl group, isopropyl group, butyl group, isobutyl group, sec-butyl group, tert-butyl group, pentyl group, isopentyl group, neopentyl group, and tert-pentyl group; trifluoromethyl group; a C_1-C_5 alkoxy group such as methoxy group, thoxy group, propoxy group, isopropoxy group, butoxy group, isobutoxy group, tert-butoxy group, pentyloxy group, and isopentyloxy group; a C_1-C_5 alkylenedioxy group such as methylenedioxy group, ethylenedioxy group, and propylenedioxy group; hydroxyl group; nitro group; a C_2-C_6 alkylcarbonyloxy group such as acetoxy group, propionyloxy group, butyryloxy group, and valeryloxy group; carboxyl group; a C_2-C_6 alkoxycarbonyl group such as methoxycarbonyl group, thoxycarbonyl group, propoxycarbonyl group, isopropoxycarbonyl group, butoxycarbonyl group, isobutoxycarbonyl group, tert-butoxycarbonyl group, and pentyloxycarbonyl group; oxo group; a C_2-C_6 alkylcarbonyl group such as acetyl group, propionyl group, butyryl group, and valeryl group; amino group; a C_1-C_5 monoalkylamino group such as methylamino group, ethylamino group, propylamino group, isopropylamino group, butylamino group, isobutylamino group, tert-butylamino group, pentylamino group, and isopentylamino group; a C_2-C_{10} dialkylamino group such as dimethylamino group, ethylmethylamino group, diethylamino group, methylpropylamino group, and diisopropylamino group; a C_2-C_6 alkylcarbonylamino group such as acetylamino group, propionylamino group, isopropionylamino group, butyrylamino group, and valerylamino group; carbamoyl group; a C_2-C_6 alkylcarbamoyl group such as methylcarbamoyl group, ethylcarbamoyl group, propylcarbamoyl group, butylcarbamoyl group, tert-butylcarbamoyl group, and pentylcarbamoyl group; and a C_6-C_{12} aryl group such as phenyl group and naphthyl group.

Among the compounds of the present invention, preferred compounds include those wherein R^1 represents a hydrogen atom, R^8-CO-, R^8-O-CO-, R^8-NH-CO-, or R^8-SO$_2$- (R^8 represents a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of a C_3-C_8 cycloalkyl group, fluorenyl group, a C_6-C_{14} aryl group which may optionally be substituted, a C_6-C_{14} aryloxy group which may optionally be substituted, a C_6-C_{14} arylthio group which may optionally be substituted, a C_6-C_{14} arylsulfonyl group which may optionally be substituted, and a residue of a heterocyclic compound which may optionally be substituted; a C_3-C_8 cycloalkyl group; a C_6-C_{14} aryl group which may optionally be substituted; a C_2-C_5 alkenyl group which may optionally be substituted with an optionally substituted C_6-C_{14} aryl group; or a residue of a heterocyclic compound which may optionally be substituted); R^2, R^4, and R^6 independently represent a hydrogen atom, a C_1-C_5 alkyl group, or a C_2-C_6 alkanoyl group; R^3 and R^5 independently represent a hydrogen atom, a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of a C_6-C_{14} aryl group which may optionally be substituted and a C_1-C_5 alkoxy group; R^7 represents a hydrogen atom, a C_1-C_5 alkyl group, or R^9-CO- (R^9 represents a C_1-C_{10} alkyl group or a C_6-C_{12} aryl group which may optionally be substituted); symbol "A" represents a C_1-C_3 alkylene group which may optionally be substituted with a C_1-C_3 alkyl group; and symbol "n" represents 0 or 1. Among these compounds, those wherein R^2, R^4, and R^6 independently represent a hydrogen atom or a C_1-C_5 alkyl group are more preferred, and those wherein symbol "n" represents 0 are further preferred. Particularly preferred compounds include:

(1) compounds wherein R^1 represents R^8-CO- (R^8 represents a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of a C_6-C_{14} aryl group which may optionally be substituted, a C_6-C_{14} aryloxy group which may optionally be substituted, a C_6-C_{14} arylthio group which may optionally be substituted, and a C_6-C_{14} arylsulfonyl group which may optionally be substituted; a C_6-C_{14} aryl group which may optionally be substituted; a C_2-C_5 alkenyl group which may optionally be substituted with an optionally substituted C_6-C_{14} aryl group; or a residue of a heterocyclic compound which may optionally be substituted); R^2, R^4, and R^6 represent hydrogen atoms; R^3 and R^5 independently represent a C_1-C_{20} alkyl group; R^7 represents a hydrogen atom or R^9-CO- (R^9 represents a C_1-C_{10} alkyl group); symbol "A" repr sents a C_1-C_3 alkylene group; and symbol "n" represents 0;

(2) compounds wherein R^1 represents R^8-O-CO- (R^8 represents a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of a C_3-C_8 cycloalkyl group, fluorenyl group, a C_6-C_{14} aryl group which may optionally b substituted, and a residue of a h terocyclic compound which may optionally be substituted; a C_3-C_8 cycloalkyl group; or a C_6-C_{14} aryl group which may optionally be substituted); R^2, R^4, and R^6 represent hydrogen atoms; R^3 and R^5 independently represent a hydrogen atom, or a C_1-C_{20} alkyl group which may optionally be substituted with on or mor substituents selected from the group consisting of a C_6-C_{14} aryl group which may optionally be substituted and a C_1-C_5 alkoxy group; R^7 represents a hydrogen atom or R^9-CO- (R^9 represents a C_1-C_{10} alkyl group); symbol "A" represents a C_1-C_3 alkylene group; and symbol "n" represents 0;

(3) compounds wherein R^1 represents R^8-NH-CO- (R^8 represents a C_6-C_{14} aryl group which may optionally be substituted); R^2, R^4, and R^6 represent hydrogen atoms; R^3 and R^5 independently represent a C_1-C_{20} alkyl group; R^7 represents a hydrogen atom or R^9-CO- (R^9 represents a C_1-C_{10} alkyl group or a C_6-C_{12} aryl group which may optionally be substituted); symbol "A" represents a C_1-C_3 alkylene group; and symbol "n" represents 0; and

(4) compounds wherein R^1 represents R^8-SO_2- (R^8 represents a C_6-C_{14} aryl group which may optionally be substituted or a residue of a heterocyclic compound which may optionally be substituted); R^2, R^4, and R^6 represent hydrogen atoms; R^3 and R^5 independently represent a C_1-C_{20} alkyl group; R^7 represents a hydrogen atom, a C_1-C_5 alkyl group, or R^9-CO- (R^9 represents a C_1-C_{10} alkyl group or a C_6-C_{12} aryl group which may optionally be substituted); symbol "A" represents a C_1-C_3 alkylene group which may optionally be substituted with a C_1-C_3 alkyl group; and symbol "n" represents 0.

Among the compounds of the above definition (1), compounds wherein R^1 represents R^8-CO- (R^8 represents a C_1-C_{20} alkyl group which may optionally be substituted with an optionally substituted C_6-C_{14} aryl group or a C_6-C_{14} aryl group which may optionally be substituted); R^2, R^4, and R^6 represent hydrogen atoms; R^3 and R^5 independently represent a C_1-C_{20} alkyl group; R^7 represents R^9-CO- (R^9 represents a C_1-C_{10} alkyl group); symbol "A" represents a C_1-C_3 alkylene group; and symbol "n" represents 0 are more preferred.

Among the compounds of the above definition (2), compounds wherein R^1 represents R^8-O-CO- (R^8 represents a C_1-C_{20} alkyl group which may optionally be substituted with one or more substituents selected from the group consisting of fluorenyl group and a C_6-C_{14} aryl group which may optionally be substituted; or a C_3-C_8 cycloalkyl group); R^2, R^4, and R^6 represent hydrogen atoms; R^3 and R^5 independently represent a C_1-C_{20} alkyl group; R^7 represents R^9-CO- (R^9 represents a C_1-C_{10} alkyl group); symbol "A" represents a C_1-C_3 alkylene group; and symbol "n" represents 0 are more preferred.

Among the compounds of the above definition (3), compounds wherein R^7 represents a hydrogen atom are more preferred.

Among the compounds of the above definition (4), compounds wherein R^1 represents R^8-SO_2- (R^8 represents a C_6-C_{14} aryl group which may optionally be substituted); and R^7 represents a C_1-C_5 alkyl group, or compounds wherein R^1 represents R^8-SO_2- (R^8 represents a C_6-C_{14} aryl group which may optionally be substituted); and R^7 represents R^9-CO- (R^9 represents a C_6-C_{12} aryl group which may optionally be substituted) are more preferred.

The oxygen-containing heterocyclic derivatives of the present invention represented by the above formula (I) may form pharmaceutically acceptable salts. As specific examples of these salts, where acidic functional groups are attached, examples include metal salts such as lithium salt, sodium salt, potassium salt, magnesium salt, or calcium salt, and ammonium salts such as ammonium salt, methylammonium salt, dimethylammonium salt, trimethylammonium salt, or dicyclohexylammonium salt. Where basic functional groups are attached, examples include inorganic acid salts such as hydrochloride, hydrobromide, sulfate, nitrate, or phosphate, and organic acid salts such as methanesulfonate, benzenesulfonate, p-toluenesulfonate, acetate, propionate, tartrate, fumarate, maleate, malate, oxalate, succinate, citrate, benzoate, mandelate, cinnamate, or lactate. The oxygen-containing heterocyclic derivatives of the present invention represented by the above formula (I) may exist in the form of hydrates or solvates.

Referring to the stereochemistry of asymmetric carbon atoms exist in the oxygen-containing heterocyclic derivatives of the present invention represented by the above formula (I), they may independently be in (R)-, (S)-, or (RS)-configuration.

A compound represented by the following formula (II), which corresponds to the oxygen-containing heterocyclic derivatives of the present invention represented by the above formula (I) wherein R^7 is a hydrogen atom (R^1, R^2, R^3, R^4, R^5, R^6, A, and n have the same meanings as those defined above), may present under equilibrium between a hydroxyaldehyde derivative represented by the following formula (III) (R^1, R^2, R^3, R^4, R^5, R^6, A, and n have the same meanings as those defined above), especially in the state of a solution. This equilibrium can be demonstrated by the experimental results as explained below. The results of NMR measurements support the chemical structure of the formula (II), whilst differences in content ratio of the st reoisomers derived from the compound of the formula (II) were observed depending on types of solvents, which is attributable to the differences in content ratio of carbon atoms with different configurations that is bound with the hydroxyl group on the lactol ring. The differences in the cont nt ratio of the stereoisom rs can b explained by the presence of the equilibrium shown b low.

$$R^1 \left(\underset{R^2}{\overset{R^3}{N}} \cdot \underset{O}{\overset{R^4}{\underset{n}{||}}} \cdot N \cdot \underset{R^5}{\overset{O}{\underset{||}{C}}} \cdot \underset{R^6}{\overset{A}{N}} \cdot \underset{OH}{\overset{A}{O}} \right)_n \rightleftharpoons$$

(II)

$$R^1 \left(\underset{R^2}{\overset{R^3}{N}} \cdot \underset{O}{\overset{R^4}{\underset{n}{||}}} \cdot N \cdot \underset{R^5}{\overset{O}{\underset{||}{C}}} \cdot \underset{R^6}{\overset{A - OH}{N}} \cdot \underset{OH}{\overset{A - OH}{H}} \right)_n$$

(III)

Specific examples of the oxygen-containing heterocyclic derivative of the present invention represented by the above formula (I) include those shown in Table 1 below wherein n is 0 and R^7 is a hydrogen atom; those shown in Table 2 below wherein n is 0 and R^7 is other than a hydrogen atom; those shown in Table 3 wherein n is 1 and R^7 is a hydrogen atom; and those shown in Table 4 wherein n is 1 and R^7 is other than a hydrogen atom.

Table 1 (where n = 0)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
1	benzyl–O–C(=O)–CH$_3$	-H	-H	-H	oxetane–OH
2	benzyl–O–C(=O)–CH$_3$	-H	-CH$_3$	-H	oxetane–OH
3	benzyl–O–C(=O)–CH$_3$	-H	-CH$_2$CH$_2$CH$_3$	-H	oxetane–OH
4	benzyl–O–C(=O)–CH$_3$	-H	-CH(CH$_3$)$_2$	-H	oxetane–OH
5	benzyl–O–C(=O)–CH$_3$	-H	-CH$_2$CH$_2$CH$_2$CH$_3$	-H	oxetane–OH
6	H-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH
7	(H$_3$C)$_3$C–O–C(=O)–CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH

9

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	(A ring) with OH
8	benzyl–O–C(=O)–CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH
9	phenyl–CH$_2$CH$_2$–C(=O)–CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH
10	phenyl–O–CH$_2$–C(=O)–CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH
11	phenyl–C(=O)–CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH
12	phenyl–SO$_2$–	-H	-CH$_2$CH(CH$_3$)$_2$	-H	oxetane–OH
13	benzyl–O–C(=O)–CH$_3$	-H	-CH$_2$–phenyl	-H	oxetane–OH
14	benzyl–O–C(=O)–CH$_3$	-H	-CH$_2$CH$_2$SCH$_3$	-H	oxetane–OH

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
15	(benzyl ester)	-H	$-CH_2OH$	-H	(oxetane with OH)
16	(benzyl ester)	-H	(phenyl)	-H	(oxetane with OH)
17	(benzyl ester)	-H	-H	-H	(tetrahydrofuran with OH)
18	(benzyl ester)	-H	$-CH_3$	-H	(tetrahydrofuran with OH)
19	(benzyl ester)	-H	$-CH_2CH_3$	-H	(tetrahydrofuran with OH)
20	(benzyl ester)	-H	$-CH_2CH_2CH_3$	-H	(tetrahydrofuran with OH)
21	(benzyl ester)	-H	$-CH(CH_3)_2$	-H	(tetrahydrofuran with OH)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure: A, O, OH)
22	(benzyl –CH₂–O–C(=O)–CH₃)	$-H$	$-CH_2CH_2CH_2CH_3$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)
23	(benzyl –CH₂–O–C(=O)–CH₃)	$-H$	$-CH<\begin{array}{l}CH_3\\CH_2CH_3\end{array}$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)
24	$H-$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)
25	($H_3C-O-C(=O)-CH_3$)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)
26	($H_3C-CH_2-O-C(=O)-CH_3$)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)
27	($(H_3C)_3C-O-C(=O)-CH_3$)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)
28	($(H_3C)_2CH-CH_2-O-C(=O)-CH_3$)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃, O, OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
29	cyclohexyl-CH$_2$-O-C(=O)-CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	methyl-tetrahydrofuran-OH
30	cyclohexyl-CH$_2$CH$_2$-O-C(=O)-CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	methyl-tetrahydrofuran-OH
31	phenyl-CH$_2$-O-C(=O)-CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	methyl-tetrahydrofuran-OH
32	phenyl-CH$_2$-O-C(=O)-CH$_3$	-CH$_3$	-CH$_2$CH(CH$_3$)$_2$	-H	methyl-tetrahydrofuran-OH
33	phenyl-CH$_2$-O-C(=O)-CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$	methyl-tetrahydrofuran-OH
34	phenyl-CH$_2$-O-C(=O)-CH$_3$	-CH$_3$	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$	methyl-tetrahydrofuran-OH
35	2-fluoro-phenyl-CH$_2$-O-C(=O)-CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	methyl-tetrahydrofuran-OH

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(ring A / OH)
36	3-F-benzyl acetate (m-fluorobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)
37	4-F-benzyl acetate (p-fluorobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)
38	2-Cl-benzyl acetate (o-chlorobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)
39	3-Cl-benzyl acetate (m-chlorobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)
40	4-Cl-benzyl acetate (p-chlorobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)
41	2-Br-benzyl acetate (o-bromobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)
42	3-Br-benzyl acetate (m-bromobenzyl –O–C(=O)–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran ring with CH₃ and OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	
43		-H	$-CH_2CH(CH_3)_2$	-H	
44		-H	$-CH_2CH(CH_3)_2$	-H	
45		-H	$-CH_2CH(CH_3)_2$	-H	
46		-H	$-CH_2CH(CH_3)_2$	-H	
47		-H	$-CH_2CH(CH_3)_2$	-H	
48		-H	$-CH_2CH(CH_3)_2$	-H	
49		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
50	phenethyl acetate (PhCH₂CH₂–O–C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)
51	fluorenylmethyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)
52	(tetrahydrofuran-2-yl)methyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)
53	(tetrahydropyran-2-yl)methyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)
54	(pyridin-2-yl)methyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)
55	(pyridin-3-yl)methyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)
56	(pyridin-4-yl)methyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH₃ and OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
57	(pyridine N-oxide with CH₂-O-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)
58	(pyridine N-oxide with CH₂-O-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)
59	(pyridine N-oxide with CH₂-O-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)
60	(cyclohexyl-O-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)
61	(phenyl-O-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)
62	(benzyl-NH-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)
63	(phenyl-NH-C(=O)CH₃)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran with OH)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	
64		-H	$-CH_2CH(CH_3)_2$	-H	
65		-H	$-CH_2CH(CH_3)_2$	-H	
66		-H	$-CH_2CH(CH_3)_2$	-H	
67		-H	$-CH_2CH(CH_3)_2$	-H	
68		-H	$-CH_2CH(CH_3)_2$	-H	
69		-H	$-CH_2CH(CH_3)_2$	-H	
70		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	[ring]
71	H$_3$C—CH$_2$CH$_2$CH$_2$CH$_2$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]
72	(H$_3$C)$_2$CH—CH$_2$CH$_2$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]
73	(H$_3$C)$_3$C—CH$_2$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]
74	CH$_3$(CH$_2$)$_5$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]
75	CH$_3$(CH$_2$)$_6$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]
76	CH$_3$(CH$_2$)$_8$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]
77	CH$_3$(CH$_2$)$_{10}$—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	[tetrahydrofuran ring, OH]

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(A—O) OH
78	$CH_3(CH_2)_{12}$ O	–H	–$CH_2CH(CH_3)_2$	–H	
79	$CH_3(CH_2)_{14}$ O	–H	–$CH_2CH(CH_3)_2$	–H	
80	O	–H	–$CH_2CH(CH_3)_2$	–H	
81	O	–H	–$CH_2CH(CH_3)_2$	–H	
82	O	–H	–$CH_2CH(CH_3)_2$	–H	
83	O	–H	–$CH_2CH(CH_3)_2$	–H	
84	O	–H	–$CH_2CH(CH_3)_2$	–H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(ring A, O, OH structure)
85	(1-naphthyl)–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)
86	(2-naphthyl)–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)
87	H$_3$C–O–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)
88	(CH$_3$)$_3$C–O–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)
89	(cyclohexyl)–O–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)
90	(phenyl)–O–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)
91	(2-fluorophenyl)–O–CH$_2$COCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydrofuran ring with CH$_3$, O, OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	
92		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
93		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
94		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
95		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
96		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
97		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
98		-H	-CH$_2$CH(CH$_3$)$_2$	-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A)—O / OH
99	CH_3O-phenyl-O-CH_2-CO-CH_3 (4-methoxyphenoxyacetone)	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH
100	CH_3O-, OCH_3-phenyl-O-CH_2-CO-CH_3	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH
101	phenyl(CH_3)-O-CH_2-CO-CH_3	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH
102	H_3C-phenyl-O-CH_2-CO-CH_3	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH
103	H_3C-phenyl-O-CH_2-CO-CH_3	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH
104	phenyl(CF_3)-O-CH_2-CO-CH_3	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH
105	F_3C-phenyl-O-CH_2-CO-CH_3	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-CH₃, OH

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	ring structure with A, O, OH
106	F₃C-phenyl-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH
107	naphthyl-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH
108	naphthyl-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH
109	phenyl-CH₂-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH
110	phenyl-S-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH
111	phenyl-S(=O)₂-CH₂CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH
112	phenyl-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with CH₃ and OH

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	structure with A, O, OH
113		-H	$-CH_2CH(CH_3)_2$	-H	
114		-H	$-CH_2CH(CH_3)_2$	-H	
115		-H	$-CH_2CH(CH_3)_2$	-H	
116		-H	$-CH_2CH(CH_3)_2$	-H	
117		-H	$-CH_2CH(CH_3)_2$	-H	
118		-H	$-CH_2CH(CH_3)_2$	-H	
119		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A, O, OH ring structure)
120	3-Cl-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH
121	4-Cl-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH
122	2-Br-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH
123	3-Br-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH
124	4-Br-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH
125	2-CH_3-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH
126	3-H_3C-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran-OH

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	
127		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
128		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
129		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
130		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
131		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
132		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
133		-H	-CH$_2$CH(CH$_3$)$_2$	-H	

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	
134	H$_3$C— (4-acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)
135	(2-CF$_3$ acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)
136	F$_3$C— (3-acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)
137	F$_3$C— (4-acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)
138	CH$_3$O— (2-acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)
139	CH$_3$O— (3-acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)
140	CH$_3$O— (4-acetylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(ring structure)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A)—O ···OH
141	CH₃O / CH₃O / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)
142	OCH₃ / CH₃O / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)
143	CH₃O / CH₃O / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)
144	OCH₃ / CH₃O / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)
145	CH₃O / OCH₃ / CH₃O / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)
146	H₃C / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)
147	H₃C—O / O (acetophenone)	-H	-CH₂CH(CH₃)₂	-H	(ring)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A, O, OH ring)
148	H₃C—CH₂—O— benzene —C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH
149	(H₃C)₃C—O— benzene —C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH
150	HO— benzene —C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH
151	methylenedioxy benzene —C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH
152	methylenedioxy benzene —C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH
153	benzene with —C(=O)— and HO—C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH
154	HO—C(=O)— benzene —C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	tetrahydrofuran ring with methyl and OH

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A) O / OH structure
155	HO-CO-C₆H₄-CO- (4-carboxybenzoyl)	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH
156	1-naphthyl-CO-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH
157	2-naphthyl-CO-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH
158	2-pyridyl-CO-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH
159	3-pyridyl-CO-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH
160	4-pyridyl-CO-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH
161	2-furyl-CO-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran with CH₃ and OH

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
162	3-acetylfuran	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran
163	2-acetylthiophene	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran
164	3-acetylthiophene	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran
165	2-acetylpyrrole (N-H)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran
166	3-acetylpyrrole (N-H)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran
167	2-acetyl-1-methylpyrrole	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran
168	3-acetyl-1-methylpyrrole	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3-methyl-2-hydroxytetrahydrofuran

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
169	(pyrazol-5-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)
170	(3-ethyl-1-methyl-pyrazol-5-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)
171	(imidazol-5-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)
172	(1-methyl-imidazol-5-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)
173	(oxazol-5-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)
174	(thiazol-5-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)
175	(chroman-2-yl acetyl)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
176	chroman-3-yl acetyl	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH
177	4-oxo-4H-chromen-2-yl acetyl	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH
178	benzofuran-2-yl acetyl	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH
179	benzothiophen-2-yl acetyl	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH
180	indol-2-yl acetyl (N-H)	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH
181	H_3C—CH=CH—C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH
182	phenyl-CH=CH—C(=O)—	-H	$-CH_2CH(CH_3)_2$	-H	methyl-tetrahydrofuran-OH

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
183	F–C₆H₄–CH=CH–CO–CH₃ (4-fluorophenyl)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)
184	Cl–C₆H₄–CH=CH–CO–CH₃ (4-chlorophenyl)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)
185	H₃C–C₆H₄–CH=CH–CO–CH₃ (4-methylphenyl)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)
186	F₃C–C₆H₄–CH=CH–CO–CH₃ (4-trifluoromethylphenyl)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)
187	CH₃O–C₆H₄–CH=CH–CO–CH₃ (4-methoxyphenyl)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)
188	(CH₃O)₂–C₆H₃–CH=CH–CO–CH₃ (2,4-dimethoxyphenyl, OCH₃)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)
189	(CH₃O)₂–C₆H₃–CH=CH–CO–CH₃ (3,4-dimethoxyphenyl)	–H	–CH₂CH(CH₃)₂	–H	(tetrahydrofuran ring, OH)

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	
190		-H	$-CH_2CH(CH_3)_2$	-H	
191		-H	$-CH_2CH(CH_3)_2$	-H	
192		-H	$-CH_2CH(CH_3)_2$	-H	
193		-H	$-CH_2CH(CH_3)_2$	-H	
194		-H	$-CH_2CH(CH_3)_2$	-H	
195		-H	$-CH_2CH(CH_3)_2$	-H	
196		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	$\begin{smallmatrix} A \\ O \\ OH \end{smallmatrix}$
197	phenyl-SO$_2$-	-CH$_3$	-CH$_2$CH(CH$_3$)$_2$	-H	
198	phenyl-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$	
199	phenyl-SO$_2$-	-CH$_3$	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$	
200	2-F-phenyl-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
201	3-F-phenyl-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
202	4-F-phenyl-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
203	2-Cl-phenyl-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A ring with O, OH)
204	3-Cl-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)
205	4-Cl-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)
206	2-Br-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)
207	3-Br-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)
208	4-Br-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)
209	2-CH₃-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)
210	3-CH₃-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran ring (methyl, OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
211	H_3C—〈benzene〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)
212	H_3C—〈benzene, CH_3〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)
213	〈benzene, CH_3, CH_3〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)
214	H_3C, H_3C—〈benzene〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)
215	H_3C—〈benzene, CH_3, CH_3〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)
216	H_3C—CH_2—〈benzene〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)
217	H_3C, H_3C—CH—〈benzene〉—SO_2—	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring, CH₃, OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(A-O-OH ring)
218	(H$_3$C)$_3$C—C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
219	2-OCH$_3$-C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
220	3-CH$_3$O-C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
221	CH$_3$O—C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
222	H$_3$C CH$_2$—O—C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
223	(H$_3$C)$_2$CH—O—C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
224	(H$_3$C)$_3$C—O—C$_6$H$_4$—SO$_2$—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(ring with A, O, OH)
225	HO—C₆H₄—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)
226	o-O₂N-C₆H₄—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)
227	m-O₂N-C₆H₄—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)
228	p-O₂N-C₆H₄—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)
229	1-naphthyl—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)
230	2-naphthyl—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)
231	2-pyridyl—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(tetrahydrofuran, methyl, OH)

41

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	$\overset{A}{\diagup}\diagdown O$ / OH
232	pyridin-3-yl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)
233	pyridin-4-yl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)
234	furan-2-yl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)
235	thiophen-2-yl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)
236	pyrrol-2-yl-SO_2- (N-H)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)
237	pyrrol-2-yl-SO_2- (N-CH_3)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)
238	oxazol-2-yl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydrofuran structure with OH)

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	
239		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
240		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
241	H-	-H	-CH$_2$⟨phenyl⟩	-H	
242	H$_3$C-O-C(=O)-	-H	-CH$_2$⟨phenyl⟩	-H	
243	(H$_3$C)$_3$C-O-C(=O)-	-H	-CH$_2$⟨phenyl⟩	-H	
244	(H$_3$C)$_2$CHCH$_2$-O-C(=O)-	-H	-CH$_2$⟨phenyl⟩	-H	
245	cyclohexyl-CH$_2$-O-C(=O)-	-H	-CH$_2$⟨phenyl⟩	-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
246		-H		-H	
247		-H		-H	
248		-H		-H	
249		-H		-H	
250		-H		-H	
251		-H		-H	
252		-H		-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
253	(benzyl acetate with OCH₃)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)
254	(CH₃O-benzyl acetate)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)
255	(N-phenyl acetamide, H, N, O)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)
256	(H₃C, H₃C, O)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)
257	(phenyl, O)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)
258	(phenoxy, O)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)
259	(phenoxy, Cl, O)	-H	-CH₂-(phenyl)	-H	(tetrahydrofuran ring, OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(A/O/OH ring)
260	4-Cl-C$_6$H$_4$-O-CH$_2$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)
261	2-CH$_3$-C$_6$H$_4$-O-CH$_2$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)
262	4-CH$_3$-C$_6$H$_4$-O-CH$_2$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)
263	2-OCH$_3$-C$_6$H$_4$-O-CH$_2$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)
264	4-CH$_3$O-C$_6$H$_4$-O-CH$_2$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)
265	C$_6$H$_5$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)
266	2-F-C$_6$H$_4$-CO-CH$_3$	-H	-CH$_2$-C$_6$H$_5$	-H	(tetrahydrofuran-OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	structure with A, O, OH
267		-H	-CH₂—	-H	
268		-H	-CH₂—	-H	
269		-H	⁼CH₂—	-H	
270		-H	-CH₂—	-H	
271		-H	-CH₂—	-H	
272		-H	-CH₂—	-H	
273		-H	-CH₂—	-H	

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	ring structure
274	H₃C–⬡–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH
275	⬡(CH₃O)–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH
276	⬡(CH₃O)–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH
277	CH₃O–⬡–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH
278	pyridin-2-yl–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH
279	pyridin-3-yl–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH
280	pyridin-4-yl–C(=O)–	-H	-CH₂–⬡	-H	tetrahydrofuran-OH

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(ring A structure with OH)
281	H$_3$C—S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH
282	(H$_3$C)$_2$CH—S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH
283	C$_6$H$_5$-CH$_2$-S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH
284	C$_6$H$_5$-S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH
285	F-C$_6$H$_4$-S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH
286	Cl-C$_6$H$_4$-S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH
287	Br-C$_6$H$_4$-S(=O)$_2$—	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydrofuran ring with OH

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
288		-H		-H	
289		-H		-H	
290		-H		-H	
291		-H		-H	
292		-H		-H	
293		-H		-H	
294		-H		-H	

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
295	(benzyl acetate structure)	-H	$-CH_2CH_2SCH_3$	-H	(tetrahydrofuran ring with OH)
296	(phenyl sulfonyl structure)	-H	$-CH_2CH_2SCH_3$	-H	(tetrahydrofuran ring with OH)
297	(benzyl acetate structure)	-H	$-CH_2OC(CH_3)_3$	-H	(tetrahydrofuran ring with OH)
298	(phenyl sulfonyl structure)	-H	$-CH_2OC(CH_3)_3$	-H	(tetrahydrofuran ring with OH)
299	(benzyl acetate structure)	-H	$-CH_2OCH_2$—(phenyl)	-H	(tetrahydrofuran ring with OH)
300	(phenyl sulfonyl structure)	-H	$-CH_2OCH_2$—(phenyl)	-H	(tetrahydrofuran ring with OH)
301	(benzyl acetate structure)	-H	$-CH_2OH$	-H	(tetrahydrofuran ring with OH)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
302		-H	-CH₂OH	-H	
303		-H	-H	-H	
304		-H	-H	-H	
305		-H	-CH₃	-H	
306		-H	-CH₃	-H	
307		-H	-CH₂CH₂CH₃	-H	
308		-H	-CH₂CH₂CH₃	-H	

Table 1   (Continued)

Compound No.	R[1]	R[4]	R[5]	R[6]	
309		-H	$-CH(CH_3)_2$	-H	
310		-H	$-CH(CH_3)_2$	-H	
311		-H	$-CH_2CH_2CH_2CH_3$	-H	
312		-H	$-CH_2CH_2CH_2CH_3$	-H	
313	H-	-H	$-CH_2CH(CH_3)_2$	-H	
314		-H	$-CH_2CH(CH_3)_2$	-H	
315		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	structure
316	(phenyl-CH2CH2-C(=O)-CH3)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH3 and OH)
317	(phenyl-O-CH2-C(=O)-CH3)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH3 and OH)
318	(phenyl-C(=O)-CH3)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH3 and OH)
319	(fluorophenyl-C(=O)-CH3)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH3 and OH)
320	(phenyl-SO2-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran ring with CH3 and OH)
321	(benzyl-O-C(=O)-CH3)	-H	$-CH_2-$phenyl	-H	(tetrahydrofuran ring with CH3 and OH)
322	(phenyl-SO2-)	-H	$-CH_2-$phenyl	-H	(tetrahydrofuran ring with CH3 and OH)

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	
323		-H	-CH$_2$CH$_2$SCH$_3$	-H	
324		-H	-CH$_2$CH$_2$SCH$_3$	-H	
325		-H	-CH$_2$OC(CH$_3$)$_3$	-H	
326		-H	-CH$_2$OC(CH$_3$)$_3$	-H	
327		-H	-CH$_2$OH	-H	
328		-H	-CH$_2$OH	-H	
329		-H		-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	
330		-H		-H	
331		-H	-H	-H	
332		-H	-H	-H	
333		-H	$-CH_3$	-H	
334		-H	$-CH_3$	-H	
335		-H	$-CH_2CH_2CH_3$	-H	
336		-H	$-CH_2CH_2CH_3$	-H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
337	benzyl acetate	-H	-CH(CH$_3$)$_2$	-H	
338	phenyl sulfonyl	-H	-CH(CH$_3$)$_2$	-H	
339	benzyl acetate	-H	-CH$_2$CH$_2$CH$_2$CH$_3$	-H	
340	phenyl sulfonyl	-H	-CH$_2$CH$_2$CH$_2$CH$_3$	-H	
341	H-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	
342		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
343	benzyl acetate	-H	-CH$_2$CH(CH$_3$)$_2$	-H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
344	(4-fluorobenzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)
345	(4-chlorobenzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)
346	(4-bromobenzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)
347	(4-methylbenzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)
348	(4-methoxybenzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)
349	(acetone structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)
350	(4-methyl-2-pentanone structure)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure with CH$_3$, H$_3$C, O, OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure)
351		-H	$-CH_2CH(CH_3)_2$	-H	
352		-H	$-CH_2CH(CH_3)_2$	-H	
353		-H	$-CH_2CH(CH_3)_2$	-H	
354		-H	$-CH_2CH(CH_3)_2$	-H	
355		-H	$-CH_2CH(CH_3)_2$	-H	
356		-H	$-CH_2CH(CH_3)_2$	-H	
357		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A)O OH
358	CH₃O—⟨⟩—O—CH₂—C(=O) with OCH₃	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH
359	⟨⟩—C(=O)	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH
360	⟨⟩(F)—C(=O)	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH
361	F—⟨⟩—C(=O)	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH
362	F—⟨⟩—C(=O)	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH
363	⟨⟩(Cl)—C(=O)	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH
364	Cl—⟨⟩—C(=O)	-H	$-CH_2CH(CH_3)_2$	-H	H₃C—CH₃ ring, O, OH

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	
365		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
366		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
367		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
368		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
369		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
370		-H	-CH$_2$CH(CH$_3$)$_2$	-H	
371		-H	-CH$_2$CH(CH$_3$)$_2$	-H	

61

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
372	3-pyridyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)
373	4-pyridyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)
374	2-furyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)
375	2-thienyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)
376	$H_3C-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)
377	$(H_3C)_2CH-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)
378	phenyl-$S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran structure)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(ring structure with A, O, OH)
379	F—⟨ ⟩—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)
380	Cl—⟨ ⟩—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)
381	Br—⟨ ⟩—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)
382	H₃C—⟨ ⟩—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)
383	H₃C—⟨ ⟩(CH₃)(CH₃)—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)
384	(H₃C)(H₃C)(H₃C)C—⟨ ⟩—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)
385	CH₃O—⟨ ⟩—S(=O)(=O)—	-H	-CH₂CH(CH₃)₂	-H	(furan ring with CH₃, H₃C, OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A)—O / —OH ring
386	3-O₂N-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran (CH₃, H₃C, O, OH)
387	pyridin-2-yl-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran (CH₃, H₃C, O, OH)
388	pyridin-3-yl-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran (CH₃, H₃C, O, OH)
389	pyridin-4-yl-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran (CH₃, H₃C, O, OH)
390	furan-2-yl-SO₂-	-H	-CH₂CH(CH₃)₂	-H	tetrahydrofuran (CH₃, H₃C, O, OH)
391	H-	-H	-CH₂-C₆H₅	-H	tetrahydrofuran (CH₃, H₃C, O, OH)
392	(CH₃)₃C-O-C(=O)-	-H	-CH₂-C₆H₅	-H	tetrahydrofuran (CH₃, H₃C, O, OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
393	(benzyl acetate structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)
394	(F-phenyl-CH₂-O-acetate structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)
395	(Cl-phenyl-CH₂-O-acetate structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)
396	(H₃C-phenyl-CH₂-O-acetate structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)
397	(CH₃O-phenyl-CH₂-O-acetate structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)
398	(phenyl-CH₂CH₂-CO-CH₃ structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)
399	(phenyl-O-CH₂-CO-CH₃ structure)	-H	-CH₂-(phenyl)	-H	(H₃C, CH₃, tetrahydrofuran-OH structure)

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	(A ring / OH group)
400	CH_3O-(4-position on phenyl)-O-CH_2-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH
401	CH_3O- and OCH_3 substituted phenyl-O-CH_2-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH
402	phenyl-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH
403	(2-F)phenyl-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH
404	(2,4-di-F)phenyl-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH
405	(2,6-di-F)phenyl-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH
406	(F_3C-, F-)phenyl-C(=O)-CH$_3$	-H	-CH_2-phenyl	-H	tetrahydrofuran ring, H_3C- and CH_3 substituents, OH

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
407	H_3C-SO_2-	-H	$-CH_2$-phenyl	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)
408	phenyl-SO_2-	-H	$-CH_2$-phenyl	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)
409	F-phenyl-SO_2-	-H	$-CH_2$-phenyl	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)
410	CH_3O-phenyl-SO_2-	-H	$-CH_2$-phenyl	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)
411	phenyl-CH_2-O-C(=O)-CH$_3$	-H	$-CH_2$-naphthyl	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)
412	phenyl-SO_2-	-H	$-CH_2$-naphthyl	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)
413	phenyl-CH_2-O-C(=O)-CH$_3$	-H	$-CH_2CH_2SCH_3$	-H	(tetrahydrofuran, CH$_3$, H$_3$C, OH)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
414	phenyl-SO₂–	-H	-CH₂CH₂SCH₃	-H	(tetrahydrofuran with CH₃, H₃C, OH)
415	benzyl-O-C(=O)-	-H	-CH₂OC(CH₃)₃	-H	(tetrahydrofuran with CH₃, H₃C, OH)
416	phenyl-SO₂–	-H	-CH₂OC(CH₃)₃	-H	(tetrahydrofuran with CH₃, H₃C, OH)
417	benzyl-O-C(=O)-	-H	-CH₂OH	-H	(tetrahydrofuran with CH₃, H₃C, OH)
418	phenyl-SO₂–	-H	-CH₂OH	-H	(tetrahydrofuran with CH₃, H₃C, OH)
419	benzyl-O-C(=O)-	-H	phenyl	-H	(tetrahydrofuran with CH₃, H₃C, OH)
420	phenyl-SO₂–	-H	phenyl	-H	(tetrahydrofuran with CH₃, H₃C, OH)

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
421	(benzyl ester structure)	-H	-H	-H	(tetrahydropyran with OH)
422	(phenyl sulfonyl structure)	-H	-H	-H	(tetrahydropyran with OH)
423	(benzyl ester structure)	-H	$-CH_3$	-H	(tetrahydropyran with OH)
424	(phenyl sulfonyl structure)	-H	$-CH_3$	-H	(tetrahydropyran with OH)
425	(benzyl ester structure)	-H	$-CH_2CH_2CH_3$	-H	(tetrahydropyran with OH)
426	(phenyl sulfonyl structure)	-H	$-CH_2CH_2CH_3$	-H	(tetrahydropyran with OH)
427	(benzyl ester structure)	-H	$-CH(CH_3)_2$	-H	(tetrahydropyran with OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
428	(phenyl-SO₂-)	-H	-CH(CH₃)₂	-H	(tetrahydropyran with OH, CH₃)
429	(benzyl-O-C(=O)-CH₃)	-H	-CH₂CH₂CH₂CH₃	-H	(tetrahydropyran with OH, CH₃)
430	(phenyl-SO-)	-H	-CH₂CH₂CH₂CH₃	-H	(tetrahydropyran with OH, CH₃)
431	-H	-H	-CH₂CH(CH₃)₂	-H	(tetrahydropyran with OH, CH₃)
432	(H₃C)₃C-O-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	(tetrahydropyran with OH, CH₃)
433	(benzyl-O-C(=O)-CH₃)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydropyran with OH, CH₃)
434	(F-phenyl-CH₂-O-C(=O)-CH₃)	-H	-CH₂CH(CH₃)₂	-H	(tetrahydropyran with OH, CH₃)

70

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(structure with A, O, OH)
435	Cl—(C$_6$H$_4$)—CH$_2$O—C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)
436	Br—(C$_6$H$_4$)—CH$_2$O—C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)
437	H$_3$C—(C$_6$H$_4$)—CH$_2$O—C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)
438	CH$_3$O—(C$_6$H$_4$)—CH$_2$O—C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)
439	H$_3$C—C(=O)—	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)
440	(H$_3$C)$_2$CH—C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)
441	(C$_6$H$_5$)—CH$_2$—C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(tetrahydropyran with OH)

71

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure with A, O, OH)
442	phenyl-CH₂CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)
443	phenyl-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)
444	F-C₆H₄-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)
445	Cl-C₆H₄-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)
446	H₃C-C₆H₄-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)
447	CH₃O-C₆H₄-O-CH₂-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)
448	phenyl-NH-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran with CH₃ and OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(structure)
449	(phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
450	(2-F-phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
451	(3-F-phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
452	(4-F-phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
453	(2-Cl-phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
454	(4-Cl-phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
455	(2-CH₃-phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	
456		-H	$-CH_2CH(CH_3)_2$	-H	
457		-H	$-CH_2CH(CH_3)_2$	-H	
458		-H	$-CH_2CH(CH_3)_2$	-H	
459		-H	$-CH_2CH(CH_3)_2$	-H	
460		-H	$-CH_2CH(CH_3)_2$	-H	
461		-H	$-CH_2CH(CH_3)_2$	-H	
462		-H	$-CH_2CH(CH_3)_2$	-H	

Table 1   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A/O structure) OH
463	(pyridyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
464	(furyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
465	(thienyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
466	$H_3C-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
467	$(CH_3)_2CH-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
468	(phenyl-$S(=O)_2-$)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)
469	$F-$(phenyl)$-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH)

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(ring with A, O, OH)
470	Cl—C₆H₄—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH
471	Br—C₆H₄—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH
472	H₃C—C₆H₄—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH
473	H₃C—C₆H₂(CH₃)₂—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH
474	(H₃C)₃C—C₆H₄—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH
475	CH₃O—C₆H₄—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH
476	O₂N—C₆H₄—SO₂—	-H	-CH₂CH(CH₃)₂	-H	tetrahydropyranyl-CH₃, OH

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	
477		-H	$-CH_2CH(CH_3)_2$	-H	
478		-H	$-CH_2CH(CH_3)_2$	-H	
479		-H	$-CH_2CH(CH_3)_2$	-H	
480		-H	$-CH_2CH(CH_3)_2$	-H	
481	H-	-H		-H	
482		-H		-H	
483		-H		-H	

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	
484		-H	-CH₂-⬡	-H	
485		-H	-CH₂-⬡	-H	
486		-H	-CH₂-⬡	-H	
487		-H	-CH₂-⬡	-H	
488		-H	-CH₂-⬡	-H	
489		-H	-CH₂-⬡	-H	
490		-H	-CH₂-⬡	-H	

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(ring with A, O, OH)
491	CH$_3$O-, OCH$_3$-substituted phenyl-O-CH$_2$-C(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH
492	C$_6$H$_5$-C(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH
493	F-substituted phenyl-C(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH
494	F-substituted phenyl-C(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH
495	F,F-disubstituted phenyl-C(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH
496	F$_3$C-substituted phenyl-C(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH
497	H$_3$C-S(=O)(=O)-	-H	-CH$_2$-C$_6$H$_5$	-H	tetrahydropyran ring with CH$_3$ and OH

Table 1 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	(A)—O with OH
498		-H		-H	
499		-H		-H	
500		-H		-H	
501		-H		-H	
502		-H		-H	
503		-H	$-CH_2CH_2SCH_3$	-H	
504		-H	$-CH_2CH_2SCH_3$	-H	

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	![A ring structure]
505	![benzyl acetate group]	-H	-CH$_2$OC(CH$_3$)$_3$	-H	![tetrahydropyran-OH ring]
506	![phenylsulfonyl group]	-H	-CH$_2$OC(CH$_3$)$_3$	-H	![tetrahydropyran-OH ring]
507	![benzyl acetate group]	-H	-CH$_2$OH	-H	![tetrahydropyran-OH ring]
508	![phenylsulfonyl group]	-H	-CH$_2$OH	-H	![tetrahydropyran-OH ring]
509	![benzyl acetate group]	-H	![phenyl group]	-H	![tetrahydropyran-OH ring]
510	![phenylsulfonyl group]	-H	![phenyl group]	-H	![tetrahydropyran-OH ring]
511	![benzyl acetate group]	-H	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$CO	![tetrahydrofuran-OH ring]

Table 1 (Continued)

Compound No.	R^1	R^4	R^5	R^6	(ring with A, O, OH)
512	benzyl—O—C(=O)— (PhCH$_2$OC(O)—)	CH$_3$CO—	—CH$_2$CH(CH$_3$)$_2$	—H	tetrahydrofuran ring, CH$_3$, OH
513	C$_6$H$_5$—S(=O)$_2$—	CH$_3$CO—	—CH$_2$CH(CH$_3$)$_2$	—H	tetrahydrofuran ring, CH$_3$, OH
514	F—C$_6$H$_4$—S(=O)$_2$—	CH$_3$CO—	—CH$_2$CH(CH$_3$)$_2$	—H	tetrahydrofuran ring, CH$_3$, OH
515	H$_3$C—C$_6$H$_4$—S(=O)$_2$—	CH$_3$CO—	—CH$_2$CH(CH$_3$)$_2$	—H	tetrahydrofuran ring, CH$_3$, OH
516	CH$_3$O—C$_6$H$_4$—S(=O)$_2$—	CH$_3$CO—	—CH$_2$CH(CH$_3$)$_2$	—H	tetrahydrofuran ring, CH$_3$, OH
517	benzyl—O—C(=O)— (PhCH$_2$OC(O)—)	CH$_3$CO—	—CH$_2$—C$_6$H$_5$	—H	tetrahydrofuran ring, CH$_3$, OH
518	C$_6$H$_5$—S(=O)$_2$—	CH$_3$CO—	—CH$_2$—C$_6$H$_5$	—H	tetrahydrofuran ring, CH$_3$, OH

Table 1   (Continued)

Compound No.	R^1	R^4	R^5	R^6	
519		CH_3CO-	$-CH_2CH(CH_3)_2$	$-H$	
520		CH_3CO-	$-CH_2CH(CH_3)_2$	$-H$	

Table 2 (where n = 0)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A
521		-H	-H	-H		
522		-H	-CH₃	-H		
523		-H	-CH₂CH₂CH₃	-H		
524		-H	-CH(CH₃)₂	-H		

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A) O O-
525	benzyl-O-C(=O)-CH₃	-H	-CH₂CH₂CH₂CH₃	-H	CH₃-C(=O)-	oxetane ring-O-
526	H-	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	oxetane ring-O-
527	(CH₃)₃C-O-C(=O)-CH₃ (H₃C/H₃C/H₃C)	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	oxetane ring-O-
528	benzyl-O-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	oxetane ring-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
529	(phenyl-CH₂CH₂-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(oxetane structure with O-)
530	(phenyl-O-CH₂-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(oxetane structure with O-)
531	(phenyl-C(=O)-)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(oxetane structure with O-)
532	(phenyl-S(=O)₂-)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(oxetane structure with O-)

86

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
533	CH₃–C(=O)–O–CH₂–C₆H₅	–H	–CH₂–C₆H₅	–H	CH₃–C(=O)–	(oxetane)–O–
534	CH₃–C(=O)–O–CH₂–C₆H₅	–H	–CH₂CH₂SCH₃	–H	CH₃–C(=O)–	(oxetane)–O–
535	CH₃–C(=O)–O–CH₂–C₆H₅	–H	–CH₂OH	–H	CH₃–C(=O)–	(oxetane)–O–
536	CH₃–C(=O)–O–CH₂–C₆H₅	–H	C₆H₅–	–H	CH₃–C(=O)–	(oxetane)–O–

Table 2 (Continued)

(structure: A–O, O⁻)	R⁷	R⁶	R⁵	R⁴	R¹	Compound No.
(tetrahydropyran, O⁻, CH₃)	CH₃–C(=O)–	–H	–H	–H	(benzyl acetate-type)	537
(tetrahydropyran, O⁻, CH₃)	CH₃–C(=O)–	–H	–CH₃	–H	(benzyl acetate-type)	538
(tetrahydropyran, O⁻, CH₃)	CH₃–C(=O)–	–H	–CH₃	–H	(O=S=O phenyl)	539
(tetrahydropyran, O⁻, CH₃)	CH₃–C(=O)–	–H	–CH₂CH₂CH₃	–H	(benzyl acetate-type)	540

88

Table 2   (Continued)

EP 0 810 221 A1 placed at top as header.

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
541		-H	-CH(CH₃)₂	-H		
542		-H	-CH₂CH₂CH₂CH₃	-H		
543		-H	$-CH(CH_3)(CH_2CH_3)$	-H		
544	H-	-H	-CH₂CH(CH₃)₂	-H		

Column headers (as printed): Compound No., R¹, R⁴, R⁵, R⁶, R⁷

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
545	H_3C–O–C(=O) (acetoxymethyl)	-H	-$CH_2CH(CH_3)_2$	-H	CH_3–C(=O) (acetyl)	(tetrahydropyran-2-yl)oxy with methyl
546	H_3C–CH$_2$–O–C(=O)	-H	-$CH_2CH(CH_3)_2$	-H	CH_3–C(=O)	(tetrahydropyran-2-yl)oxy with methyl
547	H_3C, H_3C, H_3C–C–O–C(=O)	-H	-$CH_2CH(CH_3)_2$	-H	CH_3–C(=O)	(tetrahydropyran-2-yl)oxy with methyl
548	H_3C, H_3C–CH–CH$_2$–O–C(=O)	-H	-$CH_2CH(CH_3)_2$	-H	CH_3–C(=O)	(tetrahydropyran-2-yl)oxy with methyl

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
549	cyclohexylmethyl–O–C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	tetrahydropyran-2-yl-O– (A, methyl)
550	2-cyclohexylethyl–O–C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	3-methyltetrahydropyran-2-yl-O–
551	benzyl–O–C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	3-methyltetrahydropyran-2-yl-O–
552	benzyl–O–C(=O)CH$_3$	-CH$_3$	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	3-methyltetrahydropyran-2-yl-O–

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
553	benzyl-CH$_2$-O-C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$	CH$_3$-C(=O)-	tetrahydropyranyl-methyl-O-
554	benzyl-CH$_2$-O-C(=O)CH$_3$	-CH$_3$	-CH$_2$CH(CH$_3$)$_2$	-CH$_3$	CH$_3$-C(=O)-	tetrahydropyranyl-methyl-O-
555	(2-F-benzyl)-CH$_2$-O-C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	tetrahydropyranyl-methyl-O-
556	(3-F-benzyl)-CH$_2$-O-C(=O)CH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	tetrahydropyranyl-methyl-O-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
557	$4\text{-F-C}_6\text{H}_4\text{-CH}_2\text{-O-C(=O)-CH}_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3\text{-C(=O)-}$	(A)-methyl-tetrahydropyran-O-
558	$2\text{-Cl-C}_6\text{H}_4\text{-CH}_2\text{-O-C(=O)-CH}_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3\text{-C(=O)-}$	methyl-tetrahydropyran-O-
559	$3\text{-Cl-C}_6\text{H}_4\text{-CH}_2\text{-O-C(=O)-CH}_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3\text{-C(=O)-}$	methyl-tetrahydropyran-O-
560	$4\text{-Cl-C}_6\text{H}_4\text{-CH}_2\text{-O-C(=O)-CH}_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3\text{-C(=O)-}$	methyl-tetrahydropyran-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
561	2-Br-benzyl-O-C(=O)CH$_3$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH$_3$C(=O)-	tetrahydropyran-2-yl-O⁻, 3-methyl (A)
562	3-Br-benzyl-O-C(=O)CH$_3$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH$_3$C(=O)-	tetrahydropyran-2-yl-O⁻, 3-methyl
563	4-Br-benzyl-O-C(=O)CH$_3$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH$_3$C(=O)-	tetrahydropyran-2-yl-O⁻, 3-methyl
564	2-CH$_3$-benzyl-O-C(=O)CH$_3$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH$_3$C(=O)-	tetrahydropyran-2-yl-O⁻, 3-methyl

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A
565	(H₃C-benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	(acetyl) CH₃ C=O	(tetrahydropyran ring with methyl, O-)
566	(H₃C-benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	(acetyl) CH₃ C=O	(tetrahydropyran ring with methyl, O-)
567	(OCH₃-benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	(acetyl) CH₃ C=O	(tetrahydropyran ring with methyl, O-)
568	(CH₃O-benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	(acetyl) CH₃ C=O	(tetrahydropyran ring with methyl, O-)

95

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
569	CH₃O–⟨benzyl⟩–CH₂–O–C(=O)–CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	⟨A, tetrahydropyran, CH₃, O–⟩
570	⟨phenyl⟩–CH₂CH₂–O–C(=O)–CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	⟨tetrahydropyran, CH₃, O–⟩
571	⟨fluorenyl⟩–CH₂–O–C(=O)–CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	⟨tetrahydropyran, CH₃, O–⟩
572	⟨tetrahydrofuryl⟩–CH₂–O–C(=O)–CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	⟨tetrahydropyran, CH₃, O–⟩

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A ring)
573	(tetrahydropyranylmethyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(acetyl, CH_3-C=O)	(tetrahydropyran, methyl substituted)
574	(pyridin-2-ylmethyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(acetyl, CH_3-C=O)	(tetrahydropyran, methyl substituted)
575	(pyridin-3-ylmethyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(acetyl, CH_3-C=O)	(tetrahydropyran, methyl substituted)
576	(pyridin-4-ylmethyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	(acetyl, CH_3-C=O)	(tetrahydropyran, methyl substituted)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A ring)
577	(2-pyridylmethyl acetate N-oxide)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-CO-	(3-methyltetrahydropyran-2-yl-O-)
578	(3-pyridylmethyl acetate N-oxide)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-CO-	(3-methyltetrahydropyran-2-yl-O-)
579	(4-pyridylmethyl acetate N-oxide)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-CO-	(3-methyltetrahydropyran-2-yl-O-)
580	(cyclohexyl acetate)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-CO-	(3-methyltetrahydropyran-2-yl-O-)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
581	(phenyl ester)	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-methyl-O-)
582	(benzyl amide, H-N)	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-methyl-O-)
583	(phenyl amide, H-N)	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-methyl-O-)
584	H₃C-C(=O)-	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-methyl-O-)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
585	$CH_3CH_2C(=O)CH_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	methyl-tetrahydropyran-2-yl-O^-
586	$CH_3CH_2CH_2C(=O)CH_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	methyl-tetrahydropyran-2-yl-O^-
587	$(CH_3)(CH_3CH_2)CHC(=O)-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	methyl-tetrahydropyran-2-yl-O^-
588	$CH_3CH_2CH_2CH_2C(=O)CH_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	methyl-tetrahydropyran-2-yl-O^-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
589	(H$_3$C)(H$_3$C)CH–CH$_2$–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	ring structure (tetrahydropyran with A and methyl, –O–)
590	(H$_3$C)(H$_3$C)(H$_3$C)C–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	3-methyltetrahydropyran–2–O–
591	H$_3$C–CH$_2$CH$_2$CH$_2$–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	3-methyltetrahydropyran–2–O–
592	(H$_3$C)(H$_3$C)CH–CH$_2$CH$_2$–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	3-methyltetrahydropyran–2–O–

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
593	(CH₃)₃C–CH₂–C(=O)–	-H	-CH₂CH(CH₃)₂	-H	CH₃–C(=O)–	tetrahydropyran ring with methyl, -O-
594	CH₃(CH₂)₅–C(=O)–	-H	-CH₂CH(CH₃)₂	-H	CH₃–C(=O)–	tetrahydropyran ring with methyl, -O-
595	CH₃(CH₂)₆–C(=O)–	-H	-CH₂CH(CH₃)₂	-H	CH₃–C(=O)–	tetrahydropyran ring with methyl, -O-
596	CH₃(CH₂)₉–C(=O)–	-H	-CH₂CH(CH₃)₂	-H	CH₃–C(=O)–	tetrahydropyran ring with methyl, -O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(ring structure)
597	$CH_3(CH_2)_{10}C(=O)-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	tetrahydropyran-methyl-O-
598	$CH_3(CH_2)_{12}C(=O)-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	tetrahydropyran-methyl-O-
599	$CH_3(CH_2)_{14}C(=O)-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	tetrahydropyran-methyl-O-
600	cyclohexyl-$CH_2C(=O)-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	tetrahydropyran-methyl-O-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
601	(3-cyclohexyl-propan-2-one)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(3-methyltetrahydropyran-2-yl)-O-
602	(1-phenyl-propan-2-one)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(3-methyltetrahydropyran-2-yl)-O-
603	(4-phenyl-butan-2-one)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(3-methyltetrahydropyran-2-yl)-O-
604	(5-phenyl-pentan-2-one)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(3-methyltetrahydropyran-2-yl)-O-

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	$\left(\!A\!\right)$
605	(1-naphthyl-CH₂-CO-)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyran-2-yl, 3-CH₃)
606	(2-naphthyl-CH₂-CO-)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyran-2-yl, 3-CH₃)
607	$H_3C-O-CH_2-CO-$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyran-2-yl, 3-CH₃)
608	($(CH_3)_3C-O-CH_2-CO-$)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyran-2-yl, 3-CH₃)

105

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A—O—O⁻
609	(cyclohexyl)—O—CH2—C(=O)—CH3	-H	$-CH_2CH(CH_3)_2$	-H	CH3—C(=O)—	(tetrahydropyran with CH3)—O⁻
610	(phenyl)—O—CH2—C(=O)—CH3	-H	$-CH_2CH(CH_3)_2$	-H	CH3—C(=O)—	(tetrahydropyran with CH3)—O⁻
611	(2-fluorophenyl)—O—CH2—C(=O)—CH3	-H	$-CH_2CH(CH_3)_2$	-H	CH3—C(=O)—	(tetrahydropyran with CH3)—O⁻
612	(3-fluorophenyl)—O—CH2—C(=O)—CH3	-H	$-CH_2CH(CH_3)_2$	-H	CH3—C(=O)—	(tetrahydropyran with CH3)—O⁻

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A with O-)
613	(4-F-phenyl)-O-CH₂-CO-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CO-CH₃	3-methyl-tetrahydropyran-2-yl-O-
614	(2-Cl-phenyl)-O-CH₂-CO-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CO-CH₃	3-methyl-tetrahydropyran-2-yl-O-
615	(3-Cl-phenyl)-O-CH₂-CO-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CO-CH₃	3-methyl-tetrahydropyran-2-yl-O-
616	(4-Cl-phenyl)-O-CH₂-CO-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CO-CH₃	3-methyl-tetrahydropyran-2-yl-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
617	(2-methoxyphenoxy)acetone structure, OCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–CO–	tetrahydropyran-2-yloxy (methyl)
618	(3-methoxyphenoxy)acetone structure, CH$_3$O	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–CO–	tetrahydropyran-2-yloxy (methyl)
619	(4-methoxyphenoxy)acetone structure, CH$_3$O	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–CO–	tetrahydropyran-2-yloxy (methyl)
620	(2,4-dimethoxyphenoxy)acetone structure, CH$_3$O, OCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–CO–	tetrahydropyran-2-yloxy (methyl)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
621	2-methylphenyl–O–CH₂–C(O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–C(O)CH₃	tetrahydropyranyl-O– (with A, methyl)
622	3-methylphenyl–O–CH₂–C(O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–C(O)CH₃	tetrahydropyranyl-O– (methyl)
623	4-methylphenyl–O–CH₂–C(O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–C(O)CH₃	tetrahydropyranyl-O– (methyl)
624	2-(CF₃)phenyl–O–CH₂–C(O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–C(O)CH₃	tetrahydropyranyl-O– (methyl)

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A⟩O–O–
625	3-(CF₃)phenyl–O–CH₂–C(=O)–CH₃	–H	$-CH_2CH(CH_3)_2$	–H	CH₃–C(=O)–	tetrahydropyran-methyl O–
626	4-(CF₃)phenyl–O–CH₂–C(=O)–CH₃	–H	$-CH_2CH(CH_3)_2$	–H	CH₃–C(=O)–	tetrahydropyran-methyl O–
627	naphthalen-1-yl–O–CH₂–C(=O)–CH₃	–H	$-CH_2CH(CH_3)_2$	–H	CH₃–C(=O)–	tetrahydropyran-methyl O–
628	naphthalen-2-yl–O–CH₂–C(=O)–CH₃	–H	$-CH_2CH(CH_3)_2$	–H	CH₃–C(=O)–	tetrahydropyran-methyl O–

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
629	(benzyloxyacetone structure)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(2-methyltetrahydropyranyl-O- structure)
630	(phenylthioacetone structure)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(2-methyltetrahydropyranyl-O- structure)
631	(phenylsulfonyl butanone structure)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(2-methyltetrahydropyranyl-O- structure)
632	(acetophenone structure)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(2-methyltetrahydropyranyl-O- structure)

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
633	(2-fluoroacetophenyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 acetyl	(tetrahydropyran-methyl-O⁻)
634	(3-fluoroacetophenyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 acetyl	(tetrahydropyran-methyl-O⁻)
635	(4-fluoroacetophenyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 acetyl	(tetrahydropyran-methyl-O⁻)
636	(2,6-difluoroacetophenyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 acetyl	(tetrahydropyran-methyl-O⁻)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
637	(2,4-difluoro acetophenone)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(CH$_3$C=O)	(tetrahydropyranyl-O-, methyl)
638	(2,6-difluoro-4-fluoro acetophenone)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(CH$_3$C=O)	(tetrahydropyranyl-O-, methyl)
639	(2-chloro acetophenone)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(CH$_3$C=O)	(tetrahydropyranyl-O-, methyl)
640	(3-chloro acetophenone)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(CH$_3$C=O)	(tetrahydropyranyl-O-, methyl)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
641	4-Cl-C₆H₄-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(tetrahydropyran-2-yl-oxy, 3-methyl)
642	2-Br-C₆H₄-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(tetrahydropyran-2-yl-oxy, 3-methyl)
643	3-Br-C₆H₄-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(tetrahydropyran-2-yl-oxy, 3-methyl)
644	4-Br-C₆H₄-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	(tetrahydropyran-2-yl-oxy, 3-methyl)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
645	(2-acetylphenyl)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(tetrahydropyranyl-methyl, -O⁻)
646	(3-acetylphenyl)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(tetrahydropyranyl-methyl, -O⁻)
647	(4-acetylphenyl)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(tetrahydropyranyl-methyl, -O⁻)
648	(2-methyl-4-acetylphenyl)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(tetrahydropyranyl-methyl, -O⁻)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
649	2,6-(CH₃)₂-phenyl-C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	tetrahydropyran-2-yl-O- (with A ring)
650	3,4-(H₃C)₂-phenyl-C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)
651	2-CH₃-4,6-(H₃C)₂... phenyl-C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)
652	2-CH₂-,-H₃C-phenyl-C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)

116

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
653	H₃C—⟨benzene⟩—C(=O)CH₃ (acetyl)	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	tetrahydropyran-2-yl-O⁻ with methyl
654	H₃C—⟨benzene⟩—C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	tetrahydropyran-2-yl-O⁻ with methyl
655	⟨benzene⟩ with CF₃, —C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	tetrahydropyran-2-yl-O⁻ with methyl
656	F₃C—⟨benzene⟩—C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	tetrahydropyran-2-yl-O⁻ with methyl

117

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring structure)
657	(4-F_3C-phenyl)C(=O)-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)
658	(2-CH_3O-phenyl)C(=O)-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)
659	(3-CH_3O-phenyl)C(=O)-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)
660	(4-CH_3O-phenyl)C(=O)-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-C(=O)-	tetrahydropyran-2-yl-O- (3-methyl)

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
661		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
662		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
663		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
664		-H	-CH$_2$CH(CH$_3$)$_2$	-H		

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
665	CH_3O ... OCH_3 ... CH_3O (substituted aryl with $-COCH_3$)	-H	$-CH_2CH(CH_3)_2$	-H	$-COCH_3$	(tetrahydropyranyl-O-)
666	aryl with $-COCH_3$ and H_3C-O	-H	$-CH_2CH(CH_3)_2$	-H	$-COCH_3$	(tetrahydropyranyl-O-)
667	aryl with $-COCH_3$ and H_3C-O	-H	$-CH_2CH(CH_3)_2$	-H	$-COCH_3$	(tetrahydropyranyl-O-)
668	aryl with $-COCH_3$ and H_3C-O	-H	$-CH_2CH(CH_3)_2$	-H	$-COCH_3$	(tetrahydropyranyl-O-)

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
669		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
670		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
671		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
672		-H	-CH$_2$CH(CH$_3$)$_2$	-H		

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
673		-H	$-CH_2CH(CH_3)_2$	-H		
674		-H	$-CH_2CH(CH_3)_2$	-H		
675		-H	$-CH_2CH(CH_3)_2$	-H		
676		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
677	(2-acetylnaphthyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$C(=O)-	(tetrahydropyranyl, A-O-O-)
678	(2-acetylpyridyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$C(=O)-	(2-methyltetrahydropyranyloxy)
679	(3-acetylpyridyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$C(=O)-	(2-methyltetrahydropyranyloxy)
680	(4-acetylpyridyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$C(=O)-	(2-methyltetrahydropyranyloxy)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
681		-H	$-CH_2CH(CH_3)_2$	-H		
682		-H	$-CH_2CH(CH_3)_2$	-H		
683		-H	$-CH_2CH(CH_3)_2$	-H		
684		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A ring structure)
685		-H	$-CH_2CH(CH_3)_2$	-H		
686		-H	$-CH_2CH(CH_3)_2$	-H		
687		-H	$-CH_2CH(CH_3)_2$	-H		
688		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A ring structure)
689	(2-acetyl-pyrazole)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(ring with O and O⁻, CH₃)
690	(acetyl-methyl-pyrazole, N-CH₃)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(ring with O and O⁻, CH₃)
691	(acetyl-imidazole, N-H)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(ring with O and O⁻, CH₃)
692	(acetyl-imidazole, N-CH₃)	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	(ring with O and O⁻, CH₃)

126

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
693		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
694		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
695		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
696		-H	-CH$_2$CH(CH$_3$)$_2$	-H		

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
697		-H	$-CH_2CH(CH_3)_2$	-H		
698		-H	$-CH_2CH(CH_3)_2$	-H		
699		-H	$-CH_2CH(CH_3)_2$	-H		
700		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
701	H$_3$C–CH=CH–C(=O)–	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	(A-substituted tetrahydropyranyl-O–)
702	C$_6$H$_5$–CH=CH–C(=O)–	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	(3-methyltetrahydropyran-2-yl-O–)
703	4-F–C$_6$H$_4$–CH=CH–C(=O)–	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	(3-methyltetrahydropyran-2-yl-O–)
704	4-Cl–C$_6$H$_4$–CH=CH–C(=O)–	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	(3-methyltetrahydropyran-2-yl-O–)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
705	H₃C–C₆H₄–CH=CH–CH=CH–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl (3-methyl), –O–
706	F₃C–C₆H₄–CH=CH–CH=CH–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl (3-methyl), –O–
707	CH₃O–C₆H₄–CH=CH–CH=CH–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl (3-methyl), –O–
708	CH₃O–C₆H₃(OCH₃)–CH=CH–CH=CH–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydrofuran-2-yl (3-methyl), –O–

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
709	3,4-dimethoxyphenyl-CH=CH-C(=O)-CH₃ (CH₃O, CH₃O)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl-O– (with A, methyl)
710	2,4-dimethoxy-6-methoxyphenyl-CH=CH-C(=O)-CH₃ (OCH₃, OCH₃, CH₃O)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl-O– (methyl)
711	$H_3C-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl-O– (methyl)
712	$H_3C-CH_2-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3–C(=O)–	tetrahydropyran-2-yl-O– (methyl)

131

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring structure)
713	(H₃C)₂CH–S(=O)(=O)–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring, O, O–, methyl)
714	cyclohexyl–S(=O)(=O)–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring, O, O–, methyl)
715	benzyl–S(=O)(=O)–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring, O, O–, methyl)
716	phenyl–S(=O)(=O)–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring, O, O–, methyl)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
717	phenyl–$S(=O)_2$–	$-CH_3$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	(A)-ring with –O– and CH₃
718	phenyl–$S(=O)_2$–	$-H$	$-CH_2CH(CH_3)_2$	$-CH_3$	CH_3–C(=O)–	tetrahydropyranyl –O– with CH₃
719	phenyl–$S(=O)_2$–	$-CH_3$	$-CH_2CH(CH_3)_2$	$-CH_3$	CH_3–C(=O)–	tetrahydropyranyl –O– with CH₃
720	phenyl–$S(=O)_2$–	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–CH₂–C(=O)–	tetrahydrofuranyl –O– with CH₃

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(ring)
721	phenyl–S(=O)(=O)–	-H	$-CH_2CH(CH_3)_2$	-H	(ketone structure)	(tetrahydropyran-O– with A)
722	phenyl–S(=O)(=O)–	-H	$-CH_2CH(CH_3)_2$	-H	(ketone structure)	(tetrahydropyran-O–)
723	phenyl–S(=O)(=O)–	-H	$-CH_2CH(CH_3)_2$	-H	(ketone structure)	(tetrahydropyran-O–)
724	phenyl–S(=O)(=O)–	-H	$-CH_2CH(CH_3)_2$	-H	(ketone structure)	(tetrahydropyran-O–)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
725	phenyl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3, CH_3, CH_3 — $C(CH_3)_2$ ketone	(A) ring, methyl, $-O-$
726	phenyl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3 chain ketone	tetrahydropyranyl-methyl $-O-$
727	phenyl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3, CH_3 ketone	tetrahydropyranyl-methyl $-O-$
728	phenyl-SO_2-	$-H$	$-CH_2CH(CH_3)_2$	$-H$	phenyl-$C(=O)-CH_3$	tetrahydropyranyl-methyl $-O-$

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
729	phenyl-SO_2 ($O=S=O$)	-H	$-CH_2CH(CH_3)_2$	-H	4-F-phenyl-C(=O)-	3-methyltetrahydropyran-2-yl-O-
730	phenyl-SO_2 ($O=S=O$)	-H	$-CH_2CH(CH_3)_2$	-H	4-Cl-phenyl-C(=O)-	3-methyltetrahydropyran-2-yl-O-
731	phenyl-SO_2 ($O=S=O$)	-H	$-CH_2CH(CH_3)_2$	-H	2-Cl-phenyl-C(=O)-	3-methyltetrahydropyran-2-yl-O-
732	phenyl-SO_2 ($O=S=O$)	-H	$-CH_2CH(CH_3)_2$	-H	2-CH_3-phenyl-C(=O)-	3-methyltetrahydropyran-2-yl-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
733	phenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	4-(CH_3)-phenyl-C(=O)-	tetrahydrofuranyl-(methyl)-O- (A)
734	phenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	4-(CF_3)-phenyl-C(=O)-	tetrahydrofuranyl-(methyl)-O-
735	phenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	4-(OCH_3)-phenyl-C(=O)-	tetrahydrofuranyl-(methyl)-O-
736	phenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	4-($O-C(CH_3)_2CH_3$)-phenyl-C(=O)-	tetrahydrofuranyl-(methyl)-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
737	phenyl-SO$_2$ (O=S=O)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	3,4,5-tri(CH$_3$O)phenyl-C(=O)-	2-methyltetrahydropyran-2-yl-O-
738	phenyl-SO$_2$ (O=S=O)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	4-[N(CH$_3$)CH$_3$]phenyl-C(=O)-	2-methyltetrahydropyran-2-yl-O-
739	phenyl-SO$_2$ (O=S=O)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	naphthalen-1-yl-C(=O)-	2-methyltetrahydropyran-2-yl-O-
740	phenyl-SO$_2$ (O=S=O)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	naphthalen-2-yl-C(=O)-	2-methyltetrahydropyran-2-yl-O-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	structure
741	$O{=}S{=}O$ phenyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_3$	
742	$O{=}S{=}O$ phenyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH_3$	
743	$O{=}S{=}O$ phenyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH_2CH_3$	
744	$O{=}S{=}O$ phenyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH(CH_3)_2$	

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
745	$O{=}S{=}O$ phenyl	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_2CH_3$	
746	$O{=}S{=}O$ phenyl	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	
747	$O{=}S{=}O$ phenyl	-H	$-CH_2CH(CH_3)_2$	-H	$-C(CH_3)_3$	
748	$O{=}S{=}O$ phenyl	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH(CH_3)_2$	

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
749	2-F-C₆H₄-SO₂- (2-fluorophenylsulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH₃-CO-	
750	3-F-C₆H₄-SO₂- (3-fluorophenylsulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH₃-CO-	
751	4-F-C₆H₄-SO₂- (4-fluorophenylsulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH₃-CO-	
752	2-Cl-C₆H₄-SO₂- (2-chlorophenylsulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH₃-CO-	

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A-O-O)
753	3-Cl-C$_6$H$_4$-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	tetrahydropyran-2-yl-oxy (3-methyl)
754	4-Cl-C$_6$H$_4$-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	tetrahydropyran-2-yl-oxy (3-methyl)
755	4-Cl-C$_6$H$_4$-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(CH$_3$)$_3$C-C(=O)-	tetrahydropyran-2-yl-oxy (3-methyl)
756	2-Br-C$_6$H$_4$-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	tetrahydropyran-2-yl-oxy (3-methyl)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	[ring structure]
757	(benzenesulfonyl, O=S=O, with Br)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(C(=O)CH$_3$)	(tetrahydropyran-2-yl-O⁻, methyl)
758	(benzenesulfonyl, O=S=O, with Br)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(C(=O)CH$_3$)	(tetrahydropyran-2-yl-O⁻, methyl)
759	(benzenesulfonyl, O=S=O, with CH$_3$)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(C(=O)CH$_3$)	(tetrahydropyran-2-yl-O⁻, methyl)
760	(benzenesulfonyl, O=S=O, with H$_3$C)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(C(=O)CH$_3$)	(A-O-O⁻, methyl)

143

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
761	4-H₃C–C₆H₄–SO₂– (O=S=O)	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–CO–	(lactone-O–)
762	2-CH₃-4-H₃C–C₆H₃–SO₂–	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–CO–	(lactone-O–)
763	2,6-(CH₃)₂–C₆H₃–SO₂–	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–CO–	(lactone-O–)
764	3,5-(H₃C)₂–C₆H₃–SO₂–	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–CO–	(lactone-O–)

144

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
765	2,4,6-trimethylphenyl–SO_2– (mesitylsulfonyl)	–H	–$CH_2CH(CH_3)_2$	–H	CH_3–C(=O)– (acetyl)	(ring: A–O···O– with methyl)
766	4-(CH₂CH₃ as H_3C–CH₂–)phenyl–SO_2–	–H	–$CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-2-yloxy (methyl)
767	4-(H_3C–CH–CH_3)phenyl–SO_2– (isopropylphenylsulfonyl)	–H	–$CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-2-yloxy (methyl)
768	4-(H_3C)(H_3C)(H_3C)C–phenyl–SO_2– (tert-butylphenylsulfonyl)	–H	–$CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-2-yloxy (methyl)

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
769	2-methoxyphenyl-SO₂– (benzene with OCH₃, $O=S=O$)	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydrofuran with methyl, O–)
770	3-methoxyphenyl-SO₂– (benzene with CH₃O, $O=S=O$)	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydrofuran with methyl, O–)
771	4-methoxyphenyl-SO₂– (CH₃O, $O=S=O$)	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydrofuran with methyl, O–)
772	4-(H₃C–CH₂O)phenyl-SO₂– ($O=S=O$)	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydrofuran with methyl, O–)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
773		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
774		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
775		-H	-CH$_2$CH(CH$_3$)$_2$	-H		
776		-H	-CH$_2$CH(CH$_3$)$_2$	-H		

Table 2   (Continued)

Compound No.	R[1]	R[4]	R[5]	R[6]	R[7]	
777	3-nitrophenyl-SO$_2$– (O_2N)	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran–O$^-$ (methyl)
778	4-nitrophenyl-SO$_2$– (O_2N)	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran–O$^-$ (methyl)
779	naphthalen-1-yl-SO$_2$–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran–O$^-$ (methyl)
780	naphthalen-2-yl-SO$_2$–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran–O$^-$ (methyl)

148

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
781	pyridin-2-yl-SO₂–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	tetrahydropyran-methyl-O–
782	pyridin-3-yl-SO₂–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	tetrahydropyran-methyl-O–
783	pyridin-4-yl-SO₂–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	tetrahydropyran-methyl-O–
784	furan-2-yl-SO₂–	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	tetrahydrofuran-methyl-O–

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
785	(thiophene-S,S-dioxide sulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$ (acetyl)	(3-methyltetrahydropyran-2-yl)O-
786	(1H-pyrrole sulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$ (acetyl)	(3-methyltetrahydropyran-2-yl)O-
787	(N-CH₃ pyrrole sulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$ (acetyl)	(3-methyltetrahydropyran-2-yl)O-
788	(isoxazole sulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$ (acetyl)	(3-methyltetrahydropyran-2-yl)O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
789		-H	$-CH_2CH(CH_3)_2$	-H		
790		-H	$-CH_2CH(CH_3)_2$	-H		
791	H-	-H	$-CH_2$ (phenyl)	-H		
792		-H	$-CH_2$ (phenyl)	-H		

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
793	(structure)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(structure)
794	(structure)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(structure)
795	(structure)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(structure)
796	(structure)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(structure)

152

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A)
797	(2-fluorobenzyl acetate)	$-H$	$-CH_2-$phenyl	$-H$	$CH_3-C(=O)-$	(tetrahydrofuranyl with CH₃, O-)
798	(4-fluorobenzyl acetate)	$-H$	$-CH_2-$phenyl	$-H$	$CH_3-C(=O)-$	(tetrahydrofuranyl with CH₃, O-)
799	(2-chlorobenzyl acetate)	$-H$	$-CH_2-$phenyl	$-H$	$CH_3-C(=O)-$	(tetrahydrofuranyl with CH₃, O-)
800	(4-chlorobenzyl acetate)	$-H$	$-CH_2-$phenyl	$-H$	$CH_3-C(=O)-$	(tetrahydrofuranyl with CH₃, O-)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A) structure
801	2-CH₃-benzyl acetate ester	-H	-CH₂-C₆H₅	-H	CH₃-CO-	3-methyltetrahydropyran-2-yl-O-
802	4-H₃C-benzyl acetate ester	-H	-CH₂-C₆H₅	-H	CH₃-CO-	3-methyltetrahydropyran-2-yl-O-
803	2-OCH₃-benzyl acetate ester	-H	-CH₂-C₆H₅	-H	CH₃-CO-	3-methyltetrahydropyran-2-yl-O-
804	4-CH₃O-benzyl acetate ester	-H	-CH₂-C₆H₅	-H	CH₃-CO-	3-methyltetrahydropyran-2-yl-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(ring structure)
805	phenyl-NH-C(=O)-CH3	-H	-CH2-phenyl	-H	CH3-C(=O)-	tetrahydropyran-2-yl, 3-methyl, -O-
806	H3C-CH(CH3)-... (H3C)(H3C)CH-C(=O)-	-H	-CH2-phenyl	-H	CH3-C(=O)-	tetrahydropyran-2-yl, 3-methyl, -O-
807	phenyl-CH2-CH2-C(=O)-CH3	-H	-CH2-phenyl	-H	CH3-C(=O)-	tetrahydropyran-2-yl, 3-methyl, -O-
808	phenyl-O-CH2-C(=O)-CH3	-H	-CH2-phenyl	-H	CH3-C(=O)-	tetrahydropyran-2-yl, 3-methyl, -O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	structure
809	(2-chlorophenoxy)acetone structure	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	tetrahydropyran-methyl-O⁻
810	(4-chlorophenoxy)acetone structure	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	tetrahydropyran-methyl-O⁻
811	(2-methylphenoxy)acetone structure	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	tetrahydropyran-methyl-O⁻
812	(4-methylphenoxy)acetone structure	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	tetrahydropyran-methyl-O⁻

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
813	(2-OCH$_3$-phenoxy)-CH$_2$-C(=O)-CH$_3$	-H	-CH$_2$-phenyl	-H	CH$_3$-C(=O)-	tetrahydropyranyl-CH$_3$, O⁻
814	(4-CH$_3$O-phenoxy)-CH$_2$-C(=O)-CH$_3$	-H	-CH$_2$-phenyl	-H	CH$_3$-C(=O)-	tetrahydropyranyl-CH$_3$, O⁻
815	phenyl-C(=O)-CH$_3$	-H	-CH$_2$-phenyl	-H	CH$_3$-C(=O)-	tetrahydropyranyl-CH$_3$, O⁻
816	(2-F-phenyl)-C(=O)-CH$_3$	-H	-CH$_2$-phenyl	-H	CH$_3$-C(=O)-	tetrahydropyranyl-CH$_3$, O⁻

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A—O—O⁻
817	2-F-C₆H₄-C(=O)CH₃	-H	-CH₂-C₆H₅	-H	CH₃C(=O)-	(tetrahydropyran-methyl)-O⁻
818	4-F-C₆H₄-C(=O)CH₃	-H	-CH₂-C₆H₅	-H	CH₃C(=O)-	(tetrahydropyran-methyl)-O⁻
819	2-Cl-C₆H₄-C(=O)CH₃	-H	-CH₂-C₆H₅	-H	CH₃C(=O)-	(tetrahydropyran-methyl)-O⁻
820	3-Cl-C₆H₄-C(=O)CH₃	-H	-CH₂-C₆H₅	-H	CH₃C(=O)-	(tetrahydropyran-methyl)-O⁻

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A–O–)
821	4-Cl-C₆H₄-C(=O)-CH₃	-H	-CH₂-C₆H₅	-H	-C(=O)-CH₃	tetrahydropyranyl-methyl-O-
822	2-CH₃-C₆H₄-C(=O)-CH₃	-H	-CH₂-C₆H₅	-H	-C(=O)-CH₃	tetrahydropyranyl-methyl-O-
823	3-CH₃-C₆H₄-C(=O)-CH₃	-H	-CH₂-C₆H₅	-H	-C(=O)-CH₃	tetrahydropyranyl-methyl-O-
824	4-CH₃-C₆H₄-C(=O)-CH₃	-H	-CH₂-C₆H₅	-H	-C(=O)-CH₃	tetrahydropyranyl-methyl-O-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A)
825	2-CH₃O-phenyl-C(O)-	-H	-CH₂-phenyl	-H	CH₃-C(O)-	tetrahydrofuran-O- (3-methyl)
826	3-CH₃O-phenyl-C(O)-	-H	-CH₂-phenyl	-H	CH₃-C(O)-	tetrahydrofuran-O- (3-methyl)
827	4-CH₃O-phenyl-C(O)-	-H	-CH₂-phenyl	-H	CH₃-C(O)-	tetrahydrofuran-O- (3-methyl)
828	pyridin-2-yl-C(O)-	-H	-CH₂-phenyl	-H	CH₃-C(O)-	tetrahydrofuran-O- (3-methyl)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
829	3-acetylpyridinyl (C(=O)CH₃ on pyridine, N)	–H	–CH₂– (phenyl)	–H	CH₃–C(=O)–	tetrahydropyran, CH₃, –O⁻
830	3-acetylpyridinyl (C(=O)CH₃ on pyridine, N)	–H	–CH₂– (phenyl)	–H	CH₃–C(=O)–	tetrahydropyran, CH₃, –O⁻
831	H₃C–S(=O)(=O)–	–H	–CH₂– (phenyl)	–H	CH₃–C(=O)–	tetrahydropyran, CH₃, –O⁻
832	(H₃C)₂... S(=O)(=O)– (H₃C, H₃C on S, O=S=O)	–H	–CH₂– (phenyl)	–H	CH₃–C(=O)–	tetrahydropyran, CH₃, –O⁻

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A
833	benzyl–SO_2– ($O=S=O$, CH_2-phenyl)	–H	–CH_2(phenyl)	–H	CH_3–C(=O)–	tetrahydropyranyl–O⁻ (methyl substituted)
834	phenyl–SO_2– ($O=S=O$)	–H	–CH_2(phenyl)	–H	CH_3–C(=O)–	tetrahydropyranyl–O⁻ (methyl substituted)
835	4-F-phenyl–SO_2– ($O=S=O$)	–H	–CH_2(phenyl)	–H	CH_3–C(=O)–	tetrahydropyranyl–O⁻ (methyl substituted)
836	4-Cl-phenyl–SO_2– ($O=S=O$)	–H	–CH_2(phenyl)	–H	CH_3–C(=O)–	tetrahydropyranyl–O⁻ (methyl substituted)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
837	4-Br-phenyl-SO_2-	-H	-CH_2-phenyl	-H	CH_3-C(=O)- (acetyl)	tetrahydropyranyl-O⁻ (methyl)
838	4-CH_3-phenyl-SO_2-	-H	-CH_2-phenyl	-H	CH_3-C(=O)-	tetrahydropyranyl-O⁻ (methyl)
839	4-CH_3O-phenyl-SO_2-	-H	-CH_2-phenyl	-H	CH_3-C(=O)-	tetrahydropyranyl-O⁻ (methyl)
840	3-O_2N-phenyl-SO_2-	-H	-CH_2-phenyl	-H	CH_3-C(=O)-	tetrahydropyranyl-O⁻ (methyl)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
841	benzyl ester (–CH₂–O–C(=O)–CH₃ on phenyl)	–H	1-naphthylmethyl (–CH₂)	–H	–C(=O)CH₃	tetrahydropyran ring (A) with –O⁻ and CH₃
842	phenyl sulfonyl (–S(=O)₂–phenyl)	–H	1-naphthylmethyl (–CH₂)	–H	–C(=O)CH₃	tetrahydropyran ring with –O⁻ and CH₃
843	benzyl ester (–CH₂–O–C(=O)–CH₃ on phenyl)	–H	2-naphthylmethyl (–CH₂)	–H	–C(=O)CH₃	tetrahydropyran ring with –O⁻ and CH₃
844	phenyl sulfonyl (–S(=O)₂–phenyl)	–H	2-naphthylmethyl (–CH₂)	–H	–C(=O)CH₃	tetrahydropyran ring with –O⁻ and CH₃

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
845	benzyl acetate	-H	$-CH_2CH_2SCH_3$	-H	CH_3 C=O (acetyl)	tetrahydropyran-2-yl-O- (with A) methyl
846	phenyl O=S=O	-H	$-CH_2CH_2SCH_3$	-H	CH_3 C=O	tetrahydropyran-2-yl-O- methyl
847	benzyl acetate	-H	$-CH_2OC(CH_3)_3$	-H	CH_3 C=O	tetrahydropyran-2-yl-O- methyl
848	phenyl O=S=O	-H	$-CH_2OC(CH_3)_3$	-H	CH_3 C=O	tetrahydropyran-2-yl-O- methyl

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(ring)
849	benzyl –CH₂–O–C(=O)CH₃	–H	–CH₂OCH₂(C₆H₅)	–H	CH₃–C(=O)– (acetyl)	tetrahydropyran, CH₃, –O⁻
850	phenyl–S(=O)(=O)–	–H	–CH₂OCH₂(C₆H₅)	–H	CH₃–C(=O)– (acetyl)	tetrahydropyran, CH₃, –O⁻
851	benzyl –CH₂–O–C(=O)CH₃	–H	–CH₂OH	–H	CH₃–C(=O)– (acetyl)	tetrahydropyran, CH₃, –O⁻
852	phenyl–S(=O)(=O)–	–H	–CH₂OH	–H	CH₃–C(=O)– (acetyl)	tetrahydropyran, CH₃, –O⁻

Table 2   (Continued)

	R⁷	R⁶	R⁵	R⁴	R¹	Compound No.
(CH₃-substituted tetrahydrofuranyl ring, —O—)	CH₃–C(=O)–	–H	–H	–H	benzyl acetate group (–CH₂–O–C(=O)CH₃, phenyl)	853
(CH₃-substituted tetrahydrofuranyl ring, —O—)	CH₃–C(=O)–	–H	–H	–H	phenyl–S(=O)₂–	854
(CH₃-substituted tetrahydrofuranyl ring, —O—)	CH₃–C(=O)–	–H	–CH₃	–H	benzyl acetate group (–CH₂–O–C(=O)CH₃, phenyl)	855
(CH₃-substituted tetrahydrofuranyl ring, —O—)	CH₃–C(=O)–	–H	–CH₃	–H	phenyl–S(=O)₂–	856

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
857		$-H$	$-CH_2CH_2CH_3$	$-H$		
858		$-H$	$-CH_2CH_2CH_3$	$-H$		
859		$-H$	$-CH(CH_3)_2$	$-H$		
860		$-H$	$-CH(CH_3)_2$	$-H$		

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
861	(benzyl — O—C(=O)CH$_3$)	-H	-CH$_2$CH$_2$CH$_2$CH$_3$	-H	CH$_3$—C(=O)—	(tetrahydropyran ring, CH$_3$, —O—)
862	(phenyl—S(=O)(=O)—)	-H	-CH$_2$CH$_2$CH$_2$CH$_3$	-H	CH$_3$—C(=O)—	(tetrahydropyran ring, CH$_3$, —O—)
863	H-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$—C(=O)—	(tetrahydrofuran ring, CH$_3$, —O—)
864	((H$_3$C)$_3$C—O—C(=O)CH$_3$)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$—C(=O)—	(tetrahydrofuran ring, CH$_3$, —O—)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
865	$C_6H_5CH_2\text{–}O\text{–}C(=O)CH_3$	–H	$-CH_2CH(CH_3)_2$	–H	$CH_3\text{–}C(=O)\text{–}$	tetrahydrofuran ring with CH_3 and O^-
866	$C_6H_5CH_2CH_2\text{–}C(=O)CH_3$	–H	$-CH_2CH(CH_3)_2$	–H	$CH_3\text{–}C(=O)\text{–}$	tetrahydrofuran ring with CH_3 and O^-
867	$C_6H_5\text{–}O\text{–}CH_2\text{–}C(=O)CH_3$	–H	$-CH_2CH(CH_3)_2$	–H	$CH_3\text{–}C(=O)\text{–}$	tetrahydrofuran ring with CH_3 and O^-
868	$C_6H_5\text{–}C(=O)CH_3$	–H	$-CH_2CH(CH_3)_2$	–H	$CH_3\text{–}C(=O)\text{–}$	tetrahydrofuran ring with CH_3 and O^-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
869	2-F-phenyl-C(=O)-CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	
870	phenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	
871	benzyl-O-C(=O)-CH₃	-H	$-CH_2-$phenyl	-H	$CH_3-C(=O)-$	
872	phenyl-SO_2-	-H	$-CH_2-$phenyl	-H	$CH_3-C(=O)-$	

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(ring structure)
873	phenyl-CH_2-O-C(=O)CH_3	-H	-$CH_2CH_2SCH_3$	-H	CH_3-C(=O)-	CH_3-substituted ring, O, O-
874	phenyl-S(=O)(=O)-	-H	-$CH_2CH_2SCH_3$	-H	CH_3-C(=O)-	CH_3-substituted ring, O, O-
875	phenyl-CH_2-O-C(=O)CH_3	-H	-$CH_2OC(CH_3)_3$	-H	CH_3-C(=O)-	CH_3-substituted ring, O, O-
876	phenyl-S(=O)(=O)-	-H	-$CH_2OC(CH_3)_3$	-H	CH_3-C(=O)-	CH_3-substituted ring, O, O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure, A)
877	benzyl acetate (Ph–CH$_2$–O–C(=O)CH$_3$)	–H	–CH$_2$OH	–H	acetyl (CH$_3$–C(=O)–)	CH$_3$-substituted tetrahydropyran –O–
878	phenylsulfonyl (Ph–S(=O)$_2$–)	–H	–CH$_2$OH	–H	acetyl (CH$_3$–C(=O)–)	CH$_3$-substituted tetrahydropyran –O–
879	benzyl acetate (Ph–CH$_2$–O–C(=O)CH$_3$)	–H	tolyl	–H	acetyl (CH$_3$–C(=O)–)	CH$_3$-substituted tetrahydropyran –O–
880	phenylsulfonyl (Ph–S(=O)$_2$–)	–H	tolyl	–H	acetyl (CH$_3$–C(=O)–)	CH$_3$-substituted tetrahydropyran –O–

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(A)
881	benzyl –CH₂–O–C(=O)–CH₃	–H	–H	–H	–C(=O)–CH₃	(tetrahydrofuran ring, CH_3, H_3C) –O–
882	phenyl –S(=O)(=O)–	–H	–H	–H	–C(=O)–CH₃	(tetrahydrofuran ring, CH_3, H_3C) –O–
883	benzyl –CH₂–O–C(=O)–CH₃	–H	–CH₃	–H	–C(=O)–CH₃	(tetrahydrofuran ring, CH_3, H_3C) –O–
884	phenyl –S(=O)(=O)–	–H	–CH₃	–H	–C(=O)–CH₃	(tetrahydrofuran ring, CH_3, H_3C) –O–

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
885	(benzyl acetate ester)	-H	$-CH_2CH_2CH_3$	-H	(acetyl, $CH_3C(=O)-$)	(tetrahydropyran structure)
886	(phenyl sulfonyl, $O=S=O$)	-H	$-CH_2CH_2CH_3$	-H	(acetyl, $CH_3C(=O)-$)	(tetrahydropyran structure)
887	(benzyl acetate ester)	-H	$-CH(CH_3)_2$	-H	(acetyl, $CH_3C(=O)-$)	(tetrahydropyran structure)
888	(phenyl sulfonyl, $O=S=O$)	-H	$-CH(CH_3)_2$	-H	(acetyl, $CH_3C(=O)-$)	(tetrahydropyran structure)

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A structure
889	benzyl –CH$_2$–O–C(=O)CH$_3$	–H	–CH$_2$CH$_2$CH$_2$CH$_3$	–H	–C(=O)CH$_3$	structure
890	phenyl –S(=O)$_2$–	–H	–CH$_2$CH$_2$CH$_2$CH$_3$	–H	–C(=O)CH$_3$	structure
891	–H	–H	–CH$_2$CH(CH$_3$)$_2$	–H	–C(=O)CH$_3$	structure
892	(H$_3$C)$_3$C–O–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	–C(=O)CH$_3$	structure

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
893	benzyl acetate (C₆H₅CH₂–O–C(=O)CH₃)	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)–	tetrahydrofuran ring (H₃C, CH₃, –O⁻)
894	4-fluorobenzyl acetate (4-F-C₆H₄CH₂–O–C(=O)CH₃)	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)–	tetrahydrofuran ring (H₃C, CH₃, –O⁻)
895	4-chlorobenzyl acetate (4-Cl-C₆H₄CH₂–O–C(=O)CH₃)	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)–	tetrahydrofuran ring (H₃C, CH₃, –O⁻)
896	4-bromobenzyl acetate (4-Br-C₆H₄CH₂–O–C(=O)CH₃)	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)–	tetrahydrofuran ring (H₃C, CH₃, –O⁻)

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
897	4-CH₃-benzyl acetate (H₃C–C₆H₄–CH₂–O–C(=O))	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)– (acetyl)	3,3-dimethyl-4-methyltetrahydrofuran-O–
898	4-CH₃O-benzyl acetate (CH₃O–C₆H₄–CH₂–O–C(=O))	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)– (acetyl)	3,3-dimethyl-4-methyltetrahydrofuran-O–
899	H₃C–C(=O)–	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)– (acetyl)	3,3-dimethyl-4-methyltetrahydrofuran-O–
900	(H₃C)₂CH–CH₂–C(=O)–	–H	–CH₂CH(CH₃)₂	–H	CH₃C(=O)– (acetyl)	3,3-dimethyl-4-methyltetrahydrofuran-O–

178

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
901		-H	$-CH_2CH(CH_3)_2$	-H		
902		-H	$-CH_2CH(CH_3)_2$	-H		
903		-H	$-CH_2CH(CH_3)_2$	-H		
904		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2 (Continued)

Compound No.	R1	R4	R5	R6	R7	A structure
905	Cl-phenyl-O-CH2-C(=O)-CH3	-H	-CH2CH(CH3)2	-H	CH3-C(=O)-	tetrahydrofuran ring with CH3, H3C, CH3, O-
906	H3C-phenyl-O-CH2-C(=O)-CH3	-H	-CH2CH(CH3)2	-H	CH3-C(=O)-	tetrahydrofuran ring with CH3, H3C, CH3, O-
907	CH3O-phenyl-O-CH2-C(=O)-CH3	-H	-CH2CH(CH3)2	-H	CH3-C(=O)-	tetrahydrofuran ring with CH3, H3C, CH3, O-
908	CH3O-,OCH3-phenyl-O-CH2-C(=O)-CH3	-H	-CH2CH(CH3)2	-H	CH3-C(=O)-	tetrahydrofuran ring with CH3, H3C, CH3, O-

180

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A
909	(acetyl-phenyl)	–H	–CH₂CH(CH₃)₂	–H	COCH₃	(tetrahydrofuranyl with CH₃, H₃C, CH₃)
910	(acetyl-fluorophenyl)	–H	–CH₂CH(CH₃)₂	–H	COCH₃	(tetrahydrofuranyl with CH₃, H₃C, CH₃)
911	(acetyl-fluorophenyl)	–H	–CH₂CH(CH₃)₂	–H	COCH₃	(tetrahydrofuranyl with CH₃, H₃C, CH₃)
912	(acetyl-fluorophenyl)	–H	–CH₂CH(CH₃)₂	–H	COCH₃	(tetrahydrofuranyl with CH₃, H₃C, CH₃)

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A
913	(2-Cl-phenyl)C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3C(=O)-	tetrahydropyranyl (structure)
914	(4-Cl-phenyl)C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3C(=O)-	tetrahydrofuranyl (structure)
915	(2-CH₃-phenyl)C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3C(=O)-	tetrahydropyranyl (structure)
916	(4-CH₃-phenyl)C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	CH_3C(=O)-	tetrahydrofuranyl (structure)

182

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(ring)
917	(2-CF₃-phenyl)-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(lactone ring, CH₃/H₃C)
918	(4-CF₃-phenyl)-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(lactone ring, CH₃/H₃C)
919	(4-CH₃O-phenyl)-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(lactone ring, CH₃/H₃C)
920	(2-CH₃O-phenyl)-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(lactone ring, CH₃/H₃C)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
921		-H	$-CH_2CH(CH_3)_2$	-H		
922		-H	$-CH_2CH(CH_3)_2$	-H		
923		-H	$-CH_2CH(CH_3)_2$	-H		
924		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A
925	2-acetylthiophene (thienyl–C(=O)CH$_3$)	-H	$-CH_2CH(CH_3)_2$	-H	CH$_3$–C=O (acetyl)	(tetrahydrofuranyl ring bearing CH$_3$, H$_3$C, and O$^-$)
926	H_3C–S(=O)$_2$–	-H	$-CH_2CH(CH_3)_2$	-H	CH$_3$–C=O (acetyl)	(tetrahydrofuranyl ring bearing CH$_3$, H$_3$C, and O$^-$)
927	$(H_3C)_2$CH–S(=O)$_2$–	-H	$-CH_2CH(CH_3)_2$	-H	CH$_3$–C=O (acetyl)	(tetrahydrofuranyl ring bearing CH$_3$, H$_3$C, and O$^-$)
928	C$_6$H$_5$–S(=O)$_2$– (phenylsulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	CH$_3$–C=O (acetyl)	(tetrahydrofuranyl ring bearing CH$_3$, H$_3$C, and O$^-$)

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A
929	F-phenyl-SO_2- (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C=O	(tetrahydrofuran, CH_3, H_3C) O-
930	Cl-phenyl-SO_2- (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C=O	(tetrahydrofuran, CH_3, H_3C) O-
931	Br-phenyl-SO_2- (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C=O	(tetrahydrofuran, CH_3, H_3C) O-
932	H_3C-phenyl-SO_2- (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-C=O	(tetrahydrofuran, CH_3, H_3C) O-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
933	2,4,6-trimethylphenyl-SO_2-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydrofuranyl structure)
934	4-tert-butylphenyl-SO_2-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydrofuranyl structure)
935	4-methoxyphenyl-SO_2-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydrofuranyl structure)
936	3-nitrophenyl-SO_2-	-H	-$CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydrofuranyl structure)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	A
937	![pyridine-2-sulfonyl structure]	-H	-CH$_2$CH(CH$_3$)$_2$	-H	![acetyl CH$_3$C=O]	![structure A]
938	![pyridine-3-sulfonyl structure]	-H	-CH$_2$CH(CH$_3$)$_2$	-H	![acetyl CH$_3$C=O]	![structure A]
939	![pyridine-4-sulfonyl structure]	-H	-CH$_2$CH(CH$_3$)$_2$	-H	![acetyl CH$_3$C=O]	![structure A]
940	![furan-2-sulfonyl structure]	-H	-CH$_2$CH(CH$_3$)$_2$	-H	![acetyl CH$_3$C=O]	![structure A]

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A–O–
941	H–	–H	–CH$_2$–(phenyl)	–H	CH$_3$C(=O)–	(structure)
942	(t-butyl acetate structure)	–H	–CH$_2$–(phenyl)	–H	CH$_3$C(=O)–	(structure)
943	(benzyl acetate structure)	–H	–CH$_2$–(phenyl)	–H	CH$_3$C(=O)–	(structure)
944	(4-F-benzyl acetate structure)	–H	–CH$_2$–(phenyl)	–H	CH$_3$C(=O)–	(structure)

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
945	(4-Cl-phenyl-CH$_2$-O-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(H$_3$C, CH$_3$ tetrahydrofuranyl-O-)
946	(4-H$_3$C-phenyl-CH$_2$-O-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(H$_3$C, CH$_3$ tetrahydrofuranyl-O-)
947	(4-CH$_3$O-phenyl-CH$_2$-O-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(H$_3$C, CH$_3$ tetrahydrofuranyl-O-)
948	(phenyl-CH$_2$-CH$_2$-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(H$_3$C, CH$_3$ tetrahydrofuranyl-O-)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
949	(phenoxy)CH₂C(=O)CH₃	-H	-CH₂-(phenyl)	-H	CH₃C(=O)-	(tetrahydropyranyl) CH₃, H₃C, O⁻
950	CH₃O-(phenyl)-O-CH₂C(=O)CH₃	-H	-CH₂-(phenyl)	-H	CH₃C(=O)-	H₃C / CH₃ tetrahydrofuran O⁻
951	CH₃O-(phenyl, OCH₃)-O-CH₂C(=O)CH₃	-H	-CH₂-(phenyl)	-H	CH₃C(=O)-	H₃C / CH₃ tetrahydrofuran O⁻
952	(phenyl)-C(=O)-CH₃	-H	-CH₂-(phenyl)	-H	CH₃C(=O)-	H₃C / CH₃ tetrahydrofuran O⁻

191

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A
953		-H	-CH$_2$	-H		
954		-H	-CH$_2$	-H		
955		-H	-CH$_2$	-H		
956		-H	-CH$_2$	-H		

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
957	H_3C-SO_2- (methanesulfonyl)	-H	$-CH_2-C_6H_5$	-H	CH_3-CO- (acetyl)	tetrahydrofuranyl-O⁻ (with CH_3, H_3C)
958	phenyl-SO_2- (benzenesulfonyl)	-H	$-CH_2-C_6H_5$	-H	CH_3-CO-	tetrahydrofuranyl-O⁻ (with CH_3, H_3C)
959	(4-F-phenyl)-SO_2-	-H	$-CH_2-C_6H_5$	-H	CH_3-CO-	tetrahydrofuranyl-O⁻ (with CH_3, H_3C)
960	(4-CH_3O-phenyl)-SO_2-	-H	$-CH_2-C_6H_5$	-H	CH_3-CO-	tetrahydrofuranyl-O⁻ (with CH_3, H_3C)

193

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
961	(benzyl acetate group)	-H	naphthalenyl-CH₂ (1-naphthylmethyl)	-H	CH₃–C(=O)–	(tetrahydrofuran ring with H_3C, CH_3, CH_3, O⁻)
962	(phenyl sulfonyl group, O=S=O)	-H	naphthalenyl-CH₂ (2-naphthylmethyl)	-H	CH₃–C(=O)–	(tetrahydrofuran ring with H_3C, CH_3, CH_3, O⁻)
963	(benzyl acetate group)	-H	$-CH_2CH_2SCH_3$	-H	CH₃–C(=O)–	(tetrahydrofuran ring with H_3C, CH_3, CH_3, O⁻)
964	(phenyl sulfonyl group, O=S=O)	-H	$-CH_2CH_2SCH_3$	-H	CH₃–C(=O)–	(tetrahydrofuran ring with H_3C, CH_3, CH_3, O⁻)

194

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
965	benzyl –CH₂–O–C(=O)CH₃	-H	$-CH_2OC(CH_3)_3$	-H	–C(=O)CH₃	(dimethyl tetrahydro ring, H₃C, CH₃, O–)
966	phenyl –S(=O)₂–	-H	$-CH_2OC(CH_3)_3$	-H	–C(=O)CH₃	(dimethyl tetrahydro ring, H₃C, CH₃, O–)
967	benzyl –CH₂–O–C(=O)CH₃	-H	$-CH_2OH$	-H	–C(=O)CH₃	(dimethyl tetrahydro ring, H₃C, CH₃, O–)
968	phenyl –S(=O)₂–	-H	$-CH_2OH$	-H	–C(=O)CH₃	(dimethyl tetrahydro ring, H₃C, CH₃, O–)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A ring)
969	benzyl acetate (PhCH₂–O–C(=O)–CH₃)	–H	phenyl (benzene)	–H	CH₃–C(=O)–	2-methyl-tetrahydropyran-type, (with A ring) –O⁻
970	phenyl sulfonyl (O=S(=O)–Ph)	–H	phenyl (benzene)	–H	CH₃–C(=O)–	3,3-dimethyl-tetrahydrofuran (H₃C, CH₃) –O⁻
971	benzyl acetate (PhCH₂–O–C(=O)–CH₃)	–H	–H	–H	CH₃–C(=O)–	3,3-dimethyl-tetrahydrofuran (H₃C, CH₃) –O⁻
972	phenyl sulfonyl (O=S(=O)–Ph)	–H	–H	–H	CH₃–C(=O)–	2-methyl-tetrahydropyran –O⁻

196

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
973	benzyl –O–CO–CH₃ (phenyl-CH₂–O–C(=O)–CH₃)	-H	-CH₃	-H	–C(=O)CH₃	pyran ring –O⁻ (methyl-substituted)
974	phenyl–S(=O)₂– (O=S=O)	-H	-CH₃	-H	–C(=O)CH₃	pyran ring –O⁻ (methyl-substituted)
975	benzyl –O–CO–CH₃ (phenyl-CH₂–O–C(=O)–CH₃)	-H	-CH₂CH₂CH₃	-H	–C(=O)CH₃	pyran ring –O⁻ (methyl-substituted)
976	phenyl–S(=O)₂– (O=S=O)	-H	-CH₂CH₂CH₃	-H	–C(=O)CH₃	pyran ring –O⁻ (methyl-substituted)

197

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
977	benzyl acetate structure	-H	-CH(CH₃)₂	-H	CH₃ C=O	pyranyl-O⁻
978	phenyl sulfonyl structure (O=S=O)	-H	-CH(CH₃)₂	-H	CH₃ C=O	pyranyl-O⁻
979	benzyl acetate structure	-H	-CH₂CH₂CH₂CH₃	-H	CH₃ C=O	pyranyl-O⁻
980	phenyl sulfonyl structure (O=S=O)	-H	-CH₂CH₂CH₂CH₃	-H	CH₃ C=O	pyranyl-O⁻

198

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	A O-O-
981	H-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$ C=O	O-O- structure
982	(CH$_3$)$_3$C-O-C(=O) (H$_3$C,H$_3$C,H$_3$C)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$ C=O	O-O- structure
983	benzyl-O-C(=O)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$ C=O	O-O- structure
984	4-F-benzyl-O-C(=O)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$ C=O	O-O- structure

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(A/O structure)
985	Cl—C₆H₄—CH₂—O—C(=O)—CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	
986	Br—C₆H₄—CH₂—O—C(=O)—CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	
987	H₃C—C₆H₄—CH₂—O—C(=O)—CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	
988	CH₃O—C₆H₄—CH₂—O—C(=O)—CH₃	-H	-CH₂CH(CH₃)₂	-H	CH₃—C(=O)—	

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
989	H$_3$C–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	
990	(CH$_3$)$_2$CH–CH$_2$–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	
991	C$_6$H$_5$–CH$_2$–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	
992	C$_6$H$_5$–CH$_2$CH$_2$–C(=O)–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	$\begin{array}{c} A \\ \text{(structure)} \end{array}$
993	phenoxy acetone structure	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 C=O	structure (O, O⁻, CH₃)
994	4-fluorophenoxy acetone structure (F)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 C=O	structure (O, O⁻, CH₃)
995	4-chlorophenoxy acetone structure (Cl)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 C=O	structure (O, O⁻, CH₃)
996	4-methylphenoxy acetone structure (H₃C)	-H	$-CH_2CH(CH_3)_2$	-H	CH_3 C=O	structure (O, O⁻, CH₃)

202

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
997	(2-OCH₃, 4-CH₃O phenyl-O-CH₂-CO-)	-H	-CH₂CH(CH₃)₂	-H	CH₃-CO-	(ring, 3-methyl tetrahydropyran, A-O-O-)
998	(phenyl-NH-CO-CH₃)	-H	-CH₂CH(CH₃)₂	-H	CH₃-CO-	(3-methyl tetrahydropyran-O-)
999	(phenyl-CO-CH₃)	-H	-CH₂CH(CH₃)₂	-H	CH₃-CO-	(3-methyl tetrahydropyran-O-)
1000	(2-F-phenyl-CO-CH₃)	-H	-CH₂CH(CH₃)₂	-H	CH₃-CO-	(3-methyl tetrahydropyran-O-)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1001		-H	-CH₂CH(CH₃)₂	-H		
1002		-H	-CH₂CH(CH₃)₂	-H		
1003		-H	-CH₂CH(CH₃)₂	-H		
1004		-H	-CH₂CH(CH₃)₂	-H		

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(ring)
1005	(2-methylphenyl)–C(=O)–CH$_3$	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	(A) tetrahydropyran-2-yl-O– with CH$_3$
1006	(4-methylphenyl, H$_3$C–)–C(=O)–CH$_3$	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran-2-yl-O– with CH$_3$
1007	(2-CF$_3$-phenyl)–C(=O)–CH$_3$	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran-2-yl-O– with CH$_3$
1008	(4-CF$_3$-phenyl, F$_3$C–)–C(=O)–CH$_3$	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$–C(=O)–	tetrahydropyran-2-yl-O– with CH$_3$

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	$\overset{A}{\langle}\!\!\!-\!\!O,\ -O^-$
1009	CH_3O—(4-methoxyphenyl)acetyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3—C(=O)	tetrahydropyranyl (3-methyl), $-O^-$
1010	(2-methoxyphenyl)acetyl, CH_3O	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3—C(=O)	tetrahydropyranyl (3-methyl), $-O^-$
1011	(pyridin-2-yl)acetyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3—C(=O)	tetrahydropyranyl (3-methyl), $-O^-$
1012	(pyridin-3-yl)acetyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3—C(=O)	tetrahydropyranyl (3-methyl), $-O^-$

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1013	4-(acetyl)pyridine	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	methyl-tetrahydropyranyl-O- (A)
1014	2-(acetyl)furan	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	methyl-dihydropyranyl-O-
1015	2-(acetyl)thiophene	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	methyl-dihydropyranyl-O-
1016	$H_3C-S(=O)(=O)-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	methyl-dihydropyranyl-O-

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
1017	$(H_3C)_2CH-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	
1018	phenyl-$S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	
1019	4-F-phenyl-$S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	
1020	4-Cl-phenyl-$S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
1021	Br–⟨C$_6$H$_4$⟩–SO$_2$–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$COCH$_3$ (acetyl)	tetrahydropyranyl-O–
1022	H$_3$C–⟨C$_6$H$_4$⟩–SO$_2$–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$COCH$_3$ (acetyl)	tetrahydropyranyl-O–
1023	H$_3$C,CH$_3$,CH$_3$–⟨C$_6$H$_2$⟩–SO$_2$–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$COCH$_3$ (acetyl)	tetrahydropyranyl-O–
1024	(H$_3$C)$_3$C–⟨C$_6$H$_4$⟩–SO$_2$–	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$COCH$_3$ (acetyl)	tetrahydropyranyl-O–

209

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	$\left(\!\begin{smallmatrix}A\end{smallmatrix}\!\right)$
1025	CH_3O—⟨phenyl⟩—SO_2	—H	$-CH_2CH(CH_3)_2$	—H	CH_3—CO—	⟨ring⟩—O⁻
1026	O_2N—⟨phenyl⟩—SO_2	—H	$-CH_2CH(CH_3)_2$	—H	CH_3—CO—	⟨ring⟩—O⁻
1027	⟨pyridyl⟩—SO_2	—H	$-CH_2CH(CH_3)_2$	—H	CH_3—CO—	⟨ring⟩—O⁻
1028	⟨pyridyl⟩—SO_2	—H	$-CH_2CH(CH_3)_2$	—H	CH_3—CO—	⟨ring⟩—O⁻

Table 2 (Continued)

(structure)	R7	R6	R5	R4	R1	Compound No.
(pyran ring, -O-)	CH$_3$-C(=O)-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	pyridine-SO$_2$-	1029
(pyran ring, -O-)	CH$_3$-C(=O)-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	furan-SO$_2$-	1030
(pyran ring, -O-)	CH$_3$-C(=O)-	-H	-CH$_2$- (phenyl)	-H	H-	1031
(pyran ring, -O-)	CH$_3$-C(=O)-	-H	-CH$_2$- (phenyl)	-H	(CH$_3$)$_3$C-O-C(=O)-	1032

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure)
1033	benzyl acetate (CH₂–O–C(=O)CH₃, phenyl)	–H	–CH₂–(phenyl)	–H	acetyl (CH₃–C(=O)–)	3-methyltetrahydropyran-2-yl–O⁻
1034	4-F-benzyl acetate	–H	–CH₂–(phenyl)	–H	acetyl (CH₃–C(=O)–)	3-methyltetrahydropyran-2-yl–O⁻
1035	4-Cl-benzyl acetate	–H	–CH₂–(phenyl)	–H	acetyl (CH₃–C(=O)–)	3-methyltetrahydropyran-2-yl–O⁻
1036	4-CH₃-benzyl acetate	–H	–CH₂–(phenyl)	–H	acetyl (CH₃–C(=O)–)	3-methyltetrahydropyran-2-yl–O⁻

212

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	[ring structure]
1037	(CH$_3$O-phenyl-CH$_2$-O-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(A ring with O, O-)
1038	(phenyl-CH$_2$CH$_2$-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(tetrahydropyran, CH$_3$, O-)
1039	(phenyl-O-CH$_2$-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(tetrahydropyran, CH$_3$, O-)
1040	(CH$_3$O-phenyl-O-CH$_2$-C(=O)-CH$_3$)	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(tetrahydropyran, CH$_3$, O-)

Table 2　(Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1041	(structure: CH₃O-substituted phenyl with OCH₃ and O-CH₂-C(=O)- group)	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	(pyran ring with CH₃ and O⁻)
1042	(phenyl with C(=O)CH₃)	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	(pyran ring with CH₃ and O⁻)
1043	(fluoro-substituted phenyl with C(=O)CH₃)	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	(pyran ring with CH₃ and O⁻)
1044	(fluoro-substituted phenyl with C(=O)CH₃)	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	(pyran ring with CH₃ and O⁻)

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1045	(2,6-diacetylphenyl)	-H	-CH₂(phenyl)	-H	CH₃C(=O)-	tetrahydropyran-2-yl-methyl-O-
1046	(4-trifluoromethyl-acetylphenyl)	-H	-CH₂(phenyl)	-H	CH₃C(=O)-	tetrahydropyran-2-yl-methyl-O-
1047	H₃C–S(=O)(=O)–	-H	-CH₂(phenyl)	-H	CH₃C(=O)-	tetrahydropyran-2-yl-methyl-O-
1048	(phenyl)-S(=O)(=O)–	-H	-CH₂(phenyl)	-H	CH₃C(=O)-	tetrahydropyran-2-yl-methyl-O-

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1049	4-F-C₆H₄-SO₂-	-H	-CH₂-C₆H₅	-H	CH₃-CO-	(methyl-tetrahydropyran-2-yl)-O⁻
1050	4-CH₃O-C₆H₄-SO₂-	-H	-CH₂-C₆H₅	-H	CH₃-CO-	(methyl-tetrahydropyran-2-yl)-O⁻
1051	C₆H₅-CH₂-O-CO-	-H	-CH₂-(1-naphthyl)	-H	CH₃-CO-	(methyl-tetrahydropyran-2-yl)-O⁻
1052	C₆H₅-SO₂-	-H	-CH₂-(2-naphthyl)	-H	CH₃-CO-	(methyl-tetrahydropyran-2-yl)-O⁻

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1053	benzyl ester	-H	$-CH_2CH_2SCH_3$	-H	CH_3-C(=O)-	pyranyl-O⁻
1054	phenyl sulfonyl	-H	$-CH_2CH_2SCH_3$	-H	CH_3-C(=O)-	pyranyl-O⁻
1055	benzyl ester	-H	$-CH_2OC(CH_3)_3$	-H	CH_3-C(=O)-	pyranyl-O⁻
1056	phenyl sulfonyl	-H	$-CH_2OC(CH_3)_3$	-H	CH_3-C(=O)-	pyranyl-O⁻

217

Table 2   (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	(structure with A—O⁻)
1057	benzyl –O–C(=O)–CH₃ (phenyl–CH₂–O–acetyl)	–H	–CH₂OH	–H	CH₃–C(=O)–	pyranyl ring with CH₃, –O⁻ (A ring)
1058	phenyl–S(=O)(=O)–	–H	–CH₂OH	–H	CH₃–C(=O)–	pyranyl ring with CH₃, –O⁻
1059	benzyl –O–C(=O)–CH₃ (phenyl–CH₂–O–acetyl)	–H	methylphenyl	–H	CH₃–C(=O)–	pyranyl ring with CH₃, –O⁻
1060	phenyl–S(=O)(=O)–	–H	methylphenyl	–H	CH₃–C(=O)–	pyranyl ring with CH₃, –O⁻

Table 2   (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
1061	benzyl acetate (PhCH₂–O–CO–CH₃)	H–	$-CH_2CH(CH_3)_2$	CH_3CO-	CH_3CO	tetrahydropyranyl-O⁻ (A ring)
1062	benzyl acetate (PhCH₂–O–CO–CH₃)	CH_3CO-	$-CH_2CH(CH_3)_2$	H–	CH_3CO	methyl-tetrahydropyranyl-O⁻
1063	phenyl sulfonyl (O=S=O, Ph)	CH_3CO-	$-CH_2CH(CH_3)_2$	H–	CH_3CO	methyl-tetrahydropyranyl-O⁻
1064	4-fluorophenyl sulfonyl (O=S=O, F–C₆H₄)	CH_3CO-	$-CH_2CH(CH_3)_2$	H–	CH_3CO	methyl-tetrahydropyranyl-O⁻

Table 2   (Continued)

	R⁷	R⁶	R⁵	R⁴	R¹	Compound No.
(structure, contains A)	—	—	—	—	—	—
(tetrahydropyran-2-yl, 3-methyl, -O-)	CH_3–C(=O)–	H–	$-CH_2CH(CH_3)_2$	CH_3CO-	H_3C–(phenyl)–$S(=O)_2$–	1065
(tetrahydropyran-2-yl, 3-methyl, -O-)	CH_3–C(=O)–	H–	$-CH_2CH(CH_3)_2$	CH_3CO-	CH_3O–(phenyl)–$S(=O)_2$–	1066
(tetrahydropyran-2-yl, 3-methyl, -O-)	CH_3–C(=O)–	H–	$-CH_2$(phenyl)	CH_3CO-	(phenyl)CH_2O–C(=O)–	1067
(tetrahydropyran-2-yl, 3-methyl, -O-)	CH_3–C(=O)–	H–	$-CH_2$(phenyl)	CH_3CO-	(phenyl)–$S(=O)_2$–	1068

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	(structure)
1069	(benzyl acetate structure)	CH$_3$CO–	–CH$_2$CH(CH$_3$)$_2$	H–	CH$_3$ (acetyl)	
1070	(phenyl sulfone structure)	CH$_3$CO–	–CH$_2$CH(CH$_3$)$_2$	H–	CH$_3$ (acetyl)	
1389	(benzyl acetate structure)	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$	
1390	(benzyl acetate structure)	–H	–CH$_2$CH(CH$_3$)$_2$	–H	CH$_3$ CH$_3$ CH$_3$	

Table 2 (Continued)

Compound No.	R¹	R⁴	R⁵	R⁶	R⁷	
1391		-H	$-CH_2CH(CH_3)_2$	-H		
1392		-H	$-CH_2CH(CH_3)_2$	-H		
1393		-H	$-CH_2CH(CH_3)_2$	-H		
1394		-H	$-CH_2CH(CH_3)_2$	-H		

Table 2 (Continued)

Compound No.	R^1	R^4	R^5	R^6	R^7	
1395	(2,4,6-trimethylphenyl ketone: CH₃C(O)–, H₃C, H₃C substituents)	–H	–CH₂CH(CH₃)₂	–H	(CH₃)₃C–C(O)–CH₃ (pivaloyl-type)	(oxepane, –O–)
1396	4-(tert-butyl)phenyl sulfone (O=S=O, (CH₃)₃C)	–H	–CH₂CH(CH₃)₂	–H	CH₃CH₂C(O)–	(tetrahydropyran, –O–)
1397	4-(tert-butyl)phenyl sulfone (O=S=O, (CH₃)₃C)	–H	–CH₂CH(CH₃)₂	–H	(CH₃)₃C–C(O)–CH₃	(tetrahydrofuran, –O–)
1398	4-(tert-butyl)phenyl sulfone (O=S=O, (CH₃)₃C)	–H	–CH₂CH(CH₃)₂	–H	C₆H₅C(O)–	(tetrahydropyran, –O–)

Table 3   (where n = 0)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	Structure
1071	benzyl acetate	-H	-CH₂CH(CH₃)₂	-H	-H	-H	oxetane-OH
1072	benzyl acetate	-H	-CH₂CH(CH₃)₂	-H	-CH₃	-H	oxetane-OH
1073	benzyl acetate	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂CH₃	-H	oxetane-OH
1074	benzyl acetate	-H	-CH₂CH(CH₃)₂	-H	-CH(CH₃)₂	-H	oxetane-OH

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1075	CH₃-C(=O)-O-CH₂-C₆H₅ (benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂CH₂CH₃	-H	ring with OH
1076	CH₃-C(=O)-O-CH₂-C₆H₅ (benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	ring with OH
1077	C₆H₅-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	ring with OH
1078	C₆H₅-S(=O)(=O)-	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	ring with OH

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1079	benzyl $-CH_2-O-C(=O)-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2-$ (benzyl)	-H	oxetane–OH
1080	benzyl $-CH_2-O-C(=O)-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2SCH_3$	-H	oxetane–OH
1081	benzyl $-CH_2-O-C(=O)-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2OH$	-H	oxetane–OH
1082	benzyl $-CH_2-O-C(=O)-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	phenyl	-H	oxetane–OH

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	Structure
1083	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	-H	-H	(tetrahydrofuran-OH, methyl)
1084	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_3$	-H	(tetrahydrofuran-OH, methyl)
1085	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_3$	-H	(tetrahydrofuran-OH, methyl)
1086	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_3$	-H	(tetrahydrofuran-OH, methyl)

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	structure
1087	phenyl–CH₂–O–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)_2$	-H	ring (A, O, OH, CH₃)
1088	phenyl–CH₂–O–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_2CH_3$	-H	ring (O, OH, CH₃)
1089	phenyl–CH₂–O–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)(CH_2CH_3)$	-H	ring (O, OH, CH₃)
1090	phenyl–CH₂–O–C(=O)–CH₃	-H	-H	-H	$-CH_2CH(CH_3)_2$	-H	ring (O, OH, CH₃)

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	
1091	PhCH₂-O-C(=O)-CH₃	-H	$-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran-OH with A)
1092	PhCH₂-O-C(=O)-CH₃	-H	$-CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran-OH, methyl)
1093	PhCH₂-O-C(=O)-CH₃	-H	$-CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran-OH, methyl)
1094	PhCH₂-O-C(=O)-CH₃	-H	$-CH_2CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydrofuran-OH, methyl)

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	Structure
1095	-H	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1096		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1097		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1098		$-CH_3$	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	(structure)
1099	benzyl acetate	-H	$-CH_2CH(CH_3)_2$	$-CH_3$	$-CH_2CH(CH_3)_2$	-H	2-hydroxy-3-methyltetrahydropyran (A)
1100	benzyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	$-CH_3$	2-hydroxy-3-methyltetrahydrofuran
1101	4-fluorobenzyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	2-hydroxy-3-methyltetrahydrofuran
1102	4-chlorobenzyl acetate	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	2-hydroxy-3-methyltetrahydrofuran

EP 0 810 221 A1

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	Structure
1107	$H_3C-C(=O)-$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(ring with A, O, OH, CH₃)
1108	$CH_3(CH_2)_{12}-C(=O)-$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydropyran, OH, CH₃)
1109	$C_6H_5CH_2CH_2-C(=O)-$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydropyran, OH, CH₃)
1110	$C_6H_5-O-CH_2-C(=O)-$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	(tetrahydropyran, OH, CH₃)

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	(ring structure with O, OH, CH₃)
1111	2-(CH_3-O-CH₂-C(=O)- substituted phenoxy) OCH_3	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	A ring, O, OH
1112	CH_3O / OCH_3 substituted phenoxy-CH₂-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	O ring, OH
1113	phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	O ring, OH
1114	2-F-phenyl-C(=O)-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	O ring, OH

Table 3   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	A O OH
1115		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1116		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1117		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1118		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1119	(structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(structure, A)
1120	(structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(structure)
1121	(structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(structure)
1122	(structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(structure)

236

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1123	(2-acetylfuran structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, A ring, OH, CH₃)
1124	(2-acetyl-4H-chromen-4-one structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃)
1125	$H_3C-S(=O)_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃)
1126	(phenyl-$S(=O)_2-$)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃)

237

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1127	4-F-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	3-methyltetrahydrofuran-2-ol
1128	4-Cl-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	3-methyltetrahydrofuran-2-ol
1129	4-CH₃-C₆H₄-SO₂-	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	3-methyltetrahydrofuran-2-ol
1130	2,4,6-(CH₃)₃-C₆H₂-SO₂-	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	3-methyltetrahydrofuran-2-ol

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	Structure
1131	![4-methoxyphenyl sulfonyl]	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	tetrahydropyran-OH, methyl
1132	![2-nitrophenyl sulfonyl]	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	tetrahydrofuran-OH, methyl
1133	![1-naphthyl sulfonyl]	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	tetrahydrofuran-OH, methyl
1134	![2-naphthyl sulfonyl]	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	tetrahydrofuran-OH, methyl

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	structure
1135	pyridyl-SO_2-	-H	-$CH_2CH(CH_3)_2$	-H	-$CH_2CH(CH_3)_2$	-H	tetrahydropyran (methyl, OH)
1136	quinolinyl-SO_2-	-H	-$CH_2CH(CH_3)_2$	-H	-$CH_2CH(CH_3)_2$	-H	tetrahydropyran (methyl, OH)
1137	benzyl-$O-C(=O)$-	-H	-CH_2(phenyl)	-H	-$CH_2CH(CH_3)_2$	-H	tetrahydropyran (methyl, OH)
1138	benzyl-$O-C(=O)$-	-H	-$CH_2CH_2SCH_3$	-H	-$CH_2CH(CH_3)_2$	-H	tetrahydropyran (methyl, OH)

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	
1139	(benzyl ester) CH₂–O–C(=O)CH₃	-H	$-CH_2OC(CH_3)_3$	-H	$-CH_2CH(CH_3)_2$	-H	(methyl-tetrahydropyranyl–OH)
1140	(benzyl ester) CH₂–O–C(=O)CH₃	-H	$-CH_2OH$	-H	$-CH_2CH(CH_3)_2$	-H	(methyl-tetrahydropyranyl–OH)
1141	(benzyl ester) CH₂–O–C(=O)CH₃	-H	(phenyl)	-H	$-CH_2CH(CH_3)_2$	-H	(methyl-tetrahydropyranyl–OH)
1142	(benzyl ester) CH₂–O–C(=O)CH₃	-H	-H	-H	$-CH_2-$ (benzyl)	-H	(methyl-tetrahydropyranyl–OH)

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	
1143	(benzyl acetate)	-H	-CH$_3$	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuranyl-OH, 3-methyl)
1144	(benzyl acetate)	-H	-CH$_2$CH$_2$CH$_3$	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuranyl-OH, 3-methyl)
1145	(benzyl acetate)	-H	-CH(CH$_3$)$_2$	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuranyl-OH, 3-methyl)
1146	(benzyl acetate)	-H	-CH$_2$CH$_2$CH$_2$CH$_3$	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuranyl-OH, 3-methyl)

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(ring structure)
1147	PhCH₂–O–C(=O)–CH₃	–H	–CH(CH₃)CH₂CH₃	–H	–CH₂–C₆H₅	–H	tetrahydrofuran-OH, CH₃
1148	PhCH₂–O–C(=O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–CH₂–C₆H₅	–H	tetrahydrofuran-OH, CH₃
1149	PhCH₂–O–C(=O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–CH₂–C₆H₅	–H	tetrahydrofuran-OH, CH₃
1150	C₆H₅–C(=O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–CH₂–C₆H₅	–H	tetrahydrofuran-OH, CH₃

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	(ring structure)
1151	phenyl-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuran ring with OH and CH$_3$)
1152	benzyl-O-C(=O)-	-H	-CH$_2$CH$_2$SCH$_3$	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuran ring with OH and CH$_3$)
1153	benzyl-O-C(=O)-	-H	-CH$_2$OH	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuran ring with OH and CH$_3$)
1154	benzyl-O-C(=O)-	-H	(phenyl)	-H	-CH$_2$-(phenyl)	-H	(tetrahydrofuran ring with OH and CH$_3$)

Table 3   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	(structure)
1155	PhCH$_2$-O-C(=O)- (benzyl ester)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$CH$_2$SCH$_3$	-H	tetrahydropyran-2-ol, 3-methyl (A ring)
1156	Ph-S(=O)(=O)-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$CH$_2$SCH$_3$	-H	tetrahydropyran-2-ol, 3-methyl
1157	PhCH$_2$-O-C(=O)- (benzyl ester)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$OH	-H	tetrahydrofuran-2-ol, 3-methyl
1158	Ph-S(=O)(=O)-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$OH	-H	tetrahydrofuran-2-ol, 3-methyl

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	A
1159	(benzyl ester structure)	-H	$-CH_2CH(CH_3)_2$	-H	(phenyl)	-H	(structure, OH)
1160	(phenylsulfonyl structure)	-H	$-CH_2CH(CH_3)_2$	-H	(phenyl)	-H	(structure, OH)
1161	(benzyl ester structure)	-H	$-CH_2CH(CH_3)_2$	-H	-H	-H	(structure, CH_3, OH)
1162	(benzyl ester structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_3$	-H	(structure, CH_3, OH)

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(A ring structure with $-OH$)
1163	benzyl acetate	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH_2CH_3$	$-H$	CH_3 ring, $-OH$
1164	benzyl acetate	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH(CH_3)_2$	$-H$	CH_3 ring, $-OH$
1165	benzyl acetate	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH_2CH_2CH_3$	$-H$	CH_3 ring, $-OH$
1166	benzyl acetate	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3 ring, $-OH$

247

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1167	phenyl ketone (C₆H₅–CO–)	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH(CH₃)₂	–H	tetrahydrofuran ring with H₃C, CH₃, OH
1168	phenyl sulfonyl (C₆H₅–SO₂–)	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH(CH₃)₂	–H	tetrahydrofuran ring with H₃C, CH₃, OH
1169	benzyl acetate ester	–H	–CH₂CH(CH₃)₂	–H	–CH₂– (benzyl)	–H	tetrahydrofuran ring with H₃C, CH₃, OH
1170	benzyl acetate ester	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH₂SCH₃	–H	tetrahydrofuran ring with H₃C, CH₃, OH

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1171	benzyl-O-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂OH	-H	(ring with CH₃, OH)
1172	benzyl-O-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	phenyl	-H	(ring with H₃C, CH₃, OH)
1173	benzyl-O-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	-H	-H	(pyran ring with OH)
1174	benzyl-O-C(=O)-CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₃	-H	(pyran ring with CH₃, OH)

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	
1175	⌬–CH₂–O–C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_3$	-H	(pyranol structure, A)
1176	⌬–CH₂–O–C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_3$	-H	(pyranol structure)
1177	⌬–CH₂–O–C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)_2$	-H	(pyranol structure)
1178	⌬–CH₂–O–C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_2CH_3$	-H	(pyranol structure)

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	
1179		-H	-H	-H	$-CH_2CH(CH_3)_2$	-H	
1180		-H	$-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	
1181		-H	$-CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	
1182		-H	$-CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	

251

Table 3   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	Structure
1183	benzyl–CH_2–O–C(=O)–CH_3	-H	-$CH(CH_3)_2$	-H	-$CH_2CH(CH_3)_2$	-H	pyran ring with methyl and OH
1184	benzyl–CH_2–O–C(=O)–CH_3	-H	-$CH_2CH_2CH_2CH_3$	-H	-$CH_2CH(CH_3)_2$	-H	pyran ring with methyl and OH
1185	H-	-H	-$CH_2CH(CH_3)_2$	-H	-$CH_2CH(CH_3)_2$	-H	pyran ring with methyl and OH
1186	$(H_3C)(H_3C)(H_3C)C$–O–C(=O)–CH_3	-H	-$CH_2CH(CH_3)_2$	-H	-$CH_2CH(CH_3)_2$	-H	pyran ring with methyl and OH

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	
1187		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1188		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1189		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1190		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1191		$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	
1192		$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	
1193		$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	
1194		$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	

Table 3   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	
1195		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1196		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1197		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	
1198		-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	Structure
1199	H_3C-SO_2- (methanesulfonyl)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	pyranyl-OH with CH₃
1200	phenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	pyranyl-OH with CH₃
1201	4-fluorophenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	pyranyl-OH with CH₃
1202	2,6-dimethyl-4-methylphenyl-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	pyranyl-OH with CH₃

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1203	(benzyl-O-C(=O)-CH₃)	-H	-CH₂-(phenyl)	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃, A)
1204	(benzyl-O-C(=O)-CH₃)	-H	$-CH_2CH_2SCH_3$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃)
1205	(benzyl-O-C(=O)-CH₃)	-H	$-CH_2OC(CH_3)_3$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃)
1206	(benzyl-O-C(=O)-CH₃)	-H	$-CH_2OH$	-H	$-CH_2CH(CH_3)_2$	-H	(tetrahydropyran, OH, CH₃)

Table 3   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	(structure with A, O, OH, CH₃)
1207	benzyl acetate ($-CH_2-O-C(=O)CH_3$)	$-H$	methylcyclohexenyl	$-H$	$-CH_2CH(CH_3)_2$	$-H$	tetrahydropyran, $-OH$, $-CH_3$
1208	benzyl acetate ($-CH_2-O-C(=O)CH_3$)	$-H$	$-H$	$-H$	$-CH_2-$phenyl	$-H$	tetrahydropyran, $-OH$, $-CH_3$
1209	benzyl acetate ($-CH_2-O-C(=O)CH_3$)	$-H$	$-CH_3$	$-H$	$-CH_2-$phenyl	$-H$	tetrahydropyran, $-OH$, $-CH_3$
1210	benzyl acetate ($-CH_2-O-C(=O)CH_3$)	$-H$	$-CH_2CH_2CH_3$	$-H$	$-CH_2-$phenyl	$-H$	tetrahydropyran, $-OH$, $-CH_3$

Table 3 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^G	(structure)
1211	benzyl –O–C(=O)CH$_3$	–H	–CH(CH$_3$)$_2$	–H	–CH$_2$–C$_6$H$_5$	–H	tetrahydropyran with CH$_3$ and –OH
1212	benzyl –O–C(=O)CH$_3$	–H	–CH$_2$CH$_2$CH$_2$CH$_3$	–H	–CH$_2$–C$_6$H$_5$	–H	tetrahydropyran with CH$_3$ and –OH
1213	benzyl –O–C(=O)CH$_3$	–H	–CH(CH$_3$)(CH$_2$CH$_3$)	–H	–CH$_2$–C$_6$H$_5$	–H	tetrahydropyran with CH$_3$ and –OH
1214	benzyl –O–C(=O)CH$_3$	–H	–CH$_2$CH(CH$_3$)$_2$	–H	–CH$_2$–C$_6$H$_5$	–H	tetrahydropyran with CH$_3$ and –OH

Table 3   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	
1215	C₆H₅CH₂–O–C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2$–C₆H₅	-H	(tetrahydropyran ring, A, O, OH, CH₃)
1216	C₆H₅–C(=O)CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2$–C₆H₅	-H	(tetrahydropyran ring, O, OH, CH₃)
1217	C₆H₅–S(=O)₂–	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2$–C₆H₅	-H	(tetrahydropyran ring, O, OH, CH₃)
1218	C₆H₅CH₂–O–C(=O)CH₃	-H	$-CH_2CH_2SCH_3$	-H	$-CH_2$–C₆H₅	-H	(tetrahydropyran ring, O, OH, CH₃)

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1219	benzyl acetate (–CH₂O–C(=O)CH₃ on phenyl)	-H	-CH₂OH	-H	-CH₂-phenyl	-H	tetrahydropyran, OH, methyl (A)
1220	benzyl acetate (–CH₂O–C(=O)CH₃ on phenyl)	-H	cyclohexenyl	-H	-CH₂-phenyl	-H	tetrahydropyran, OH, methyl
1221	benzyl acetate (–CH₂O–C(=O)CH₃ on phenyl)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂SCH₃	-H	tetrahydropyran, OH, methyl
1222	phenylsulfonyl (phenyl–S(=O)(=O)–)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂SCH₃	-H	tetrahydropyran, OH, methyl

Table 3 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	(structure)
1223	benzyl–O–C(=O)–CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2OH$	$-H$	(2-hydroxy-3-methyltetrahydropyran)
1224	phenyl–S(=O)(=O)–	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2OH$	$-H$	(2-hydroxy-3-methyltetrahydropyran)
1225	benzyl–O–C(=O)–CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	phenyl	$-H$	(2-hydroxy-3-methyltetrahydropyran)
1226	phenyl–S(=O)(=O)–	$-H$	$-CH_2CH(CH_3)_2$	$-H$	phenyl	$-H$	(2-hydroxy-3-methyltetrahydropyran)

Table 4 (where n = 0)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	A
1227	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	-H	-H	(acetyl, CH_3-C=O)	(oxetane structure)
1228	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_3$	-H	(acetyl, CH_3-C=O)	(oxetane structure)
1229	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_3$	-H	(acetyl, CH_3-C=O)	(oxetane structure)
1230	(benzyl acetate structure)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)_2$	-H	(acetyl, CH_3-C=O)	(oxetane structure)

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	
1231	benzyl –O–C(=O)–CH₃ (phenyl-CH₂–O–C(=O)–)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_2CH_3$	-H	CH₃–C(=O)–	(oxetane ring with –O–)
1232	benzyl –O–C(=O)–CH₃ (phenyl-CH₂–O–C(=O)–)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	CH₃–C(=O)–	(oxetane ring with –O–)
1233	phenyl–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	CH₃–C(=O)–	(oxetane ring with –O–)
1234	phenyl–S(=O)(=O)–	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	CH₃–C(=O)–	(oxetane ring with –O–)

264

Table 4   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	(ring)
1235	CH$_3$C(=O)O-CH$_2$-(phenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$-(phenyl)	-H	CH$_3$-C(=O)-	(oxo ring, -O-)
1236	CH$_3$C(=O)O-CH$_2$-(phenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$CH$_2$SCH$_3$	-H	CH$_3$-C(=O)-	(oxo ring, -O-)
1237	CH$_3$C(=O)O-CH$_2$-(phenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$OH	-H	CH$_3$-C(=O)-	(oxo ring, -O-)
1238	CH$_3$C(=O)O-CH$_2$-(phenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(phenyl)	-H	CH$_3$-C(=O)-	(oxo ring, -O-)

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1239	$CH_3C(=O)O\text{-}CH_2\text{-}C_6H_5$	-H	$-CH_2CH(CH_3)_2$	-H	-H	-H	$CH_3C(=O)-$	tetrahydropyranyl-methyl-O-
1240	$CH_3C(=O)O\text{-}CH_2\text{-}C_6H_5$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_3$	-H	$CH_3C(=O)-$	tetrahydropyranyl-methyl-O-
1241	$CH_3C(=O)O\text{-}CH_2\text{-}C_6H_5$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_3$	-H	$CH_3C(=O)-$	tetrahydropyranyl-methyl-O-
1242	$CH_3C(=O)O\text{-}CH_2\text{-}C_6H_5$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_3$	-H	$CH_3C(=O)-$	tetrahydropyranyl-methyl-O-

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	(ring)
1243	PhCH₂–O–CO–CH₃ (benzyl acetate)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)_2$	-H	CH₃–CO–	tetrahydropyranyl (A, O, O–)
1244	PhCH₂–O–CO–CH₃ (benzyl acetate)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_2CH_3$	-H	CH₃–CO–	tetrahydropyranyl (O, O–)
1245	PhCH₂–O–CO–CH₃ (benzyl acetate)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)(CH_2CH_3)$	-H	CH₃–CO–	tetrahydropyranyl (O, O–)
1246	PhCH₂–O–CO–CH₃ (benzyl acetate)	-H	-H	-H	$-CH_2CH(CH_3)_2$	-H	CH₃–CO–	tetrahydropyranyl (O, O–)

Table 4　(Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1247	$PhCH_2-O-CO-CH_3$	-H	$-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyranyl-O-)
1248	$PhCH_2-O-CO-CH_3$	-H	$-CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyranyl-O-)
1249	$PhCH_2-O-CO-CH_3$	-H	$-CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyranyl-O-)
1250	$PhCH_2-O-CO-CH_3$	-H	$-CH_2CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	CH_3-CO-	(tetrahydropyranyl-O-)

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(structure)
1251	H–	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(structure)
1252	$H_3C-C(CH_3)_2-O-C(=O)-CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(structure)
1253	benzyl–O–C(=O)–CH₃	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(structure)
1254	benzyl–O–C(=O)–CH₃	$-CH_3$	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(structure)

Table 4 (Continued)

Compound No.	R1	R2	R3	R4	R5	R6	R7	(ring)
1255	benzyl–O–C(=O)–CH3	-H	-CH2CH(CH3)2	-CH3	-CH2CH(CH3)2	-H	CH3–C(=O)–	(A ring, O–O⁻)
1256	benzyl–O–C(=O)–CH3	-H	-CH2CH(CH3)2	-H	-CH2CH(CH3)2	-CH3	CH3–C(=O)–	(ring, O–O⁻)
1257	4-F-benzyl–O–C(=O)–CH3	-H	-CH2CH(CH3)2	-H	-CH2CH(CH3)2	-H	CH3–C(=O)–	(ring, O–O⁻)
1258	4-Cl-benzyl–O–C(=O)–CH3	-H	-CH2CH(CH3)2	-H	-CH2CH(CH3)2	-H	CH3–C(=O)–	(ring, O–O⁻)

Table 4 (Continued)

Structure	R^7	R^6	R^5	R^4	R^3	R^2	R^1	Compound No.
(tetrahydropyran, methyl, —O—)	CH_3—C(=O)—	—H	—$CH_2CH(CH_3)_2$	—H	—$CH_2CH(CH_3)_2$	—H	4-CH_3-benzyl acetoxymethyl (H_3C—phenyl—CH$_2$—O—C(=O))	1259
(tetrahydropyran, methyl, —O—)	CH_3—C(=O)—	—H	—$CH_2CH(CH_3)_2$	—H	—$CH_2CH(CH_3)_2$	—H	4-CH_3O-benzyl acetoxymethyl (CH_3O—phenyl—CH$_2$—O—C(=O))	1260
(tetrahydropyran, methyl, —O—)	CH_3—C(=O)—	—H	—$CH_2CH(CH_3)_2$	—H	—$CH_2CH(CH_3)_2$	—H	fluorenyl acetoxymethyl	1261
(tetrahydropyran, methyl, —O—)	CH_3—C(=O)—	—H	—$CH_2CH(CH_3)_2$	—H	—$CH_2CH(CH_3)_2$	—H	pyridyl acetoxymethyl	1262

271

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1263	H_3C–C(=O)–	–H	$-CH_2CH(CH_3)_2$	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-methyl –O–
1264	$CH_3(CH_2)_{12}$–C(=O)–	–H	$-CH_2CH(CH_3)_2$	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-methyl –O–
1265	phenyl–CH₂CH₂–C(=O)–CH₃	–H	$-CH_2CH(CH_3)_2$	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-methyl –O–
1266	phenyl–O–CH₂–C(=O)–CH₃	–H	$-CH_2CH(CH_3)_2$	–H	$-CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyran-methyl –O–

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1267	(2-methoxyphenoxy)acetyl, OCH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)	tetrahydropyran-2-yl-O-
1268	(4-methoxy-2-methoxyphenoxy)acetyl, OCH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)	tetrahydropyran-2-yl-O-
1269	phenyl-C(=O)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)	tetrahydropyran-2-yl-O-
1270	(2-fluorophenyl)-C(=O), F	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)	tetrahydropyran-2-yl-O-

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1271		-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H		
1272		-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H		
1273		-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H		
1274		-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H		

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	(ring)
1275	2-methoxyphenyl acetyl (CH₃O-substituted acetophenone)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	acetyl ($CH_3C{=}O$)	(tetrahydropyran ring with A, methyl, O^-)
1276	(OCH₃-substituted methoxyphenyl acetyl)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	acetyl ($CH_3C{=}O$)	(tetrahydropyran ring, methyl, O^-)
1277	(tert-butyl–oxy phenyl acetyl)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	acetyl ($CH_3C{=}O$)	(tetrahydropyran ring, methyl, O^-)
1278	(pyridyl acetyl)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	acetyl ($CH_3C{=}O$)	(tetrahydropyran ring, methyl, O^-)

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1279	2-acetyl-furan	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	(ring structure)
1280	2-acetyl-4-oxo-chromene	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	(ring structure)
1281	H_3C-SO_2-	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	(ring structure)
1282	$C_6H_5-SO_2-$	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3CH_2C(=O)-$	(ring structure)

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1283	phenyl-SO_2 (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3CH_2C(=O)CH_2-$	tetrahydropyran ring with -O-
1284	phenyl-SO_2 (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$(CH_3)_2CHC(=O)CH_3$	tetrahydropyran ring with -O-
1285	phenyl-SO_2 (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$(CH_3)_3CC(=O)CH_3$	tetrahydropyran ring with -O-
1286	phenyl-SO_2 (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$C_6H_5C(=O)CH_3$	tetrahydropyran ring with -O-

Table 4   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	(ring)
1287	phenyl–SO_2–	–H	–$CH_2CH(CH_3)_2$	–H	–$CH_2CH(CH_3)_2$	–H	–CH_3	tetrahydropyranyl-methyl–O–
1288	phenyl–SO_2–	–H	–$CH_2CH(CH_3)_2$	–H	–$CH_2CH(CH_3)_2$	–H	–$C(CH_3)_3$	tetrahydropyranyl-methyl–O–
1289	4-F-phenyl–SO_2–	–H	–$CH_2CH(CH_3)_2$	–H	–$CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyranyl-methyl–O–
1290	4-Cl-phenyl–SO_2–	–H	–$CH_2CH(CH_3)_2$	–H	–$CH_2CH(CH_3)_2$	–H	CH_3–C(=O)–	tetrahydropyranyl-methyl–O–

278

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1291	4-methylphenyl-SO₂ (H₃C–C₆H₄–SO₂–)	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring with CH₃ and –O–)
1292	2,4,6-trimethylphenyl-SO₂ (H₃C, CH₃, CH₃ substituted)	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring with CH₃ and –O–)
1293	4-methoxyphenyl-SO₂ (CH₃O–C₆H₄–SO₂–)	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring with CH₃ and –O–)
1294	3-nitrophenyl-SO₂ (O₂N–C₆H₄–SO₂–)	–H	–CH₂CH(CH₃)₂	–H	–CH₂CH(CH₃)₂	–H	CH₃–C(=O)–	(tetrahydropyran ring with CH₃ and –O–)

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(structure)
								(A ring, O-)
1295	(naphthalene-1-sulfonyl, O=S=O)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-O-, CH₃)
1296	(naphthalene-2-sulfonyl, O=S=O)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-O-, CH₃)
1297	(quinoline-sulfonyl, O=S=O)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-O-, CH₃)
1298	(quinoline-5-sulfonyl, O=S=O)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(tetrahydropyran-O-, CH₃)

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1299	$-CH_2-C_6H_5$–OC(=O)CH$_3$	-H	-CH$_2$- (phenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	ring with O (A)
1300	$-CH_2-C_6H_5$–OC(=O)CH$_3$	-H	-CH$_2$CH$_2$SCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	ring with O
1301	$-CH_2-C_6H_5$–OC(=O)CH$_3$	-H	-CH$_2$OC(CH$_3$)$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	ring with O
1302	$-CH_2-C_6H_5$–OC(=O)CH$_3$	-H	-CH$_2$OH	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	ring with O

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1303	benzyl–O–C(=O)CH₃	-H	phenyl	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(=O)-$	tetrahydropyranyl–O– (2-methyl)
1304	benzyl–O–C(=O)CH₃	-H	-H	-H	$-CH_2$–phenyl	-H	$CH_3C(=O)-$	tetrahydropyranyl–O– (2-methyl)
1305	benzyl–O–C(=O)CH₃	-H	$-CH_3$	-H	$-CH_2$–phenyl	-H	$CH_3C(=O)-$	tetrahydropyranyl–O– (2-methyl)
1306	benzyl–O–C(=O)CH₃	-H	$-CH_2CH_2CH_3$	-H	$-CH_2$–phenyl	-H	$CH_3C(=O)-$	tetrahydropyranyl–O– (2-methyl)

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	A
1307	[benzyl acetate]	-H	$-CH(CH_3)_2$	-H	$-CH_2$ (phenyl)	-H	$C(=O)CH_3$	[ring structure]
1308	[benzyl acetate]	-H	$-CH_2CH_2CH_2CH_3$	-H	$-CH_2$ (phenyl)	-H	$C(=O)CH_3$	[ring structure]
1309	[benzyl acetate]	-H	$-CH(CH_3)(CH_2CH_3)$	-H	$-CH_2$ (phenyl)	-H	$C(=O)CH_3$	[ring structure]
1310	[benzyl acetate]	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2$ (phenyl)	-H	$C(=O)CH_3$	[ring structure]

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring structure)
1311	benzyl –CH₂–O–C(=O)–CH₃	–H	–CH₂CH(CH₃)₂	–H	–CH₂–C₆H₅	–H	CH₃–C(=O)–	tetrahydropyran-2-yl-O–, 3-methyl
1312	phenyl –C(=O)CH₃ (acetophenone)	–H	–CH₂CH(CH₃)₂	–H	–CH₂–C₆H₅	–H	CH₃–C(=O)–	tetrahydropyran-2-yl-O–, 3-methyl
1313	phenyl–SO₂–	–H	–CH₂CH(CH₃)₂	–H	–CH₂–C₆H₅	–H	CH₃–C(=O)–	tetrahydropyran-2-yl-O–, 3-methyl
1314	benzyl –CH₂–O–C(=O)–CH₃	–H	–CH₂CH₂SCH₃	–H	–CH₂–C₆H₅	–H	CH₃–C(=O)–	tetrahydropyran-2-yl-O–, 3-methyl

284

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1315	benzyl acetate	-H	-CH₂OH	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	ring
1316	benzyl acetate	-H	cyclohexenyl	-H	-CH₂-C₆H₅	-H	CH₃-C(=O)-	ring
1317	benzyl acetate	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂SCH₃	-H	CH₃-C(=O)-	ring
1318	phenyl-SO₂-	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂SCH₃	-H	CH₃-C(=O)-	ring

285

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	$\overset{A}{\frown}$ structure
1319	benzyl–O–C(=O)CH₃ (phenyl-CH₂-O-acetyl)	-H	-CH₂CH(CH₃)₂	-H	-CH₂OH	-H	-C(=O)CH₃	tetrahydropyran ring (3-methyl, 2-O–)
1320	phenyl–S(=O)(=O)–	-H	-CH₂CH(CH₃)₂	-H	-CH₂OH	-H	-C(=O)CH₃	tetrahydropyran ring (3-methyl, 2-O–)
1321	benzyl–O–C(=O)CH₃ (phenyl-CH₂-O-acetyl)	-H	-CH₂CH(CH₃)₂	-H	phenyl	-H	-C(=O)CH₃	tetrahydropyran ring (3-methyl, 2-O–)
1322	phenyl–S(=O)(=O)–	-H	-CH₂CH(CH₃)₂	-H	phenyl	-H	-C(=O)CH₃	tetrahydropyran ring (3-methyl, 2-O–)

Table 4   (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	A
1323	(benzyl acetate: $C_6H_5CH_2$–O–CO–CH_3)	-H	$-CH_2CH(CH_3)_2$	-H	-H	-H	CH_3–CO–	(ring with CH_3) –O–
1324	(benzyl acetate: $C_6H_5CH_2$–O–CO–CH_3)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_3$	-H	CH_3–CO–	(ring with CH_3) –O–
1325	(benzyl acetate: $C_6H_5CH_2$–O–CO–CH_3)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2CH_3$	-H	CH_3–CO–	(ring with CH_3) –O–
1326	(benzyl acetate: $C_6H_5CH_2$–O–CO–CH_3)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH(CH_3)_2$	-H	CH_3–CO–	(ring with CH_3) –O–

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1327	(benzyl-O-C(=O)CH₃)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂CH₂CH₃	-H	CH₃-C(=O)-	(ring with O, A, O-)
1328	(benzyl-O-C(=O)CH₃)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(CH₃ ring, O, O-)
1329	(phenyl-C(=O)CH₃)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(CH₃ ring, O, O-)
1330	(phenyl-S(=O)(=O)-)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃-C(=O)-	(H₃C, CH₃ ring, O, O-)

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(structure)
1331	benzyl–O–C(=O)CH₃ (benzyloxy-acetyl)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2$–C₆H₅	$-H$	CH_3–C(=O)–	H_3C, CH_3 tetrahydrofuran O–O–
1332	benzyl–O–C(=O)CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH_2SCH_3$	$-H$	CH_3–C(=O)–	H_3C, CH_3 tetrahydrofuran O–O–
1333	benzyl–O–C(=O)CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2OH$	$-H$	CH_3–C(=O)–	H_3C, CH_3 tetrahydrofuran O–O–
1334	benzyl–O–C(=O)CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	C₆H₅ (phenyl)	$-H$	CH_3–C(=O)–	H_3C, CH_3 tetrahydrofuran O–O–

Table 4 (Continued)

Compound No.	R1	R2	R3	R4	R5	R6	R7	(A)O—O-
1335	benzyl–O–C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	-H	-H	CH₃–C(=O)–	methyl-pyranyl-O-
1336	benzyl–O–C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₃	-H	CH₃–C(=O)–	methyl-pyranyl-O-
1337	benzyl–O–C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₃	-H	CH₃–C(=O)–	methyl-pyranyl-O-
1338	benzyl–O–C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH₂CH₃	-H	CH₃–C(=O)–	methyl-pyranyl-O-

Table 4 (Continued)

Compound No.	R1	R2	R3	R4	R5	R6	R7	(structure)
1339	benzyl –CH2–O–CO–CH3	–H	–CH2CH(CH3)2	–H	–CH(CH3)2	–H	CH3–CO–	pyran (A)
1340	benzyl –CH2–O–CO–CH3	–H	–CH2CH(CH3)2	–H	–CH2CH2CH2CH3	–H	CH3–CO–	pyran
1341	benzyl –CH2–O–CO–CH3	–H	–H	–H	–CH2CH(CH3)2	–H	CH3–CO–	pyran
1342	benzyl –CH2–O–CO–CH3	–H	–CH3	–H	–CH2CH(CH3)2	–H	CH3–CO–	pyran

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(A)
1343	benzyl acetate ester	-H	$-CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(O)-$	pyran ring with methyl, O-
1344	benzyl acetate ester	-H	$-CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(O)-$	pyran ring with methyl, O-
1345	benzyl acetate ester	-H	$-CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(O)-$	pyran ring with methyl, O-
1346	benzyl acetate ester	-H	$-CH_2CH_2CH_2CH_3$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3C(O)-$	pyran ring with methyl, O-

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1347	-H	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	(pyranyl, A)-O⁻
1348	$(CH_3)_3C-C(=O)-O-$ type	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	pyranyl-O⁻
1349	benzyl acetate type	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	pyranyl-O⁻
1350	F-substituted benzyl acetate type	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH(CH_3)_2$	-H	$CH_3-C(=O)-$	pyranyl-O⁻

293

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(A structure)
1351	(structure: CH₃O–benzyl acetate)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃–C=O (acetyl)	(methyl-pyran O⁻)
1352	(structure: fluorenylmethyl acetate)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃–C=O (acetyl)	(methyl-pyran O⁻)
1353	(structure: H₃C–C=O, acetonyl)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃–C=O (acetyl)	(methyl-pyran O⁻)
1354	(structure: phenyl-CH₂CH₂–C(=O)CH₃)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃–C=O (acetyl)	(methyl-pyran O⁻)

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1355	phenoxy-CH₂-C(=O)CH₃ (C₆H₅O–CH₂–CO–CH₃)	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	tetrahydropyranyl (methyl-substituted) –O–
1356	CH₃O–(aryl)(OCH₃)–O–CH₂–C(=O)CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	tetrahydropyranyl (methyl-substituted) –O–
1357	phenyl–C(=O)CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	tetrahydropyranyl (methyl-substituted) –O–
1358	(F-substituted phenyl)–C(=O)CH₃	$-H$	$-CH_2CH(CH_3)_2$	$-H$	$-CH_2CH(CH_3)_2$	$-H$	CH_3–C(=O)–	tetrahydropyranyl (methyl-substituted) –O–

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	Structure
1359	4-CH₃-C₆H₄-C(=O)CH₃ (H₃C–phenyl–acetyl)	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	tetrahydropyran-O⁻ with methyl (ring A)
1360	2-OCH₃-C₆H₄-C(=O)CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	tetrahydropyran-O⁻ with methyl
1361	H₃C–S(=O)₂–	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	tetrahydropyran-O⁻ with methyl
1362	C₆H₅–S(=O)₂–	-H	-CH₂CH(CH₃)₂	-H	-CH₂CH(CH₃)₂	-H	CH₃C(=O)-	tetrahydropyran-O⁻ with methyl

EP 0 810 221 A1

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	
1363	4-F-C$_6$H$_4$-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	(pyranyl)-O-
1364	2,4,6-(CH$_3$)$_3$-C$_6$H$_2$-SO$_2$-	-H	-CH$_2$CH(CH$_3$)$_2$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	(pyranyl)-O-
1365	C$_6$H$_5$-CH$_2$-O-C(=O)CH$_3$	-H	-CH$_2$(C$_6$H$_5$)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	(pyranyl)-O-
1366	C$_6$H$_5$-CH$_2$-O-C(=O)CH$_3$	-H	-CH$_2$CH$_2$SCH$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$-C(=O)-	(pyranyl)-O-

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	(structure)
1367	benzyl–CH$_2$OC(=O)CH$_3$	-H	-CH$_2$OC(CH$_3$)$_3$	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	pyranyl (CH$_3$) –O-
1368	benzyl–CH$_2$OC(=O)CH$_3$	-H	-CH$_2$OH	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	pyranyl (CH$_3$) –O-
1369	benzyl–CH$_2$OC(=O)CH$_3$	-H	(methylphenyl)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	CH$_3$–C(=O)–	pyranyl (CH$_3$) –O-
1370	benzyl–CH$_2$OC(=O)CH$_3$	-H	-H	-H	-CH$_2$–(phenyl)	-H	CH$_3$–C(=O)–	pyranyl (CH$_3$) –O-

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring)
1371	benzyl-OC(=O)CH₃	-H	-CH₃	-H	-CH₂-phenyl	-H	CH₃C(=O)	(A) ring
1372	benzyl-OC(=O)CH₃	-H	-CH₂CH₂CH₃	-H	-CH₂-phenyl	-H	CH₃C(=O)	tetrahydropyran ring
1373	benzyl-OC(=O)CH₃	-H	-CH(CH₃)₂	-H	-CH₂-phenyl	-H	CH₃C(=O)	tetrahydropyran ring
1374	benzyl-OC(=O)CH₃	-H	-CH₂CH₂CH₂CH₃	-H	-CH₂-phenyl	-H	CH₃C(=O)	tetrahydropyran ring

Table 4   (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	(ring A)
1375	benzyl–CH₂–O–C(=O)–CH₃	-H	-CH(CH₃)CH₂CH₃	-H	-CH₂–(phenyl)	-H	CH₃–C(=O)–	(A-ring pyran with O⁻, CH₃)
1376	benzyl–CH₂–O–C(=O)–CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂–(phenyl)	-H	CH₃–C(=O)–	pyran O⁻, CH₃
1377	benzyl–CH₂–O–C(=O)–CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂–(phenyl)	-H	CH₃–C(=O)–	pyran O⁻, CH₃
1378	phenyl–C(=O)–CH₃	-H	-CH₂CH(CH₃)₂	-H	-CH₂–(phenyl)	-H	CH₃–C(=O)–	pyran O⁻, CH₃

Table 4 (Continued)

Compound No.	R¹	R²	R³	R⁴	R⁵	R⁶	R⁷	
1379	(phenyl)-SO₂-	-H	-CH₂CH(CH₃)₂	-H	-CH₂-(phenyl)	-H	CH₃-CO-	(pyran ring with A, CH₃, O⁻)
1380	(phenyl)-CH₂-O-CO-CH₃	-H	-CH₂CH₂SCH₃	-H	-CH₂-(phenyl)	-H	CH₃-CO-	(pyran ring, CH₃, O⁻)
1381	(phenyl)-CH₂-O-CO-CH₃	-H	-CH₂OH	-H	-CH₂-(phenyl)	-H	CH₃-CO-	(pyran ring, CH₃, O⁻)
1382	(phenyl)-CH₂-O-CO-CH₃	-H	(cyclohexenyl)	-H	-CH₂-(phenyl)	-H	CH₃-CO-	(pyran ring, CH₃, O⁻)

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	(ring structure)
1383	benzyl ester (—CH₂—O—C(=O)—)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2SCH_3$	-H	$-C(=O)CH_3$	pyran ring, —O⁻
1384	phenyl sulfonyl (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2CH_2SCH_3$	-H	$-C(=O)CH_3$	pyran ring, —O⁻
1385	benzyl ester (—CH₂—O—C(=O)—)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2OH$	-H	$-C(=O)CH_3$	pyran ring, —O⁻
1386	phenyl sulfonyl (O=S=O)	-H	$-CH_2CH(CH_3)_2$	-H	$-CH_2OH$	-H	$-C(=O)CH_3$	pyran ring, —O⁻

Table 4 (Continued)

Compound No.	R^1	R^2	R^3	R^4	R^5	R^6	R^7	
1387	(benzyl ester structure)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(phenyl)	-H	(CH$_3$ acetyl)	(oxygen heterocyclic structure)
1388	(phenyl sulfonyl structure)	-H	-CH$_2$CH(CH$_3$)$_2$	-H	(phenyl)	-H	(CH$_3$ acetyl)	(oxygen heterocyclic structure)

Method for preparation of the compounds of the present invention will be explained. The oxygen-containing heterocyclic derivatives represented by the above formula (I) can be prepared, for example, according to the method described below.

Preparation m thod 1: Preparation of the compounds wherein R^7 is a hydrogen atom

(IV)

(V)

(VI)

(II)

In the above formulas, R^1, R^2, R^3, R^4, R^5, R^6, A, and n have the same meanings as those defined above.

An amino acid derivative represented by the formula (IV) is reacted with a condensing agent such as dicyclohexylcarbodiimide, diphenylphosphoryl azide, carbonyldiimidazole, oxalyl chloride, isobutyl chloroformate, or thionyl chloride optionally in the presence of a base such as triethylamine or pyridine to activate the carboxylic acid. A compound represented by the above formula (VI) can be obtained by reacting the above-obtained product with a lactone derivative represented by the above formula (V). A solvent used for this condensation reaction may be appropriately chosen so as to be suitable for a condensing agent used, and reaction conditions or other may also be applied so as to be suitable for a condensing agent used. The oxygen-containing heterocyclic derivative represented by the formula (II) can be obtained by treating the above-obtained compound of the formula (VI) with a reducing agent such as diisobutyl aluminium hydride or sodium borohydride/cerium chloride.

Preparation method 2: Preparation of the compounds wherein R^7 is R^9-CO-

(II)

(VII)

In the above formulas, R^1, R^2, R^3, R^4, R^5, R^6, R^9, A, and n have the same meanings as those defined above.

The oxygen-containing heterocyclic derivative represented by the above formula (VII) can be obtained by dissolving the oxygen-containing heterocyclic derivative produced by Preparation method 1 in an organic solvent such as methylene chloride, 1,2-dichloroethane, dimethylformamide, N-methylpyrrolidone, tetrahydrofuran, ethyl acetate, acetonitrile, or toluene, and then the solution is allowed to react with an acid anhydride represented by the formula of (R^{9}CO)$_2$O in the presence of a base such as pyridine, triethylamine, or 4-dimethylaminopyridine. This reaction can also be carried out without a solvent.

Preparation method 3: Preparation of the compounds wherein R^7 is a C$_1$-C$_5$ alkyl group

(VII)

(VIII)

In the above formulas, R^1, R^2, R^3, R^4, R^5, R^6, R^9, A, and n have the same meaning as those defined above and R^{10} represents a C_1-C_5 alkyl group.

The compound represented by the above formula (VIII) can be obtained by dissolving the compound of the formula (VII) obtained in Preparation method 2 in an alcohol compound represented by the formula $R^{10}OH$, and then adding a catalytic amount of an acid such as hydrochloric acid or sulfuric acid and stirring the mixture.

In the above-mentioned series of reaction steps, some reactions may require protection and deprotection of one or more functional groups. For that purpose, any protective group suitable for the functional group may be chosen, and any known method described in literatures may be applied as manufacturing procedures.

Among the oxygen-containing heterocyclic derivatives of the present invention obtained as described above, the compounds of the formula (II) wherein R^7 is hydrogen atom have potent inhibitory activity against a cysteine protease. The compounds of formula (VIII) having a C_1-C_5 alkyl group as R^7 and the compounds of formula (VII) having R^9-CO- (R^9 represents a C_1-C_{10} alkyl group or a C_6-C_{12} aryl group which may optionally be substituted) as R^7 can be used as pro-drugs of the oxygen-containing heterocyclic derivatives (II) that have potent inhibitory activity against a cysteine protease. More specifically, when the compound of formula (VII) or (VIII) is orally administered, the compound is absorbed from the intestinal tract or other, and then the oxygen-containing heterocyclic derivative of the formula (II) as an active form is rapidly released by the functions of enzymes and other in a living body.

(VII)

In a living body ⟶

(II)

(VIII)

When the compound of the present invention is clinically used, a ratio of the therapeutically useful ingredient based on one or more carrier components may vary in a range of 1-90% by weight. For example, the compounds of the present invention can be orally administered as formulations in the form of granules, fine granules, powders, hard capsules, soft capsules, syrups, emulsions, suspensions, liquid preparations or the like, or alternatively, administered intravenously, intramuscularly, or subcutaneously as injections. They may be used as suppositories. They may also be formulated as powders for injection and used as injections prepared before use. Organic or inorganic and solid or liquid carriers or diluents for the preparation of formulations suitable for oral, intestinal, or parenteral administration may be used for the preparation of the medicament of the present invention. For example, as excipients used for the preparation of solid formulations, lactose, sucrose, starch, talc, cellulose, dextrin, kaolin, calcium carbonate or other may be used. Liquid formulations for oral administration such as emulsions, syrups, suspensions, solutions or other may contain a conventional inert diluent such as water or vegetable oil. These pharmaceutical preparations may contain, in addition to the inert diluent, auxiliaries such as moistening agents, suspension aids, sweeteners, aromatics, colorants, or preservatives. The medicament may be formulated as a liquid preparation and filled in capsules made of an absorbable material such as gelatin. As solvents or suspension mediums used for formulations for parenteral administration, i.e., injection, suppositories and other, for example, propylene glycol, polyethylene glycol, benzyl alcohol, ethyl oleate, lecithin and the like may be used. As base materials used for suppositories, for example, cacao butter, emulsified cacao butt r, lauric lipid, witepsol and the lik may be used. The pharmaceutical pr parations may be prepared according to

ordinary methods.

Clinical dose may generally be in the range of 0.01-1,000 mg per day, when orally administered, for an adult as the weight of the compound of the pr sent invention . However, it is further preferred that the dose may appropriately be increased or decreased depending the age, conditions, and symptom of a patient. The daily dose of the medicament of the present invention may be administered once a day, or twice o or three times a day with appropriate intervals, or alternatively, administered intermittently.

When the medicament is used as an injection, it is desirable that 0.001-100 mg per day for an adult as the weight of the compound of the present invention is administered continuously or intermittently.

Brief Description of the Drawings

Figure 1 is a HPLC chart obtained after 5 minutes' incubation of the compound of Example 88 in rat serum.
Figure 2 is a HPLC chart of a sample obtained by dissolving the compound of Example 1 and the compound of Example 88 in the absence of serum.

Best Mode for Carrying Out the Invention

The present invention will be further detailed by referring to reference examples and examples. However, these reference examples and examples are non-limiting so far that the scope of the present invention falls within the gist of the present invention.

Reference Example 1: Preparation of (S)-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-tetrahydrofuranone

6 ml of thionyl chloride was cooled to -5°C, and the reagent was added with 998 mg of N-phenylsulfonyl-L-leucine. The reaction mixture was stirred at -5°C for 10 minutes, then warmed up to room temperature and stirring was further continued for 3 hours. The reaction mixture was then concentrated under reduced pressure, and the resulting residue was added with 10 ml of toluene and further concentrated to obtain crude N-phenylsulfonyl-L-leucyl chloride as a residue. The resulting crude N-phenylsulfonyl-L-leucyl chloride was dissolved in 20 ml of methylene chloride, and the solution was added with 443 mg of L-homoserinelactone hydrochloride and 0.946 ml of triethylamine under ice cooling. The reaction mixture was stirred for 15 minutes under ice cooling and then stirring was further continued for 1.5 hours at room temperature. After the completion of the reaction, the reaction mixture was added with diluted hydrochloric acid and extracted with methylene chloride. The extract was washed successively with water, saturated aqueous sodium hydrogencarbonate, and saturated brain, dried over magnesium sulfate, and then filtered. The filtrate was concentrated and the resulting residue was added with 10 ml of ethyl acetate and 20 ml of hexane and stirred. Precipitated crystals were collected by filtration to obtain the desired compound (861 mg).

Yield: 76%
Melting point: 183-184°C
IR (KBr, cm^{-1}): 3331, 3256, 1772, 1649.
NMR (CDCl$_3$, δ): 0.67 (d, J=6.0Hz, 3H), 0.84 (d, J=6.3Hz, 3H), 1.45-1.56 (m, 3H), 2.01 (m, 1H), 2.63 (m, 1H), 4.26 (m, 1H), 4.36 (m, 1H), 4.45 (ddd, J=9.3Hz, 9.3Hz, 1.8Hz, 1H), 5.28 (d, J=8.1 Hz, 1H), 6.67 (d, J=6.0Hz, 1H), 7.52 (m, 2H), 7.60 (m, 1H), 7.88 (dd, J=7.2Hz, 1.5Hz, 2H).

Example 1: Preparation of (3S)-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 196 in Table-1)

413 mg of (S)-3-((S)-2-phenylsulfonylamino-4-methylvalerylamino)-2-tetrahydrofuranone obtained in Reference Example 1 was dissolved in 60 ml of methylene chloride and the solution was cooled to -78 °C . 3.81 ml of a solution of diisobutylalminium hydride in toluene (1.01 mol/L) was added to the reaction solution. After three hours, the reaction mixture was added with a saturated aqueous solution of ammonium chloride and ethyl acetate, and then warmed up to room temperature and filtered through celite. The celite was washed thoroughly with ethyl acetate. The filtrate was washed with saturated brain, dried over magnesium sulfate, and then filtered. The filtrate was concentrated, and the resulting residue was purified by silica gel column chromatography (eluent: ethyl acetate containing 30% hexane) to give 191 mg of the desired compound.

Yield: 46%
Melting point: 162°C
IR (KBr, cm^{-1}): 3337, 3260, 1649.
NMR (CDCl$_3$+DMSO-d$_6$, δ): 0.72 (d, J=6.6Hz, 2.7H), 0.78 (d, J=6.3Hz, 0.3H), 0.84 (d, J=6.6Hz, 2.7H), 0.86 (d,

J=6.3Hz, 0.3H), 1.46 (t, J=7.2Hz, 2H), 1.63 (m, 2H), 2.09 (m, 0.9H), 2.25 (m, 0.1H), 3.68-3.81 (m, 2H), 3.99-4.12 (m, 2H), 4.95 (br s, 0.1H), 5.03 (d, J=3.6Hz, 0.1H), 5.15 (dd, J=3.9Hz, 3.9Hz, 0.9H), 5.63 (d, J=3.9Hz, 0.9H), 6.68 (d, J=9.3Hz, 0.1H), 6.81 (d, J=8.1Hz, 0.9H), 6.89 (d, J=7.8Hz, 0.9H), 7.14 (d, J=7.2Hz, 0.1H), 7.46-7.58 (m, 3H), 7.85 (m, 2H).

The ratio of the isomers in this solvent was about 9:1.

NMR (CD$_3$OD, δ): 0.74 (d, J=6.5Hz, 3H), 0.80 (d, J=6.5Hz, 3H), 0.86 (d, J=7.1Hz, 3H), 0.88 (d, J=6.8Hz, 3H), 1.34-1.50 (m, 3H), 1.64 (m, 1H), 2.01 (m, 0.4H), 2.17 (m, 0.6H), 3.73-3.97 (m, 4H), 4.96 (s, 0.6H), 5.11 (d, J=4Hz, 0.4H), 7.53-7.61 (m, 3H), 7.85 (m, 2H).

The ratio of the isomers in this solvent was about 6:4.

Compounds of Example 2 to Example 87 were prepared in the same manners as those of Reference Example 1 and Example 1. Physicochemical data of the compounds will be described below.

Example 2: Preparation of (3S)-3-benzyloxycarbonylaminoacethylamino-2-tetrahydrofuranol (Compound No. 17 in Table-1)

Melting point: 119-121°C
IR (KBr, cm^{-1}): 3314, 1692, 1649, 1541.
NMR (CDCl$_3$, δ): 1.84 (m, 1H), 2.22-2.50 (m, 1H), 2.86 (s, 0.3H), 2.97 (s, 0.7H), 3.79-4.00 (m, 3H), 4.11 (m, 1H), 4.37 (m, 1H), 5.14 (s, 2H), 5.27 (m, 1.3H), 5.39 (s, 0.7H), 6.12 (s, 0.3H), 6.42 (s, 0.7H), 7.36 (m, 5H).

Example 3: Preparation of (3S)-3-((S)-2-bezyloxycarbonylaminopropionylamino)-2-tetrahydrofuranol (Compound No. 18 in Table-1)

Melting point: 161-163°C
IR (KBr, cm^{-1}): 3312, 1688, 1647, 1561, 1530.
NMR (CDCl$_3$, δ): 1.36 (d, J=7.2Hz, 0.75H), 1.39 (d, 7.2Hz, 2.25H), 1.81 (m, 1H), 2.34 (m, 0.75H), 2.43 (m, 0.25H), 2.99 (s, 0.25H), 3.09 (s, 0.75H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.75H), 4.00 (m, 0.25H), 4.11 (m, 1H), 4.22 (m, 1H), 4.37 (m, 1H), 5.11 (s, 2H), 5.29 (m, 2H), 6.28 (s, 0.25H), 6.45 (s, 0.75H), 7.35 (m, 5H).

Example 4: Preparation of (3S)-3-((S)-2-benzyloxycarbonylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 20 in Table-1)

Melting point: 148-149°C
IR (KBr, cm^{-1}): 3299, 1694, 1645, 1539.
NMR (CDCl$_3$, δ): 0.92 (m, 3H), 1.38 (m, 2H), 1.62 (m, 1H), 1.78 (m, 2H), 2.31 (m, 0.7H), 2.42 (m, 0.3H), 3.27 (s, 0.3H), 3.42 (s, 0.7H), 3.86 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.7H), 3.98 (m, 0.3H), 4.11 (m, 2H), 4.33 (m, 1H), 5.10 (s, 2H), 5.27 (m, 1.3H), 5.38 (d, J=7.1Hz, 0.7H), 6.30 (s, 0.3H), 6.46 (d, J=8.0Hz, 0.7H), 7.35 (m, 5H).

Example 5: Preparation of 3-((S)-2-benzyloxycarbonylamino-3-methylbutyrylamino)-2-tetrahydrofuranol (Compound No. 21 in Table-1)

Melting point: 122-123°C
IR (KBr, cm^{-1}): 3302, 1694, 1647, 1537.
NMR (CDCl$_3$, δ): 0.94 (m, 6H), 1.83 (m, 1H), 2.12 (m, 1H), 2.30 (m, 0.6H), 2.44 (m, 0.4H), 3.40 (s, 0.4H), 3.49 (s, 0.6H), 3.82-4.06 (m, 2H), 4.10 (m, 1H), 4.38 (m, 1H), 5.09 (m, 2H), 5.26 (s, 0.4H), 5.32 (s, 0.6H), 5.50 (m, 1H), 6.32 (s, 0.2H), 6.45 (s, 0.4H), 6.54 (s, 0.2H), 6.68 (s, 0.2H), 7.34 (m, 5H).

Example 6: Preparation of (3S)-3-((S)-2-benzyloxycarbonylaminohexanoylamino)-2-tetrahydrofuranol (Compound No. 22 in Table-1)

Melting point: 165-166°C
IR (KBr, cm^{-1}): 3304, 1694, 1645, 1539.
NMR (CDCl$_3$, δ): 0.87 (m, 3H), 1.32 (m, 3H), 1.64 (m, 2H), 1.78 (m, 2H), 2.29 (m, 0.8H), 2.42 (m, 0.2H), 3.22 (s, 0.2H), 3.39 (s, 0.8H), 3.86 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.8H), 3.99 (m, 0.2H), 4.11 (m, 2H), 4.33 (m, 1H), 5.10 (s, 2H), 5.17-5.44 (m, 2H), 6.27 (s, 0.2H), 6.47 (d, J=7.8Hz, 0.8H), 7.34 (m, 5H).

Example 7: Preparation of (3S)-3-((2S)-2-benzyloxycarbonylamino-3-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 23 in Table-1)

Melting point: 169-171°C

IR (KBr, cm⁻¹): 3299, 1694, 1649, 1539.

NMR (CDCl₃, δ): 0.91 (m, 6H), 1.12 (m, 1H), 1.50 (m, 1H), 1.84 (m, 2H), 2.31 (m, 0.7H), 2.44 (m, 0.3H), 3.40 (s, 1H), 3.82-4.16 (m, 3H), 4.35 (m, 1H), 5.10 (s, 2H), 5.26 (m, 1H), 5.41 (s, 1H), 6.17 (s, 0.3H), 6.40 (s, 0.7H), 7.38 (m, 5H).

Example 8: Preparation of (3S)-3-((S)-2-amino-4-methylvalerylamino)-2-tetrahydrofuranol hydrochloride (Compound No. 24 in Table-1)

NMR (CD₃OD, δ): 1.00 (d, J=5.6Hz, 6H), 1.69-1.83 (m, 3H), 1.98 (m, 1H), 2.24 (m, 0.5H), 2.34 (m, 0.5H), 3.84-4.07 (m, 4H), 4.95 (s, 0.5H), 4.99 (d, J=4.0Hz, 0.5H).

Example 9: Preparation of 3-((S)-2-methoxycarbonylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 25 in Table-1)

Melting point: 139-141°C

IR (KBr, cm⁻¹): 3287, 3081, 1686, 1653, 1553.

NMR (CDCl₃, δ): 0.94 (m, 6H), 1.50-1.74 (m, 3H), 1.81 (m, 1H), 2.36 (m, 0.6H), 2.48 (m, 0.4H), 3.44 (s, 0.4H), 3.58 (s, 0.6H), 3.68 (s, 3H), 3.99 (m, 0.6H), 4.01 (m, 0.4H), 4.13 (m, 2H), 4.34 (m, 1H), 5.27 (d, J=3.0Hz, 0.6H), 5.34 (m, 1.4H), 6.49 (s, 0.4H), 6.57 (d, J=7.8Hz, 0.4H), 6.68 (s, 0.2H).

Example 10: Preparation of (3S)-3-((S)-2-tert-butoxycarbonylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 27 in Table-1)

Melting point: 65-70°C

IR (KBr, cm⁻¹): 3310, 1698, 1657.

NMR (CDCl₃, δ): 0.95 (m, 6H), 1.44 (s, 4.5H), 1.44 (s, 4.5H), 1.48 (m, 1H), 1.66 (m, 2H), 1.81 (m, 1H), 2.33 (m, 0.5H), 2.46 (m, 0.5H), 3.88 (ddd, J=7.5Hz, 7.5Hz, 7.5Hz, 0.5H), 4.01 (ddd, J=8.4Hz, 8.4Hz, 8.4Hz, 0.5H), 4.12 (m, 2H), 4.36 (m, 1H), 4.96 (d, J=7.8Hz, 0.5H), 5.03 (m, 0.5H), 5.26 (s, 0.5H), 5.33 (s, 0.5H), 6.47 (m, 0.5H), 6.61 (m, 0.5H).

Example 11: Preparation of (3S)-3-((S)-2-isobutoxycarbonylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 28 in Table-1)

Melting point: 31-33°C

IR (KBr, cm⁻¹): 3310, 1699, 1657, 1543.

NMR (CDCl₃, δ): 0.94 (m, 12H), 1.47-2.00 (m, 5H), 2.36 (m, 0.5H), 2.45 (m, 0.5H), 2.96 (s, 0.5H), 3.17 (s, 0.5H), 3.90 (m, 1.5H), 4.02 (m, 0.5H), 4.15 (m, 2H), 4.36 (m, 1H), 5.08 (m, 1H), 5.27 (s, 0.5H), 5.35 (s, 0.5H), 6.24 (s, 0.5H), 6.50 (s, 0.5H).

Example 12: Preparation of (3S)-3-((S)-2-cyclohexylmethoxycarbonylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 29 in Table-1)

Melting point: 52-54°C

IR (KBr, cm⁻¹): 3310, 1703, 1659, 1545.

NMR (CDCl₃, δ): 0.93 (m, 8H), 1.26-1.33 (m, 4H), 1.43-1.92 (m, 9H), 2.36 (m, 0.5H), 2.50 (m, 0.5H), 3.22 (s, 0.5H), 3.49 (s, 0.5H), 3.87-4.23 (m, 5H), 4.34 (m, 1H), 5.13 (m, 1H), 5.26 (d, J=2.7Hz, 0.5H), 5.32 (dd, J=3.9Hz, 3.9Hz, 0.5H), 5.34 (s, 0.5H), 6.53 (s, 0.5H).

Example 13: Preparation of (3S)-3-((S)-2-benzyloxycarbonylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 31 in Table-1)

Melting point: 40-43°C

IR (KBr, cm⁻¹): 3306, 1705, 1657.

NMR (CDCl₃, δ): 0.92 (d, J=6.1Hz, 3H), 0.94 (d, J=5.9Hz, 3H), 1.52 (m, 1H), 1.64 (m, 2H), 1.78 (m, 1H), 2.29 (m, 0.5H), 2.41 (m, 0.5H), 3.51 (s, 0.5H), 3.74 (s, 0.5H), 3.85 (ddd, J=8.0Hz, 8.0Hz, 8.0Hz, 0.5H), 3.97 (m, 0.5H), 4.10

(m, 2H), 4.32 (m, 1H), 5.09 (s, 1H), 5.10 (s, 1H), 5.24 (s, 0.5H), 5.29 (s, 0.5H), 5.35 (d, J=6.5Hz, 0.5H), 5.38 (d, J=8.2Hz, 0.5H), 6.45 (d, J=6.0Hz, 0.5H), 6.57 (d, J=6.0Hz, 0.5H), 7.33 (m, 5H).

Example 14: Preparation of (3S)-3-{(S)-2-(N-benzyloxycarbonyl-N-methyl)amino-4-methylvalerylamino}-2-tetrahydro-furanol (Compound No. 32 in Table-1)

IR (neat): 3335, 1669.
NMR (CDCl₃, δ): 0.93 (m, 6H), 1.49 (m, 1H), 1.69 (m, 3H), 2.28 (m, 0.5H), 2.39 (m, 0.5H), 2.85 (s, 1.5H), 2.86 (s, 1.5H), 3.12 (s, 0.5H), 3.30 (s, 0.5H), 3.85 (m, 1H), 4.07 (m, 1H), 4.29 (m, 1H), 4.60 (m, 0.5H), 4.70 (m, 0.5H), 5.09-5.26 (m, 3H), 6.24 (s, 0.5H), 6.53 (s, 0.5H), 7.36 (m, 5H).

Example 15: Preparation of (3S)-3-{(S)-2-(4-fluorobenzyloxycarbonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 37 in Table-1)

Melting point: 50-52°C
IR (KBr, cm⁻¹): 3310, 1705, 1657, 1607, 1541, 1514.
NMR (CDCl₃, δ): 0.93 (m, 6H), 1.47-1.90 (m, 4H), 2.32 (m, 0.5H), 2.44 (m, 0.5H), 3.10 (s, 0.5H), 3.35 (s, 0.5H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 3.98 (m, 0.5H), 4.11 (m, 2H), 4.33 (m, 1H), 5.04 (s, 1H), 5.06 (s, 1H), 5.25 (m, 2H), 6.20 (bs, 0.5H), 6.46 (d, J=8.1Hz, 0.5H), 7.03 (dd, J=8.7Hz, 2H), 7.33 (m, 2H).

Example 16: Preparation of (3S)-3-{(S)-2-(2-chlorobenzyloxycarbonylamino)-4-methylvalerylamino}-2-tetrahydrofura-nol (Compound No. 38 in Table-1)

Melting point: 46-49°C
IR (KBr, cm⁻¹): 3308, 1707, 1657, 1541.
NMR (CDCl₃, δ): 0.94 (m, 6H), 1.48-1.86 (m, 4H), 2.32 (m, 0.6H), 2.48 (m, 0.4H), 2.90 (s, 0.4H), 3.09 (s, 0.6H), 3.88 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.6H), 4.01 (m, 0.4H), 4.11 (m, 2H), 4.34 (m, 1H), 5.23 (s, 2H), 5.25 (m, 2H), 6.18 (s, 0.4H), 6.45 (d, J=7.6Hz, 0.6H), 7.26 (m, 2H), 7.40 (m, 2H).

Example 17: Preparation of (3S)-3-{(S)-2-(4-chlorobenzyloxycarbonylamino)-4-methylvalerylamino}-2-tetrahydrofura-nol (Compound No. 40 in Table-1)

Melting point: 47-49°C
IR (KBr, cm⁻¹): 3308, 1705, 1657, 1541.
NMR (CDCl₃, δ): 0.94 (m, 6H), 1.50-1.86 (m, 4H), 2.32 (m, 0.7H), 2.44 (m, 0.3H), 2.88 (s, 0.3H), 3.03 (s, 0.7H), 3.88 (ddd, J=7.8Hz, 7.8HZ, 7.8Hz, 0.7H), 3.95 (m, 0.3H), 4.12 (m, 2H), 4.35 (m, 1H), 5.06 (s, 0.6H), 5.07 (s, 1.4H), 5.22 (m, 1.3H), 5.30 (s, 0.7H), 6.09 (s, 0.3H), 6.39 (d, J=8.7Hz, 0.7H), 7.31 (m, 4H).

Example 18: Preparation of (3S)-3-{(S)-4-methyl-2-(2-methylbenzyloxycarbonylamino)valerylamino}-2-tetrahydrofura-nol (Compound No. 44 in Table-1)

Melting point: 120-122°C
IR (KBr, cm⁻¹): 3302, 1694, 1645, 1541.
NMR (CDCl₃, δ): 0.93 (m, 6H), 1.52 (m, 1H), 1.59 (m, 3H), 2.16-2.49 (m, 1H), 2.34 (s, 3H), 3.28 (s, 0.5H), 3.52 (s, 0.5H), 3.86 (dd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 3.98 (m, 0.5H), 4.13 (m, 2H), 4.31 (m, 1H), 5.12 (s, 2H), 5.27 (m, 2H), 6.34 (s, 0.5H), 6.51 (s, 0.5H), 7.19 (m, 2H), 7.31 (m, 2H).

Example 19: Preparation of (3S)-3-{(S)-4-methyl-2-(4-methylbenzyloxycarbonylamino)valerylamino}-2-tetrahydrofura-nol (Compound No. 46 in Table-1)

IR (KBr, cm⁻¹): 3310, 1703, 1657, 1539.
NMR (CDCl₃, δ): 0.93 (m, 6H), 1.52 (m, 1H), 1.58-1.86 (m, 3H), 2.29 (m, 0.5H), 2.35 (s, 3H), 2.43 (m, 0.5H), 2.91 (s, 0.5H), 3.03 (s, 0.5H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 3.97 (m, 0.5H), 4.10 (m, 2H), 4.33 (m, 1H), 5.06 (s, 2H), 5.14 (m, 1H), 5.26 (m, 1H), 6.20 (s, 0.5H), 6.44 (s, 0.5H), 7.16 (m, 2H), 7.23 (m, 2H).

Example 20: Preparation of (3S)-3-{(S)-2-(2-methoxybenzyloxycarbonylamino)-4-methylvalerylamino)-2-tetrahydro-furanol (Compound No. 47 in Table-1)

Melting point: 36-38°C

IR (KBr, cm^{-1}): 3308, 1703, 1657, 1539.

NMR (CDCl$_3$, δ): 0.93 (m, 6H), 1.51 (m, 1H), 1.60-1.91 (m, 3H), 2.30 (m, 0.6H), 2.42 (m, 0.4H), 3.15 (s, 0.4H), 3.31 (d, J=3.0Hz, 0.6H), 3.83 (s, 3H), 3.87 (m, 0.6H), 3.91 (m, 0.4H), 4.13 (m, 2H), 4.32 (m, 1H), 5.16 (s, 0.8H), 5.18 (s, 1.2H), 5.23 (m, 2H), 6.65 (s, 0.4H), 6.53 (d, J=7.5Hz, 0.6H), 6.91 (m, 2H), 7.31 (m, 2H).

Example 21: Preparation of (3S)-3-{(S)-2-(4-methoxybenzyloxycarbonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 49 in Table-1)

Melting point: 30-33°C

IR (KBr, cm^{-1}): 3310, 1701, 1657, 1516.

NMR (CDCl$_3$, δ): 0.92 (m, 6H), 1.50 (m, 1H), 1.59-1.88 (m, 3H), 2.29 (m, 0.5H), 2.42 (m, 0.5H), 3.01 (s, 0.5H), 3.20 (d, J=3.0Hz, 0.5H), 3.80 (s, 3H), 3.88 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 3.96 (m, 0.5H), 4.11 (m, 2H), 4.33 (m, 1H), 5.03 (s, 2H), 5.15 (m, 1H), 5.27 (s, 0.5H), 5.32 (s, 0.5H), 6.22 (s, 0.5H), 6.66 (s, 0.5H), 6.87 (m, 2H), 7.28 (m, 2H).

Example 22: Preparation of (3S)-3-{(S)-2-(9-fluorenylmethoxycarbonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 51 in Table-1)

Melting point: 150-153°C

IR (KBr, cm^{-1}): 3314, 1725, 1696, 1653, 1534.

NMR (CDCl$_3$, δ): 0.93 (m, 6H), 1.65 (m, 3H), 1.91 (m, 1H), 2.32 (m, 0.7H), 2.42 (m, 0.3H) 3.08 (s, 0.3H), 3.28 (d, J=3Hz, 0.7H), 3.86 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.7H), 4.01 (m, 0.3H), 4.08-4.24 (m, 3H), 4.32-4.52 (m, 3H), 5.27 (m, 2H), 6.21 (s, 0.3H), 6.38 (s, 0.7H), 7.31 (m, 2H), 7.40 (m, 2H), 7.56 (dd, J=7.4Hz, 2H), 7.76 (d, J=7.4Hz, 2H).

Example 23: Preparation of (3S)-3-((S)-4-methyl-2-tetrahydrofurfuryloxycarbonylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 52 in Table-1)

Melting point: 40-43°C

IR (KBr, cm^{-1}): 3308, 1705, 1659, 1543.

NMR (CDCl$_3$, δ): 0.94 (m, 6H), 1.46-2.07 (m, 8H), 2.32 (m, 0.5H), 2.46 (m, 0.5H), 3.36 (s, 0.5H), 3.39 (s, 0.5H), 3.78-4.25 (m, 8H), 4, 29-4.42 (m, 1H), 5.28 (m, 1.5H), 5.40 (s, 0.5H), 6.34 (s, 0.5H), 5.56 (s, 0.5H).

Example 24: Preparation of (3S)-3-{(S)-4-methyl-2-(2-tetrahydropyranylmethoxycarbonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 53 in Table-1)

IR (KBr, cm^{-1}): 3306, 1705, 1659, 1539.

NMR (CDCl$_3$, δ): 0.94 (m, 6H), 1.30 (m, 1H), 1.53 (m, 3H), 1.68 (m, 4H), 1.84 (m, 2H), 2.32 (m, 0.5H), 2.47 (m, 0.5H), 3.44 (m, 1H), 3.55 (m, 1H), 3.65 (s, 0.5H), 3.74 (s, 0.5H), 3.98 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.00 (m, 3.5H), 4.12 (m, 2H), 4.35 (m, 1H), 5.28 (d, J=3.0Hz, 0.5H), 5.31 (m, 1H), 5.46 (m, 0.5H), 6.45 (s, 0.5H), 6.61 (m, 0.5H).

Example 25: Preparation of (3S)-3-{(S)-4-methyl-2-(2-pyridylmethoxycarbonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 54 in Table-1)

Melting point: 54-56°C

IR (KBr, cm^{-1}): 3306, 1711, 1657, 1541.

NMR (CDCl$_3$, δ): 0.95 (m, 6H), 1.53 (m, 2H), 1.68 (m, 3H), 2.28 (m, 0.5H), 2.47 (m, 0.5H), 3.85 (ddd, J=8.1Hz, 8.1Hz, 8.1Hz, 0.5H), 3.97 (ddd, J=8.1Hz, 8.1Hz, 8.1Hz, 0.5H), 4.08-4.44 (m, 3H), 5.10-5.36 (m, 3H), 5.75 (s, 1H), 6.67 (s, 0.5H), 6.69 (s, 0.5H), 7.19-7.40 (m, 2H), 7.69 (m, 1H), 8.50 (d, J=3.9Hz, 0.5H), 8.56 (d, J=3.9Hz, 0.5H).

Example 26: Preparation of 3-{(S)-4-methyl-2-(2-pyridylmethoxycarbonylamino)valerylamino)-2-tetrahydrofuranol N-oxide (Compound No. 57 in Table-1)

NMR (CDCl$_3$, δ): 0.93 (m, 6H), 1.50-1.92 (m, 4H), 2.10-2.48 (m, 1H), 2.86 (s, 0.8H), 3.30 (s, 0.2H), 3.84 (m, 1H), 4.08 (m, 1H), 4.36 (m, 2H), 5.10-5.68 (m, 3H), 5.99 (s, 0.5H), 6.12 (s, 0.3H), 6.30 (m, 0.2H), 6.82-7.07 (m, 1H), 7.35 (m, 3H), 8.28 (d, J=8.6Hz, 1H).

Example 27: Preparation of (3S)-3-((S)-2-cyclohexyloxycarbonylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 60 in Table-1)

Melting point: 28-30°C

IR (KBr, cm^{-1}): 3310, 1696, 1657, 1539.

NMR (CDCl$_3$, δ): 0.95 (m, 6H), 1.29 (m, 2H), 1.36 (m, 4H), 1.53 (m, 2H), 1.69 (m, 4H), 1.85 (m, 2H), 2.36 (m, 0.5H), 2.48 (m, 0.5H), 2.85 (s, 0.5H), 3.06 (s, 0.5H), 3.89 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.02 (m, 0.5H), 4.14 (m, 2H), 4.36 (m, 1H), 4.63 (m, 1H), 4.99 (m, 1H), 5.26 (m, 0.5H), 5.32 (m, 0.5H), 6.22 (s, 0.5H), 6.47 (s, 0.5H).

Examle28: Preparation of (3S)-3-((S)-4-methyl-2-phenoxycarbonylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 61 in Table-1)

Melting point: 68-70°C

IR (KBr, cm^{-1}): 3308, 1723, 1659, 1539.

NMR (CDCl$_3$, δ): 0.98 (m, 6H) 1.52 (m, 1H), 1.58-1.86 (m, 3H), 2.30-2.52 (m, 1H), 2.98 (s, 0.6H), 3.23 (s, 0.4H), 3.89 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.6H), 4.01 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.4H), 4.09-4.27 (m, 2H), 4.38 (m, 1H), 5.27 (d, J=2.7Hz, 0.4H), 5.33 (dd, J=3.6Hz, 3.6Hz, 0.6H), 5.60 (s, 1H), 6.10 (s, 0.4H), 6.24 (s, 0.6H), 7.12 (m, 2H), 7.20 (m, 1H), 7.35 (m, 2H).

Example 29: Preparation of (3S)-3-((S)-4-methyl-2-phenylureidovalerylamino)-2-tetrahydrofuranol (Compound No. 63 in Table-1)

Melting point: 181-182°C

IR (KBr, cm^{-1}): 3291, 1638, 1555.

NMR (CDCl$_3$, δ): 0.96 (m, 6H), 1.50 (m, 2H), 1.67 (m, 1H), 1.86 (m, 1H), 2.26 (m, 0.5H), 2.41 (m, 0.5H), 2.88 (s, 0.5H), 3.40 (s, 0.5H), 3.82 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.07 (m, 1.5H), 4.29 (m, 2H), 5.20 (s, 0.5H), 5.26 (d, J=4.5Hz, 0.5H), 5.98 (m, 1H), 7.02 (m, 1H), 7.24 (m, 5H), 7.73 (s, 0.5H), 7.91 (s, 0.5H).

Example 30: Preparation of (3S)-3-{(S)-2-(3,3-dimethylbutyrylamino)-4-methylvlerylamino}-2-tetrahydrofuranol (Compound No. 73 in Table-1)

Melting point: 152-153°C

IR (KBr, cm^{-1}): 3293, 1642, 1549.

NMR (CDCl$_3$, δ): 0.93 (m, 6H), 1.02 (s, 9H), 1.49-1.94 (m, 4H), 2.07 (s, 1H), 2.08 (s, 1H), 2.32 (m, 0.5H), 2.45 (m, 0.5H), 3.17 (d, J=3.0Hz, 0.5H), 3.73 (d, J=3.0Hz, 0.5H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.02 (m, 0.5H), 4.10 (m, 1H), 4.23-4.54 (m, 3H), 5.28 (d, J=2.4Hz, 0.5H), 5.31 (dd, J=2.4Hz, 2.4Hz, 0.5H), 6.62 (s, 0.5H), 6.69 (s, 0.5H).

Eample31: Preparation of (3S)-3-((S)-4-methyl-2-tetradecanoylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 78 in Table-1)

Melting point: 96-98°C

IR (KBr, cm^{-1}): 3292, 1636, 1543.

NMR (CDCl$_3$, δ): 0.88 (t, J=5.3Hz, 3H), 0.94 (m, 6H), 1.25 (m, 22H), 1.51-1.94 (m, 4H), 2.20 (m, 2H), 2.31 (m, 0.5H), 2.43 (m, 0.5H), 3.10 (d, J=2.7Hz, 0.5H), 3.65 (d, J=2.7Hz, 0.5H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.00 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.11 (m, 1H), 4.26-4.55 (m, 2H), 5.27 (d, J=2.7Hz, 0.5H), 5.31 (dd, J=4.1Hz, 4.1Hz, 0.5H), 5.97 (s, 1H), 6.52 (s, 0.5H), 6.62 (s, 0.5H).

Example 32: Preparation of (3S)-3-{(S)-4-methyl-2-(3-phenylpropionylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 83 in Table-1)

Melting point: 60-62°C

IR (KBr, cm^{-1}): 3291, 1644, 1549.

NMR (CDCl$_3$, δ): 0.89 (m, 6H), 1.81 (m, 1H), 1.44-1.62 (m, 3H), 2.28-2.46 (m, 1H), 2.50 (m, 2H), 2.77 (s, 0.6H), 2.93 (m, 2H), 3.08 (s, 0.4H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.6H), 4.05 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.4H), 4.12 (m, 1H), 4.24-4.48 (m, 2H), 5.24 (s, 0.4H), 5.29 (m, 0.6H), 5.79 (m, 1H), 6.28 (d, J=7.5Hz, 0.4H), 6.45 (d, J=7.5Hz, 0.6H), 7.23 (m, 5H).

Example 33: Preparation of (3S)-3-{(S)-4-methyl-2-(1-naphthylacetylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 85 in Table-1)

Melting point: 164-167°C
IR (KBr, cm^{-1}): 3279, 1638.
NMR (CDCl$_3$+DMSO-d$_6$, δ): 0.76 (d, J=6.0Hz, 3H), 0.78 (d, J=5.8Hz, 3H), 1.35 (m, 2H), 1.48 (m, 1H), 1.72 (m, 1H), 2.19 (m, 0.7H), 2.31 (m, 0.3H), 3.78 (ddd, J=8.0Hz, 8.0Hz, 8.0Hz, 0.7H), 3.85 (ddd, J=8.0Hz, 8.0Hz, 8.0Hz, 0.3H), 3.97-4.09 (m, 3H), 4.20 (m, 1H), 4.41 (m, 1H), 5.09 (d, J=4.0Hz, 0.3H), 5.15 (d, J=3.7Hz, 0.3H), 5.22 (dd, J=4.4Hz, 4.4Hz, 0.7H), 5.38 (d, J=4.3Hz, 0.7H), 6.42 (d, J=7.8Hz, 0.7H), 6.56 (d, J=9.0Hz, 0.3H), 6.71 (d, J=7.4Hz, 0.7H), 6.97 (d, J=7.2Hz, 0.3H), 7.42-7.53 (m, 4H), 7.55-7.88 (m, 2H), 7.98 (d, J=7.4Hz, 1H).

Example 34: Preparation of (3S)-3-((S)-4-methyl-2-phenoxyacetylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 90 in Table-1)

Melting point: 30°C
IR (KBr, cm^{-1}): 3297, 1653.
NMR (CDCl$_3$ δ): 0.92 (d, J=6.0Hz, 3H), 0.92 (d, J=5.7Hz, 3H), 1.55-1.70 (m, 2H), 1.84 (m, 2H), 2.32 (m, 0.6H), 2.45 (m, 0.4H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.6H), 4.01 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.4H), 4.11 (m, 1H), 4.33 (m, 1H), 4.51 (s, 0.8H), 4.52 (s, 1.2H), 4.55 (m, 1H), 5.28 (s, 0.4H), 5.33 (d, J=4.5Hz, 0.6H), 6.63 (d, J=7.5Hz, 0.4H), 6.68 (d, J=8.1Hz, 0.6H), 6.93 (m, 2H), 7.03 (m, 2H), 7.31 (m, 2H).

Example 35: Preparation of (3S)-3-{(S)-2-(2-chlorophenoxyacetylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 94 in Table-1)

Melting point: 49-52°C
IR (KBr, cm^{-1}): 3302, 3074, 1655, 1537.
NMR (CDCl$_3$, δ): 0.94 (m, 6H), 1.60-1.91 (m, 4H), 2.31 (m, 0.7H), 2.45 (m, 0.3H), 3.27 (d, J=2.7Hz, 0.3H), 3.72 (d, J=2.7Hz, 0.7H), 3.87 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.7H), 4.06 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.3H), 4.11 (m, 1H), 4.40 (m, 1H), 4.52 (m, 1H), 4.52 (s, 0.6H), 4.57 (s, 1.4H), 5.29 (s, 0.3H), 5.33 (dd, J=3.6Hz, 3.6Hz, 0.7H), 6.49 (s, 0.3H), 6.64 (s, 0.7H), 6.90 (m, 1H), 7.02 (m, 1H), 7.26 (m, 2H), 7.41 (m, 1H).

Example 36: Preparation of (3S)-3-{(S)-2-(4-chlorophenoxyacetylamino)-4-methylvalerylamino}-2-tertahydrofuranol (Compound No. 96 in Table-1)

Melting point: 53-55°C
IR (KBr, cm^{-1}): 3301, 1653, 1541.
NMR (CDCl$_3$, δ): 0.94 (m, 6H), 1.69-1.88 (m, 4H), 2.32 (m, 0.5H), 2.58 (m, 0.5H), 2.91 (d, J=2.8Hz, 0.5H), 3.28 (d, J=3.0Hz, 0.5H), 3.88 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.06 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.12 (m, 1H), 4.34 (m, 1H), 4.47 (s, 1H), 4.49 (s, 1H), 4.53 (m, 1H), 5.28 (d, J=2.6Hz, 0.5H), 5.33 (dd, J=4.5Hz, 4.5Hz, 0.5H), 6.29 (s, 0.5H), 6.47 (s, 0.5H), 6.88 (m, 3H), 7.27 (m, 2H).

Example 37: Preparation of (3S)-3-((S)-4-methyl-2-phenylthioacetylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 110 in Table-1)

Melting point: 45-47°C
IR (KBr, cm^{-1}): 3287, 1645, 1551.
NMR (CDCl$_3$, δ): 0.81 (m, 6H), 1.35 (m, 1H), 1.52 (m, 2H), 1.70 (m, 1H), 2.25 (m, 0.5H), 2.39 (m, 0.5H), 3.23 (s, 0.5H), 3.66 (m, 2.5H), 3.85 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 3.96 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.5H), 4.09 (m, 1H), 4.40-4.42 (m, 2H), 5.23 (d, J=2.3Hz, 0.5H), 5.28 (dd, J=3.9Hz, 3.9Hz, 0.5H), 6.39 (s, 0.5H), 6.53 (s, 0.5H), 7.10 (m, 1H), 7.21 (m, 1H), 7.29 (m, 3H), 7.30 (s, 1H).

Example 38: Preparation of (3S)-3-{(S)-4-methyl-2-(3-pheylsulfonylpropionylamino)valerylamino)-2-tetrahydorofuranol (Compound No. 111 in Table-1)

IR (KBr, cm^{-1}): 3301, 1649, 1547.
NMR (CDCl$_3$, δ): 0.93 (m, 6H), 1.48-1.74 (m, 5H), 1.80-1.92 (m, 1H), 2.30 (m, 0.5H), 2.43 (m, 0.5H), 2.72 (m, 2H), 3.41 (m, 1H), 3.52 (m, 1H), 3.86 (ddd, J=8.1Hz, 8.1Hz, 8.1Hz, 0.5H), 4.02 (m, 0.5H), 4.09 (m, 2H), 4.28-4.48 (m, 2H), 5.29 (s, 0.5H), 5.33 (dd, J=4.2Hz, 4.2Hz, 0.5H), 6.27 (s, 1H), 6.46 (s, 0.5H), 6.58 (s, 0.5H), 7.59 (m, 2H), 7.69 (m, 1H), 7.93 (dd, J=7.2Hz, 5.4Hz, 2H).

Example 39: Preparation of (3S)-3-((S)-2-benzoylamino-4-methylvalerylamino)-2-t trahydrofuranol (Compound No. 112 in Table-1)

Melting point: 154- 156°C

IR (KBr, cm^{-1}): 3300, 1665, 1636.

NMR (CDCl$_3$, δ): 0.98 (d, J=5.5Hz, 6H), 1.71 (m, 3H), 1.87 (m, 1H), 2.31 (m, 0.7H), 2.42 (m, 0.3H), 3.16 (s, 0.3H), 3.60 (s, 0.7H), 3.87 (ddd, J=8.2Hz, 8.2Hz, 8.2Hz, 0.7H), 4.03 (ddd, J=7.5Hz, 7.5Hz, 7.5Hz, 0.3H), 4.12 (ddd, J=8.6Hz, 8.6Hz, 3.5Hz, 1H), 4.36 (m, 1H), 4.68 (m, 1H), 5.30 (s, 0.3H), 5.35 (d, J=4.7Hz, 0.7H), 6.67-6.75 (m, 2H), 7.43 (m, 2H), 7.52 (m, 1H), 7.79 (m, 2H).

Example 40: Preparation of (3S)-3-{(S)-2-(2-fluorobenzoylamino)-4-metylvalerylamino}-2-tetrahydrofuranol (Compound No. 113 in Table-1)

Melting point: 62-64°C

IR (KBr, cm^{-1}): 3306, 1644, 1534.

NMR (CDCl$_3$, δ): 0.80-1.07 (m, 6H), 1.59-2.01 (m, 4H), 2.40 (m, 1H), 3.82 (m, 1H), 3.97-4.20 (m, 1.6H), 4.24-4.45 (m, 1.4H), 4.68 (m, 1H), 5.31 (d, J=2.5Hz, 0.6H), 5.35 (dd, J=4.2Hz, 4.1Hz, 0.4H), 6.87 (m, 1H), 7.02-7.20 (m, 2H), 7.25 (m, 1H), 7.50 (m, 1H), 8.00 (m, 1H).

Example 41: Preparation of (3S)-3-{(S)-2-(3-fluorobenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 114 in Table-1)

Melting point: 159-161°C

IR (KBr, cm^{-1}): 3304, 3076, 1638, 1588, 1547.

NMR (CDCl$_3$, δ): 0.87-1.07 (m, 6H), 1.61-1.98 (m, 4H), 2.39 (m, 1H), 3.39 (d, J=2.6Hz, 0.5H), 3.87 (m, 1H), 3.98-4.18 (m, 1.5H), 4.35 (m, 1H), 4.67 (m, 1H), 5.31 (d, J=2.6Hz, 0.5H), 5.35 (dd, J=4.1Hz, 4.1Hz, 0.5H), 6.71 (d, J=7.6Hz, 0.5H), 6.76 (d, J=7.6Hz, 0.5H), 6.90 (d, J=8.2Hz, 0.5H), 6.99 (d, J=8.2Hz, 0.5H), 7.20 (ddd, J=8.2Hz, 8.2Hz, 2.7Hz, 1H), 7.40 (m, 1H), 7.43-7.60 (m, 2H).

Example 42: Preparation of (3S)-3-{(S)-2-(4-fluorobenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 115 in Table-1)

Melting point: 151-153°C

IR (KBr, cm^{-1}): 3422, 3301, 1640, 1545, 1503.

NMR (CDCl$_3$, δ): 0.85-1.17 (m, 6H), 1.60-1.97 (m, 4H), 2.37 (m, 1H), 3.79 (d, J=2.7Hz, 0.55H), 3.87 (m, 0.55H), 4.06 (m, 0.45H), 4.10 (m, 1H), 4.27 (ddd, J=6.8Hz, 6.8Hz, 2.1Hz, 0.45H), 4.39 (m, 1H), 5.31 (d, J=2.7Hz, 0.55H), 5.35 (d, J=4.1Hz, 4.1Hz, 0.45H), 6.28 (d, J=8.3Hz, 0.45H), 6.96 (d, J=8.3Hz, 0.55H), 7.02-7.19 (m, 3H), 7.75-7.89 (m, 2H).

Example 43: Preparation of (3S)-3-{(S)-2-(4-chlorobenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 121 in Table-1)

Melting point: 93-96°C

IR (KBr, cm^{-1}): 3295, 1636.

NMR (CDCl$_3$, δ): 0.95 (d, J=6.0Hz, 3H), 0.97 (d, J=5.7Hz, 3H), 1.73 (m, 3H), 1.86 (m, 1H), 2.31 (m, 0.7H), 2.42 (m, 0.3H), 3.41 (s, 0.3H), 3.86 (s, 0.7H), 3.87 (ddd, J=8.4Hz, 8.4Hz, 8.4Hz, 0.7H), 4.02 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.3H), 4.12 (m, 1H), 4.35 (m, 1H), 4.68 (m, 1H), 5.30 (s, 0.3H), 5.35 (d, J=4.5Hz, 0.7H), 6.71 (d, J=8.1Hz, 0.7H), 6.76 (d, J=6.9Hz, 0.3H), 6.87 (d, J=6.4Hz, 0.7H), 6.95 (d, J=8.1Hz, 0.3H), 7.39 (dd, J=8.4Hz, 1.8Hz, 2H), 7.73 (dd, J=8.4Hz, 2.1Hz, 2H).

Example 44: Preparation of (3S)-3-{(S)-4-methyl-2-(2-methylbenzoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 125 in Table-1)

Melting point: 73-74°C

IR (KBr, cm^{-1}): 3298, 1638, 1541.

NMR (CD$_3$OD, δ): 0.98 (d, J=6.1Hz, 6H), 1.50-1.99 (m, 4H), 2.30 (m, 1H), 2.38 (s, 1.65H), 2.39 (s, 1.35H), 3.85 (m, 0.55H), 3.98-4.17 (m, 1.45H), 4.23 (m, 1H), 4.60 (m, 1H), 5.16 (s, 0.55H), 5.25 (d, J=4.7Hz, 0.45H), 7.18-7.29 (m, 2H), 7.30-7.42 (m, 2H).

Example 45: Preparation of (3S)-3-{(S)-4-methyl-2-(3-methylbenzoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 126 in Table-1)

Melting point: 85-87°C
IR (KBr, cm⁻¹): 3299, 1638, 1586, 1541.
NMR (CDCl₃, δ): 0.83-1.07 (m, 6H), 1.60-1.97 (m, 3H), 2.07 (m, 1H), 2.30 (m, 1H), 2.35 (s, 1.8H), 2.36 (s, 1.2H), 3.82 (m, 0.6H), 3.96-4.18 (m, 1.6H), 4.18-4.42 (m, 1.4H), 4.75 (m, 1H), 4.93 (m, 0.4H), 5.31 (d, J=2.9Hz, 0.6H), 5.35 (dd, J=4.4Hz, 4.3Hz, 0.4H), 7.01 (m, 1H), 7.14 (m, 1H), 7.32-7.40 (m, 2H), 7.57-7.67 (m, 2H).

Example 46: Preparation of (3S)-3-{(S)-4-methyl-2-(4-methylbenzoylamino)valerylamino}-2-tetrahydorofuranol (Compound No. 127 in Table-1)

Melting point: 101-102°C
IR (KBr, cm⁻¹): 3304, 1634, 1545, 1504.
NMR (CDCl₃, δ): 0.87-1.07 (m, 6H), 1.60-1.95 (m, 4H), 2.37 (m, 1H), 2.39 (s, 3H), 3.45 (d, J=2.9Hz, 0.6H), 3.90 (m, 0.4H), 3.97-4.19 (m, 2H), 4.38 (m, 1H), 4.71 (m, 1H), 5.30 (d, J=2.9Hz, 0.6H), 5.35 (dd, J=6.7Hz, 6.7Hz, 0.6H), 6.74 (d, J=8.5Hz, 0.4H), 6.80 (d, J=8.5Hz, 1H), 6.89 (d, J=6.7Hz, 0.6H), 7.22 (d, J=8.2Hz, 2H), 7.68 (d, J=8.2Hz, 2H).

Example 47: Preparation of (3S)-3-{(S)-2-(2,6-dimethylbenzoylamino)-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 129 in Table-1)

Melting point: 89-91°C
IR (KBr, cm⁻¹): 3389, 1638, 1539.
NMR (CDCl₃, δ): 0.83-1.10 (m, 6H), 1.58-1.99 (m, 4H), 2.25 (s, 3H), 2.27 (s, 3H), 2.32 (m, 1H), 3.65-3.93 (m, 1.55H) 4.07 (m, 1H), 4.22-4.45 (m, 1.45H), 4.70 (m, 1H), 5.25 (d, J=3.1Hz, 0.55H), 5.31 (dd, J=4.3Hz, 4.2Hz, 0.45H), 6.36 (d, J=8.2Hz, 1H), 6.85-7.07 (m, 3H), 7.14 (dd, J=7.8Hz, 7.3Hz, 1H).

Example 48: Preparation of (3S)-3-{(S)-2-(3,4-dimethylbenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 130 in Table-1)

Melting point: 92-94°C
IR (KBr, cm⁻¹): 3299, 1636, 1541.
NMR (CDCl₃, δ): 0.83-1.10 (m, 6H), 1.57-1.99 (m, 4H), 2.28 (s, 3H), 2.29 (s, 3H), 2.34 (m, 1H), 3.57 (s, 0.5H), 3.86 (m, 0.5H), 3.97-4.20 (m, 1.5H), 4.21-4.45 (m, 1.5H), 4.70 (m, 1H), 5.31 (s, 0.5H), 5.35 (d, J=2.1Hz, 0.5H), 6.74 (d, J=8.3Hz, 0.5H), 6.80 (d, J=8.2Hz, 0.5H), 6.85 (d, J=8.5Hz, 0.5H), 6.93 (d, J=7.1Hz, 0.5H), 7.17 (d, J=7.8Hz, 1H), 7.43-7.60 (m, 2H).

Example 49: Preparation of (3S)-3-{(S)-4-methyl-2-(2,4,6-trimethylbenzoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 131 in Table-1)

Melting point: 150-152°C
IR (KBr, cm⁻¹): 3295, 1638, 1522.
NMR (CDCl₃, δ): 0.87-1.07 (m, 6H), 1.58-1.97 (m, 4H), 2.18-2.35 (m, 9H), 2.36 (m, 1H), 3.09 (s, 0.45H), 3.48 (s, 0.55H), 3.85 (m, 0.55H), 3.97-4.20 (m, 1.45H), 4.35 (m, 1H), 4.65 (m, 1H), 5.29 (d, J=2.5Hz, 0.45H), 5.34 (dd, J=4.1Hz, 3.8Hz, 0.55H), 6.08 (d, J=5.2Hz, 1H), 6.75 (m, 1H), 6.83 (s, 2H).

Example 50: Preparation of (3S)-3-{(S)-2-(4-ethylbenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 134 in Table-1)

Melting point: 98-99°C
IR (KBr, cm⁻¹): 3304, 1672, 1634, 1545, 1505.
NMR (CDCl₃, δ): 0.83-1.05 (m, 6H), 1.23 (t, J=7.6Hz, 3H), 1.60-1.99 (m, 4H), 2.38 (m, 1H), 2.68 (q, J=7.6Hz, 2H), 3.34 (s, 0.4H), 3.87 (m, 1H), 4.00-4.20 (m, 1.6H), 4.35 (m, 1H), 4.68 (m, 1H), 5.30 (s, 0.6H), 5.34 (d, J=3.7Hz, 0.4H), 6.65-6.90 (m, 2H), 7.21-7.27 (m, 2H), 7.69-7.74 (m, 2H).

Example 51: Preparation of (3S)-3-{(S)-4-methyl-2-(4-trifluoromethylbenzoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 137 in Table-1)

Melting point: 135-136°C

IR (KBr, cm^{-1}): 3310, 1640, 1548, 1508.

NMR (CDCl$_3$, δ): 0.87-1.07 (m, 6H), 1.61-1.97 (m, 4H), 2.37 (m, 1H), 3.43 (s, 0.4H), 3.82-4.08 (m, 1.6H), 4.17 (m, 1H), 4.29 (m, 1H), 4.70 (m, 1H), 5.31 (d, J=2.6Hz, 0.4H), 5.36 (d, J=4.2Hz, 4.2Hz, 0.6H), 6.71 (d, J=7.9Hz, 0.6H), 6.73 (d, J=7.9Hz, 0.4H), 7.03 (d, J=8.3Hz, 0.6H), 7.14 (d, J=8.3Hz, 0.4H), 7.61-7.86 (m, 2H), 7.87-7.93 (m, 2H).

Example 52: Preparation of (3S)-3-{(S)-2-(2-methoxybenzoylamino)-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 138 in Table-1)

Melting point: 65-66°C

IR (KBr, cm^{-1}): 3376, 1640, 1601, 1532.

NMR (CDCl$_3$, δ): 0.83-1.05 (m, 6H), 1.60-1.95 (m, 4H), 2.31 (m, 1H), 3.85 (m, 0.6H), 3.98 (s, 3H), 4.00-4.20 (m, 1.6H), 4.24-4.45 (m, 1.4H), 4.69 (m, 1H), 5.00 (m, 0.4H), 5.34 (m, 1H), 6.93-7.17 (m, 2.4H), 7.22 (m, 0.6H), 7.50 (m, 1H), 8.15 (m, 1H), 8.30 (m, 1H).

Example 53: Preparation of (3S)-3-{(S)-2-(4-methoxybenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 140 in Table-1)

Melting point: 85-88°C
IR (KBr, cm^{-1}): 3295, 1632.

NMR (CDCl$_3$, δ): 0.95 (d, J=6.0Hz, 3H), 0.96 (d, J=4.8Hz, 3H), 1.72 (m, 3H), 1.86 (m, 1H), 2.29 (m, 0.5H), 2.40 (m, 0.5H), 3.84 (s, 3H), 3.86 (ddd, J=8.1Hz, 8.1Hz, 8.1Hz, 0.5H), 4.01 (ddd, J=8.1Hz, 8.1Hz, 8.1Hz, 0.5H), 4.08 (m, 1H), 4.28 (m, 0.5H), 4.35 (m, 0.5H), 4.68 (m, 1H), 5.30 (s, 0.5H), 5.34 (d, J=4.8Hz, 0.5H), 6.71 (d, J=8.4Hz, 0.5H), 6.78 (d, J=8.4Hz, 0.5H), 6.82 (d, J=8.1Hz, 0.5H), 6.91 (dd, J=9.0Hz, 2.4Hz, 2H), 6.94 (d, J=8.7Hz, 0.5H), 7.76 (m, 2H).

Example 54: Preparation of (3S)-3-{(S)-2-(2,4-dimethoxybenzoylamino)-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 141 in Table-1)

Melting point: 65-67°C
IR (KBr, cm^{-1}): 3382, 1638, 1604, 1534.

NMR (CDCl$_3$, δ): 0.83-1.05 (m, 6H), 1.59-1.98 (m, 4H), 2.40 (m, 1H), 3.83 (m, 0.6H), 3.84 (s, 1.2H), 3.86 (s, 1.8H), 3.95 (s, 1.8H), 3.97 (s, 1.2H), 4.00-4.21 (m, 2H), 4.35 (m, 1H), 4.65 (m, 1H), 4.80 (m, 0.4H), 5.32 (d, J=3.2Hz, 0.6H), 5.35 (dd, J=4.6Hz, 4.6Hz, 0.5H), 6.48 (s, 0.4H), 6.49 (s, 0.6H), 6.59 (m, 1H), 6.99 (d, J=8.1Hz, 0.6H), 7.15 (d, J=6.7Hz, 0.4H), 8.07-8.22 (m, 2H).

Example 55: Preparation of (3S)-3-{(S)-2-(2,6-dimethoxybenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 142 in Table-1)

Melting point: 166-168°C
IR (KBr, cm^{-1}): 3299, 3279, 1645, 1597, 1508.

NMR (DMSO-d$_6$, δ): 0.87 (d, J=6.2Hz, 6H), 1.38-1.60 (m, 2H), 1.60-1.83 (m, 2H), 2.15 (m, 1H), 3.65 (m, 1H), 3.71 (s, 2.4H), 3.72 (s, 3.6H), 3.93 (m, 1H), 4.10 (m, 1H), 4.39 (m, 1H), 5.14 (m, 1H), 6.55 (m, 1H), 6.58-6.70 (m, 2H), 7.18 (d, J=5.7Hz, 1H), 7.29 (m, 1H), 8.33 (d, J=8.4Hz, 1H).

Example 56: Preparation of (3S)-3-{(S)-2-(3,5-dimethoxybenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 144 in Table-1)

Melting point: 88-90°C
IR (KBr, cm^{-1}): 3407, 1639, 1595, 1539.

NMR (CDCl$_3$, δ): 0.83-1.05 (m, 6H), 1.60-1.98 (m, 4H), 2.40 (m, 1H), 3.41 (s, 0.6H), 3.70-3.93 (m, 0.8H), 3.79 (s, 6H), 3.95-4.08 (m, 0.6H), 4.10 (m, 1H), 4.25 (m, 1H), 4.65 (m, 1H), 5.30 (d, J=2.2Hz, 0.6H), 5.34 (m, 0.4H), 6.58 (dd, J=1.9Hz, 1.9Hz, 1H), 6.67-6.87 (m, 2H), 6.87-6.97 (m, 2H).

Example 57: Preparation of (3S)-3-{(S)-2-(4-ethoxybenzoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 148 in Table-1)

Melting point: 84-85°C
IR (KBr, cm^{-1}): 3299, 1634, 1609, 1547, 1504.

NMR (CDCl$_3$, δ): 0.83-1.07 (m, 6H), 1.43 (t, J=7.0Hz, 3H), 1.60-1.99 (m, 4H), 2.38 (m, 1H), 3.82 (m, 1H), 3.92-4.20

(m, 3.5H), 4.55 (s, 0.5H), 4.69 (m, 1H), 5.31 (d, J=2.8Hz, 0.5H), 5.35 (dd, J=4.3Hz, 4.1Hz, 0.5H), 6.80 (d, J=8.3Hz, 0.5H), 6.84-7.00 (m, 3H), 7.14 (d, J=7.0Hz, 0.5H), 7.76 (d, J=8.7Hz, 2H).

Example 58: Preparation of (3S)-3-{(S)-4-methyl-2-(3,4-methylenedioxybenzoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 152 in Table-1)

Melting point: 94-96°C
IR (KBr, cm⁻¹): 3410, 3111, 1753, 1659.
NMR (CDCl₃, δ): 0.95 (d, J=6.0Hz, 3H), 0.96 (d, J=5.1Hz, 3H), 1.69-1.81 (m, 3H), 1.86 (m, 1H), 2.30 (m, 0.5H), 2.41 (m, 0.5H), 3.89 (ddd, J=8.1Hz, 8.1Hz, 8.1Hz, 0.5H), 4.05 (ddd, J=7.6Hz, 7.6Hz, 7.6Hz, 0.5H), 4.11 (m, 1H), 4.30 (m, 0.5H), 4.36 (m, 0.5H), 4.64 (m, 1H), 5.30 (s, 0.5H), 5.32 (d, J=4.6Hz, 0.5Hz), 6.02 (s, 2H), 6.61-6.80 (m, 2H), 6.82 (d, J=7.8Hz, 1H), 7.28 (dd, J=2.7Hz, 2.7Hz, 1H), 7.33 (ddd, J=8.0Hz, 2.0Hz, 2.0Hz, 1H).

Example 59: Preparation of (3S)-3-{(S)-4-methyl-2-(1-naphthoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 156 in Table-1)

Melting point: 85-86°C
IR (KBr, cm⁻¹): 3293, 1638, 1535.
NMR (CDCl₃, δ): 0.83-1.05 (m, 6H), 1.59-1.99 (m, 4H), 2.38 (m, 1H), 3.47 (s, 0.45H), 3.88 (m, 1H), 3.94-4.20 (m, 1.55H), 4.38 (m, 1H), 4.80 (m, 1H), 5.29 (d, J=2.9Hz, 0.55H), 5.35 (dd, J=4.4Hz, 3.9Hz, 0.45H), 6.64 (d, J=8.2Hz, 0.55H), 6.73 (d, J=8.1Hz, 0.45H), 6.89 (m, 1H), 7.44 (m, 1H), 7.47-7.68 (m, 3H), 7.80-7.97 (m, 2H), 8.25 (m, 1H).

Example 60: Preparation of (3S)-3-{(S)-4-methyl-2-(2-naphthoylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 157 in Table-1)

Melting point: 98-100°C
IR (KBr, cm⁻¹): 3293, 1640, 1539, 1512.
NMR (CDCl₃, δ): 0.83-1.13 (m, 6H), 1.60-1.99 (m, 4H), 2.36 (m, 1H), 3.53 (s, 0.3H), 3.85 (m, 0.7H), 3.99-4.21 (m, 2H), 4.40 (m, 1H), 4.80 (m, 1H), 5.34 (dd, J=4.4Hz, 4.3Hz, 0.3H), 5.38 (d, J=2.7Hz, 0.7H), 6.88 (d, J=8.5Hz, 0.7H), 6.96 (d, J=7.2Hz, 0.3H), 7.04 (d, J=8.3Hz, 0.7H), 7.12 (d, J=8.2Hz, 0.3H), 7.43-7.64 (m, 2H), 7.79-7.97 (m,4H), 8.32 (s, 1H).

Example 61: Preparation of (3S)-3-{(S)-2-(2-furoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 161 in Table-1)

Melting point: 92-95°C
IR (KBr, cm⁻¹): 3420, 3300, 1645, 1595, 1529.
NMR (CDCl₃, δ): 0.85-1.05 (m, 6H), 1.58-2.00 (m, 4H), 2.25-2.55 (m, 1H), 3.52 (d, J=2.7Hz, 0.55H), 3.87 (dt, J=7.8Hz, 7.8Hz, 0.55H), 3.95-4.17 (m, 1.9H), 4.27-4.45 (m, 1H), 4.45-4.73 (m, 1H), 5.31 (d, J=2.7Hz, 0.55H), 5.35 (dd, J=4.0Hz, 4.0Hz, 0.45H), 6.51 (dd, J=3.1Hz, 1.4Hz, 1H), 6.77 (d, J=7.7Hz, 1H), 6.80-6.93 (m, 1H), 7.14 (dd, J=3.1Hz, 1.9Hz, 1H), 7.46 (dd, J=1.9Hz, 1.4Hz, 1H).

Example 62: Preparation of (3S)-3-{(S)-2-(3-ethyl-1-methyl-5-pyrazole)carbonylamino-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 170 in Table-1)

Melting point: 89-91°C
IR (KBr, cm⁻¹): 3301, 1643.
NMR (CDCl₃, δ): 0.97 (d, J=5.0Hz, 6H), 1.23 (t, J=7.7Hz, 2.1H), 1.23 (t, J=7.6Hz, 0.9H), 1.60-1.90 (m, 4H), 2.34 (m, 0.7H), 2.56 (m, 0.3H), 2.62 (q, J=7.6Hz, 2H), 3.89 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 1H), 4.01-4.17 (m, 2H), 4.08 (s, 0.9H), 4.08 (s, 2.1H), 4.35 (m, 1H), 4.58 (m, 1H), 5.29 (s, 0.3H), 5.35 (d, J=3.9Hz, 0.7H), 6.39 (s, 1H), 6.44 (d, J=6.8Hz, 0.3H), 6.55 (d, J=8.4Hz, 0.7H), 6.60 (d, J=8.4Hz, 1H).

Example 63: Preparation of (3S)-3-{(S)-2-(2-chromancarbonylamino-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 175 in Table-1)

Melting point: 84-85°C
IR (KBr, cm⁻¹): 3299, 1786, 1532.
NMR (CDCl₃, δ): 0.78-0.93 (m, 3H), 0.95-1.05 (m, 3H), 1.35-2.07 (m, 5H), 2.20-2.54 (m, 2H), 2.70-3.00 (m, 2H), 3.80-4.20 (m, 2H), 4.24-4.60 (m, 3H), 5.15 (m, 1H), 5.31 (m, 1H), 6.40 (m, 1H), 6.65 (m, 1H), 6.83-6.97 (m, 2H),

6.98-7.20 (m, 2H).

Example 64: Preparation of (3S)-3-((S)-2-cinnamoylamino-4-methylvalerylamino)-2-tetrahydrofuranol (Compound No. 182 in Table-1)

Melting point: 102-104°C
IR (KBr, cm⁻¹): 3293, 1651, 1624, 1543.
NMR (CDCl₃, δ): 0.96 (m, 6H), 1.63-1.92 (m, 4H), 2.32 (m, 0.5H), 2.47 (m, 0.5H), 3.35 (d, J=2.7Hz, 0.5H), 3.43 (d, J=2.7Hz, 0.5H), 3.92 (m, 1H), 4.02-4.18 (m, 2H), 4.34 (m, 1H), 4.62 (m, 1H), 5.31 (s, 0.5H), 5.35 (m, 0.5H), 6.43 (d, J=15.0Hz, 1H), 6.44 (m, 1H), 6.90 (m, 1H), 7.32 (m, 2H), 7.51 (m, 2H), 7.62 (m, 1H).

Example 65: Preparation of (3S)-3-{(S)-2-(4-methoxycinnamoylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 187 in Table-1)

Melting point: 101-103°C
IR (KBr, cm⁻¹): 3283, 1651, 1602, 1543, 1512.
NMR (CDCl₃, δ): 0.85-1.05 (m, 6H), 1.60-1.99 (m, 4H), 2.30 (m, 1H), 3.77 (s, 1.5H), 3.79 (s, 1.5H), 3.82 (m, 0.5H), 4.10 (m, 1H), 4.30 (m, 0.5H), 4.41 (m, 1H), 4.72 (m, 1H), 5.35 (m, 1H), 6.38 (d, J=15.6Hz, 0.5H), 6.41 (d, J=15.6Hz, 0.5H), 6.65-6.85 (m, 2H), 6.90 (d, J=8.5Hz, 0.5H), 7.15 (d, J=8.5Hz, 0.5H), 7.20 (d, J=8.5Hz, 0.5H), 7.43 (dd, J=8.8Hz, 3.1Hz, 2H), 7.49 (d, J=8.5Hz, 0.5H), 7.58 (d, J=15.6Hz, 0.5H), 7.59 (d, J=15.6Hz, 0.5H).

Example 66: Preparation of (3S)-3-{(S)-2-(4-fluorophenylsulfonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 202 in Table-1)

Melting point: 156-158°C
IR (KBr, cm⁻¹): 3347, 3256, 1649, 1593, 1541.
NMR (DMSO-d₆, δ): 0.71 (d, J=6.5Hz, 3H), 0.80 (d, J=6.6Hz, 3H), 1.28 (m, 2H), 1.45-1.68 (m, 2H), 1.86 (m, 1H), 3.60-3.95 (m, 4H), 4.98 (dd, J=4.4Hz, 4.1Hz, 1H), 6.37 (d, J=4.1Hz, 1H), 7.39 (dd, J=8.8Hz, 8.3Hz, 2H), 7.75 (d, J=7.5Hz, 1H), 7.81 (dd, J=8.8Hz, 5.3Hz, 2H), 8.00 (d, J=9.0Hz, 1H).

Example 67: Preparation of (3S)-3-{(S)-2-(2-chlorophenylsulfonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 203 in Table-1)

IR (KBr, cm⁻¹): 3365, 1657, 1541.
NMR (CDCl₃, δ): 0.65 (d, J=6.2Hz, 1.2H), 0.67 (d, J=6.4Hz, 1.8H), 0.82 (d, J=6.1Hz, 1.2H), 0.84 (d, J=6.4Hz, 1.8H), 1.43-1.72 (m, 4H), 2.13 (m, 0.6H), 2.36 (m, 0.4H), 3.62-4.28 (m, 5H), 5.24 (d, J=3.0Hz, 0.4H), 5.31 (dd, J=3.8Hz, 3.8Hz, 0.6H), 5.91 (d, J=8.8Hz, 0.4H), 5.94 (d, J=7.9Hz, 0.6H), 6.47 (d, J=7.5Hz, 0.4H), 6.65 (d, J=8.1Hz, 0.6H), 7.38-7.45 (m, 1H), 7.45-7.60 (m, 2H), 8.06 (dd, J=7.3Hz, 1.1Hz, 1H).

Example 68: Preparation of (3S)-3-{(S)-2-(4-chlorophenylsulfonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 205 in Table-1)

Melting point: 112-115°C
IR (KBr, cm⁻¹): 3335, 3264, 1649.
NMR (CDCl₃, δ): 0.76 (d, J=6.3Hz, 0.6H), 0.80 (d, J=6.3Hz, 2.4H), 0.87 (d, J=6.9Hz, 0.6H), 0.89 (d, J=6.6Hz, 2.4H), 1.48 (m, 3H), 1.68 (m, 1H), 2.11 (m, 0.8H), 2.40 (m, 0.2H), 3.67 (ddd, J=6.9Hz, 6.9Hz, 6.9Hz, 1H), 3.85 (m, 0.8H), 3.94 (m, 0.2H), 4.05-4.21 (m, 2H), 5.18 (s, 0.2H), 5.25 (d, J=4.5Hz, 0.8H), 5.31 (d, J=9.9Hz, 0.2H), 5.35 (d, J=8.4Hz, 0.8H), 5.94 (d, J=7.8Hz, 0.2H), 6.23 (d, J=7.8Hz, 0.8H), 7.48 (d, J=8.4Hz, 2H), 7.80 (d, J=8.4Hz, 2H).

Example 69: Preparation of (3S)-3-{(S)-2-(4-bromophenylsulfonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 208 in Table-1)

Melting point: 139-140°C
IR (KBr, cm⁻¹): 3478, 3362, 3264, 1647, 1576, 1537.
NMR (DMSO-d₆, δ): 0.63-0.90 (m, 6H), 1.15-1.49 (m, 2H), 1.42-1.66 (m, 2H), 1.83 (m, 0.65H), 2.02 (m, 0.35H), 3.45-3.92 (m, 4H), 4.80 (d, J=4.1Hz, 0.35H), 4.96 (dd, J=4.4Hz, 4.4Hz, 0.65H), 6.10 (d, J=4.1Hz, 0.35H), 6.36 (d, J=4.4Hz, 0.65H), 7.60-7.85 (m, 4.65H), 7.95-8.15 (m, 1.35H).

Example 70: Preparation of (3S)-3-{(S)-4-methyl-2-(4-methylphenylsulfonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 211 in Table-1)

Melting point: 137-138°C
IR (KBr, cm⁻¹): 3343, 3264, 1649, 1541.
NMR (DMSO-d₆, δ): 0.66 (d, J=6.5Hz, 0.9H), 0.74 (d, J=6.7Hz, 3H), 0.80 (d, J=6.7Hz, 2.1H), 1.10-1.40 (m, 3H), 1.53 (m, 1H), 1.85 (m, 0.3H), 1.98 (m, 0.7H), 2.35 (s, 3H), 3.51 (m, 0.7H), 3.58-3.85 (m, 3.3H), 4.78 (d, J=4.4Hz, 0.7H), 4.96 (m, 0.3H), 6.07 (d, J=4.4Hz, 0.7H), 6.39 (d, J=3.9Hz, 0.3H), 7.31 (d, J=7.9Hz, 2H), 7.61 (d, J=7.9Hz, 2H), 7.66 (d, J=6.2Hz, 0.3H), 7.80 (m, 1H), 7.92 (d, J=6.4Hz, 0.7H).

Example 71: Preparation of (3S)-3-{(S)-4-methyl-2-(2,4,6-trimethylphenylsulfonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 215 in Table-1)

Melting point: 75-77°C
IR (KBr, cm⁻¹): 3328, 1657, 1605, 1541.
NMR (CDCl₃, δ): 0.67 (d, J=6.3Hz, 1.35H), 0.68 (d, J=6.3Hz, 1.65H), 0.83 (d, J=6.4Hz, 1.35H), 0.84 (d, J=6.4Hz, 1.65H), 1.38-1.72 (m, 4H), 2.11 (m, 0.65H), 2.29 (s, 3H), 2.31 (m, 0.45H), 2.63 (s, 6H), 3.61 (m, 1H), 3.72 (d, J=3.0Hz, 0.45H), 3.74-3.98 (m, 1H), 3.98-4.24 (m, 2.55H), 5.23 (d, J=3.0Hz, 0.45H), 5.28 (d, J=3.9Hz, 3.9H, 0.55H), 5.46 (d, J=8.6Hz, 0.45H), 5.60 (d, J=8.0Hz, 0.55H), 6.38 (d, J=7.4Hz, 0.45H), 6.54 (d, J=8.0Hz, 0.55H), 6.95 (s, 2H).

Example 72: Preparation of (3S)-3-{(S)-2-(4-tert-butylphenylsulfonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 218 in Table-1)

Melting point: 140-141°C
IR (KBr, cm⁻¹): 3362, 3161, 1647, 1535.
NMR (DMSO-d₆, δ): 0.74 (d, J=6.5Hz, 3H), 0.81 (d, J=6.6Hz, 3H), 1.15-1.41 (m, 3H), 1.29 (s, 9H), 1.54 (m, 1H), 1.95 (m, 1H), 3.46 (m, 1H), 4.62-4.82 (m, 3H), 4.80 (d, J=4.3Hz, 1H), 6.06 (d, J=4.3Hz, 1H), 7.53 (d, J=8.5Hz, 2H), 7.66 (d, J=8.5Hz, 2H), 7.82 (d, J=9.4Hz, 1H), 7.95 (d, J=6.7Hz, 1H).

Example 73: Preparation of (3S)-3-{(S)-2-(4-methoxyphenylsulfonylamino)-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 221 in Table-1)

Melting point: 153-155°C
IR (KBr, cm⁻¹): 3362, 3150, 1647, 1597, 1535, 1501.
NMR (DMSO-d₆, δ): 0.69 (m, 0.6H), 0.76 (d, J=6.5Hz, 2.7H), 0.82 (d, J=6.7Hz, 2.7H), 1.15-1.45 (m, 3H), 1.56 (m, 1H), 2.01 (m, 1H), 3.54 (m, 1H), 3.60-3.90 (m, 3H), 3.81 (s, 3H), 4.89 (d, J=4.6Hz, 0.9H), 4.99 (m, 0.1H), 6.09 (d, J=4.6Hz, 0.9H), 6.41 (d, J=2.7Hz, 0.1H), 7.04 (d, J=8.8Hz, 2H), 7.60-7.80 (m, 3.1H), 7.94 (d, J=6.9Hz, 0.9H).

Example 74: Preparation of (3S)-3-{(S)-4-methyl-2-(3-nitrophenylsulfonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 227 in Table-1)

Melting point: 164-165°C
IR (KBr, cm⁻¹): 3358, 3264, 1649, 1537.
NMR (DMSO-d₆, δ): 0.73-0.95 (m, 6H), 1.20-1.80 (m, 4.75H), 1.95 (m, 0.25H), 3.47-3.65 (m, 2H), 3.77 (m, 1H), 3.94 (m, 1H), 4.73 (d, J=2.7Hz, 0.25H), 4.92 (dd, J=4.1Hz, 4.1Hz, 0.75H), 6.06 (d, J=2.7Hz, 0.25H), 6.29 (d, J=4.1Hz, 0.75H), 7.80-7.95 (m, 2H), 8.05-8.21 (m, 1.25H), 8.35 (d, J=8.4Hz, 0.75H), 8.46 (d, J=7.9Hz, 1H), 8.50 (s, 1H).

Example 75: Preparation of (3S)-3-{(S)-4-methyl-2-(1-naphtylsulfonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 229 in Table-1)

Melting point: 89-91°C
IR (KBr, cm⁻¹): 3580, 3520, 3470, 3281, 1647, 1553.
NMR (DMSO-d₆, δ): 0.34 (d, J=6.0Hz, 1.95H), 0.51 (d, J=6.2Hz, 1.05H), 0.62 (d, J=6.1Hz, 1.95H), 0.71 (d, J=6.4Hz, 1.05H), 1.10-1.60 (m, 4H), 1.75-1.95 (m, 1H), 3.47-3.83 (m, 4H), 4.73 (d, J=4.5Hz, 0.35H), 4.94 (dd, J=4.3Hz, 4.3Hz, 0.65H), 6.03 (d, J=4.5Hz, 0.35H), 6.34 (d, J=4.3Hz, 0.65H), 7.50-7.75 (m, 4H), 7.86 (d, J=8.2Hz, 0.35H), 7.98-8.32 (m, 3.65H), 8.66 (d, J=7.4Hz, 1H).

319

Example 76: Preparation of (3S)-3-{(S)-4-methyl-2-(2-naphtylsulfonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 230 in Table-1)

Melting point: 102- 104°C

IR (KBr, cm^{-1}): 3351, 1655, 1541.

NMR (DMSO-d$_6$, δ): 0.65 (d, J=6.4Hz, 1.5H), 0.65-0.90 (m, 4.5H), 1.14-1.68 (m, 5H), 3.25 (m, 0.5H), 3.57-3.91 (m, 3.5H), 4.73 (s, 0.5H), 4.89 (dd, J=4.3Hz, 4.3Hz, 0.5H), 5.98 (s, 0.5H), 6.32 (d, J=4.3Hz, 0.5H), 7.60-7.80 (m, 3.5H), 7.93 (d, J=6.7Hz, 0.5H), 7.98-8.17 (m, 4H), 8.38 (d, J=8.3Hz, 1H).

Example 77: Preparation of (3S)-3-{(S)-4-methyl-2-(3-pyridylsulfonylamino)valerylamino}-2-tetrahydrofuranol (Compound No. 232 in Table-1)

Melting point: 127-130°C

IR (KBr, cm^{-1}): 3268, 1647.

NMR (CDCl$_3$, δ): 0.79 (d, J=6.3Hz, 0.6H), 0.84 (d, J=6.6Hz, 2.4H), 0.86 (d, J=6.0Hz, 0.6H), 0.90 (d, J=6.6Hz, 2.4H), 1.49-1.61 (m, 3H), 1.71 (m, 1H), 2.09 (m, 0.8H), 2.37 (m, 0.2H), 3.78-3.87 (m, 1.8H), 3.96 (q, J=6.6Hz, 0.2H), 4.05-4.15 (m, 2H), 5.17 (s, 0.2H), 5.22 (d, J=4.8Hz, 0.8Hz), 5.75 (d, J=8.7Hz, 0.8H), 5.79 (d, J=9.3Hz, 0.2H), 6.13 (d, J=6.6Hz, 0.2H), 6.36 (d, J=8.1Hz, 0.8H), 7.45 (dd, J=7.8Hz, 4.8Hz, 1H), 8.17 (ddd, J=7.8Hz, 1.8Hz, 1.8Hz, 1H), 8.80 (dd, J=4.8Hz, 1.2Hz, 1H), 9.07 (d, J=2.1Hz, 1H).

Example 78: Preparation of (3S)-3-((S)-2-benzyloxycarbonylamino-3-phenylpropionylamino)-2-tetrahydrofuranol (Compound No. 246 in Table-1)

Melting point: 144-146°C

IR (KBr, cm^{-1}): 3302, 1696, 1649, 1537.

NMR (CDCl$_3$, δ): 1.69 (m, 1H), 2.20-2.41 (m, 1H), 2.60 (s, 0.8H), 2.82 (s, 0.2H), 2.97 (dd, J=14.7Hz, 7.8Hz, 1H), 3.13 (m, 1H), 3.81 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 0.8H), 4.02 (m, 1.2H), 4.26 (m, 1H), 4.37 (m, 1H), 5.09 (m, 3H), 5.40 (s, 1H), 5.70 (s, 0.2H), 6.08 (s, 0.8H), 7.33 (m, 10H).

Example 79: Preparation of (3S)-3-((S)-2-benzyloxycarbonylamino-3-tert-butoxypropionylamino)-2-tetrahydrofuranol (Compound No. 297 in Table-1)

IR (KBr, cm^{-1}): 3322, 1719, 1661, 1534.

NMR (CDCl$_3$, δ): 1.17 (s, 9H), 1.69-1.98 (m, 1H), 2.30 (m, 0.6H), 2.44 (m, 0.4H), 3.40 (m, 1H), 3.62-3.98 (m, 3H), 4.11 (m, 1H), 4.22 (m, 1H), 4.35 (m, 1H), 5.11 (s, 2H), 5.22 (s, 0.4H), 5.29 (s, 0.6H), 5.76 (s, 1H), 6.80 (s, 0.4H), 7.08 (bs, 0.6H), 7.35 (m, 5H).

Example 80: Preparation of (3S)-4-methyl-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-tetrahydrofuranol (Compound No. 320 in Table-1)

Melting point: 116-121°C

IR (KBr, cm^{-1}): 3356, 3272, 1655.

NMR (CDCl$_3$, δ): 0.70 (d, J=6.0Hz, 2.1H), 0.71 (d, J=6.0Hz, 0.9H), 0.86 (d, J=6.6Hz, 3H), 0.95 (d, J=6.6Hz, 2.1H), 1.04 (d, J=6.6Hz, 0.9H), 1.50 (m, 2H), 1.62 (m, 1H), 2.12 (m, 1H), 3.43 (m, 1H), 3.70 (m, 1H), 3.91 (m, 1H), 4.17 (dd, J=8.4Hz, 8.4Hz, 1H), 5.12 (d, J=4.5Hz, 0.3H), 5.25 (d, J=4.5Hz, 0.7H), 5.35 (d, J=7.2Hz, 0.7H), 5.40 (d, J=7.2Hz, 0.3H), 6.34 (d, J=8.7Hz, 0.3H), 6.37 (d, J=8.7Hz, 0.7H), 7.49-7.62 (m, 3H), 7.88 (m, 2H).

Example 81: Preparation of (3S)-3-((S)-2-benzyloxycarbonylamino-4-methylvalerylamino)-2-tetrahydropyrazole (Compound No. 433 in Table-1)

IR (KBr, cm^{-1}): 3298, 1691, 1649, 1541.

NMR (CDCl$_3$, δ): 0.90-0.92 (m, 6H), 1.57-1.70 (m, 7H), 3.43-3.70 (m, 2H), 3.87-3.99 (m, 2H), 4.08-4.12 (m, 1H), 5.02 (s, 0.4H), 5.08 (s, 1.6H), 5.66 (s, 1H), 6.57 (s, 0.8H), 6.88 (s, 0.2H), 7.31 (s, 5H).

Example 82: Preparation of (3S)-3-{(S)-2-(2-fluorobenzoylamino)-4-methylvalerylamino}-2-tetrahydropyrazole (Compound No. 450 in Table-1)

IR (KBr, cm^{-1}): 3408, 1600, 1495.

NMR (CDCl$_3$, δ): 0.92-0.98 (m, 6H), 1.64-1.82 (m, 7H), 3.48-3.59 (m, 2H), 3.84-4.13 (m, 2H), 4.68-4.70 (m, 1H),

5.01 (m, 0.3H), 5.05 (m, 0.7H), 6.55 (d, J=8.3Hz, 0.7H), 6.82 (d, J=8.2Hz, 0.3H), 7.07-7.28 (m, 2H), 7.41-7.52 (m, 1H), 7.99-8.05 (m, 1H).

Example 83: Preparation of (3S)-3-((S)-4-methyl-2-phenylsufonylaminovalerylamino)-2-tetrahydropyranole (Compound No. 468 in Table-1)

Melting point: 156-157°C
IR (KBr, cm⁻¹): 3335, 3261, 1649, 1545.
NMR (CDCl₃, δ): 0.67 (d, J=8.5Hz, 3H), 0.85 (d, J=6.0Hz, 3H), 1.21-2.05 (m, 7H), 3.21 (d, J=4.1Hz, 0.8H), 3.41-3.78 (m, 2H), 3.80-4.07 (m, 2H), 4.19 (m, 0.2H), 4.95 (s, 1H), 5.23 (d, J=6.8Hz, 1H), 6.26 (m, 1H), 7.42-7.68 (m, 3H), 7.87 (d, J=8.5Hz, 2H).

Example 84: Preparation of (3S)-3-{(S)-4-methyl-2-(2,4,6-trimethylphenylsulfonylamino)valerylamino}-2-tetrahydropyranol (Compound No. 473 in Table-1)

IR (KBr, cm⁻¹): 3331, 1655, 1541.
NMR (CDCl₃, δ): 0.66-0.72 (m, 3H), 0.83-0.85 (m, 3H), 1.44-1.97 (m, 7H), 2.28 (s, 3H), 2.63 (s, 6H), 3.42-3.75 (m, 2H), 3.84-3.98 (m, 2H), 4.40-4.47 (m, 1H), 4.93 (s, 1H), 5.47 (d, J=8.4Hz, 0.3H), 5.53 (d, J=7.8Hz, 0.7H), 6.34 (m, 1H), 6.94 (s, 2H).

Example 85: Preparation of (3S)-3-{(S)-2-(N-acetyl-N-4-methylphenylsulfonyl)amino-4-methylvalerylamino}-2-tetrahydorofuranol (Compound No. 515 in Table-1)

Melting point: 49-51°C
IR (KBr, cm⁻¹): 3414, 1701, 1674, 1524.
NMR (CDCl₃, δ): 0.81-1.04 (m, 6H), 1.54-2.02 (m, 4H), 2.19-2.55 (m, 1H), 2.25 (s, 1.95H), 2.29 (s, 1.05H), 2.46 (s, 3H), 2.85 (d, J=2.9Hz, 0.35H), 3.23 (d, J=3.2Hz, 0.65H), 3.80-4.20 (m, 2H), 4.27-4.44 (m, 1H), 4.86 (t, J=7.7Hz, 0.35H), 4.97 (dd, J=7.7Hz, 6.3Hz, 0.65H), 5.24 (d, J=2.9Hz, 0.35H), 5.34 (dd, J=3.2Hz, 3.2Hz, 0.65H), 6.02 (d, J=7.4Hz, 0.35H), 6.41 (d, J=7.9Hz, 0.65H), 7.37 (d, J=8.4Hz, 2H), 7.98 (d, J=8.4Hz, 0.7H), 8.05 (d, J=8.4Hz, 1.3H).

Example 86: Preparation of (3S)-3-{(S)-2-(N-acetyl-N-4-methoxyphenylsulfonyl)amino-4-methylvalerylamino}-2-tetrahydrofuranol (Compound No. 516 in Table-1)

Melting point: 48-51°C
IR (KBr, cm⁻¹): 3414, 1701, 1595, 1522, 1501.
NMR (CDCl₃, δ): 0.84-1.06 (m, 6H), 1.57-2.01 (m, 4H), 2.24 (s, 1.95H), 2.29 (s, 1.05H), 2.26-2.58 (m, 1H), 3.04 (d, J=2.6Hz, 0.35H), 3.42 (d, J=3.1Hz, 0.65H), 3.80-4.18 (m, 2H), 3.89 (s, 1.95H), 3.90 (s, 1.05H), 4.24-4.43 (m, 1H), 4.87 (dd, J=7.7Hz, 6.0Hz, 0.35H), 4.95 (t, J=6.9Hz, 0.65H), 5.24 (d, J=2.6Hz, 0.35H), 5.34 (dd, J=3.1Hz, 3.1Hz, 0.65H), 6.04 (d, J=7.1Hz, 0.35H), 6.42 (d, J=8.0Hz, 0.65H), 7.04 (d, J=9.0Hz, 2H), 8.04 (d, J=9.0Hz, 0.7H), 8.12 (d, J=9.0Hz, 1.3H).

Example 87: Preparation of (3S)-3-{(S)-2-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-4-methylvalerylamino]-2-tetrahydrofuranol (Compound No. 1126 in Table-3)

Melting point: 186-188°C
IR (KBr, cm⁻¹): 3285, 1644, 1549.
NMR (CDCl₃, δ): 0.40 (d, J=6.3Hz, 1.2H), 0.52 (d, J=5.6Hz, 1.8H), 0.75-1.00 (m, 9H), 1.32-2.07 (m, 7H), 2.25 (m, 0.6H), 2.40 (m, 0.4H), 3.48 (d, J=2.8Hz, 0.4H), 3.59 (m, 1H), 3.73-3.92 (m, 1.2H), 4.02-4.20 (m, 1.4H), 4.21-4.56 (m, 2H), 5.30-5.38 (m, 1H), 5.59 (d, J=4.8Hz, 0.4H), 5.68 (d, J=5.9Hz, 0.6H), 6.82 (d, J=8.3Hz, 0.6H), 6.95 (d, J=9.7Hz, 0.4H), 6.99 (d, J=8.6Hz, 0.6H), 7.08 (d, J=7.0Hz, 0.4H), 7.47-7.72 (m, 3H), 7.91 (d, J=8.5Hz, 2H).

Example 88: Preparation of (2S,3S)-2-acetoxy-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)tetrahydrofuran (Compound No. 716 in Table-2)

244 mg of (S)-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-tetrahydrofuranone obtained in Reference Example 1 was dissolved in 35 ml of methylene chloride and the solution was cooled to -78°C. 1.91 ml of a solution of diisobutylalminium hydride in toluene (1.01 mol/L) was added to th reaction solution. After stirring for 3 hours at -78°C, the reaction mixture was added with a saturated aqueous solution of ammonium chloride and ethyl acetate. The mixture was warmed up to room temp rature and then filtered through celit . Th celite was thoroughly washed with ethyl ace-

tate. The filtrate was washed with saturated brine, dried over magnesium sulfate, and then filtered. The filtrate was concentrated to obtain crude (3S)-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-tetrahydrofuranol (compound of Example 1). This compound was dissolved in 1 ml of pyridine and added with 1.5 ml of acetic anhydride under ice cooling, and then, the mixture was stirred for 9 hours under ice cooling and further added with 1.5 ml of methanol and concentrated. The resulting residue was dissolved in ethyl acetate, and the solution was washed successively with diluted hydrochloric acid, water, saturated aqueous solution of sodium hydrogencarbonate, and saturated brine, and then dried over magnesium sulfate and filtered. The filtrate was concentrated, and the resulting residue was added with 2.5 ml of ethyl acetate and 2.5 ml of hexane, and then the mixture was stirred. Precipitated crystals were collected by filtration to obtain the desired compound (120 mg).

Yield: 44%

Melting point: 177-178°C

IR (KBr, cm^{-1}): 3409, 3100, 1753, 1659.

NMR (CDCl$_3$, δ): 0.54 (d, J=6.3Hz, 3H), 0.80 (d, J=6.3Hz, 3H), 1.37 (m, 2H), 1.59 (m, 1H), 1.81 (m, 1H), 2.16 (s, 3H), 2.24 (m, 1H), 3.61 (m, 1H), 3.95 (ddd, J=9.3Hz, 9.0Hz, 7.5Hz, 1H), 4.14 (ddd, J=9.3Hz, 9.3Hz, 3.0Hz, 1H), 4.53 (m, 1H), 4.87 (d, J=6.6Hz, 1H), 6.16 (d, J=4.5Hz, 1H), 6.56 (d, J=8.7Hz, 1H), 7.54 (m, 2H), 7.62 (m, 1H), 7.86 (dd, J=7.2Hz, 1.5Hz, 2H).

Compounds of Example 89 to Example 117 were prepared in the same manners as that of Example 88. Physicochemical data of the compounds will be described below.

Example 89: Preparation of (2S,3S)-2-acetoxy-3-((S)-2-tert-butoxycarbonylamino-4-methylvalerylamino)tetrahydrofuran (Compound No. 547 in Table-2)

Melting point: 143-145°C

IR (KBr, cm^{-1}): 3297, 1748, 1659.

NMR (CDCl$_3$, δ): 0.93 (d, J=6.0Hz, 3H), 0.94 (d, J=6.0Hz, 3H), 1.45 (s, 9H), 1.49 (m, 1H), 1.67 (m, 2H), 1.83 (m, 1H), 2.11 (s, 3H), 2.36 (m, 1H), 3.96 (ddd, J=9.3Hz, 9.3Hz, 9.3Hz, 1H), 4.06 (m, 1H), 4.14 (ddd, J=9.3Hz, 9.3Hz, 3.0Hz, 1H), 4.57 (m, 1H), 4.84 (s, 1H), 6.17 (s, J=4.8Hz, 1H), 6.45 (s, 1H).

Example 90: Preparation of (2S,3S)-2-acetoxy-3-((S)-2-benzyloxycarbonylamino-4-methylvalerylamino)tetrahydrofuran (Compound No. 551 in Table-2)

Melting point: 162-164°C

IR (KBr, cm^{-1}): 3310, 1687, 1655, 1535.

NMR (CDCl$_3$, δ): 0.94 (d, J=6.2Hz, 3H), 0.95 (d, J=6.2Hz, 3H), 1.83 (m, 1H), 2.07 (s, 3H), 2.11 (m, 3H), 2.36 (m, 1H), 3.93 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 1H), 4.09 (m, 2H), 4.79 (m, 1H), 5.12 (s, 2H), 5.32 (s, 1H), 6.04 (d, J=4.2Hz, 1H), 6.17 (d, J=4.4Hz, 1H), 7.35 (m, 5H).

Example 91: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-(2-chlorobenzyloxycarbonylamino)-4-methylvalerylamino}tetrahydrofuran (Compound No. 558 in Table-2)

Melting point: 144-147°C

IR (KBr, cm^{-1}): 3314, 3076, 1699, 1655, 1535.

NMR (CDCl$_3$, δ): 0.92-0.95 (m, 6H), 1.51-1.84 (m, 4H), 2.08 (s, 3H), 2.34 (m, 1H), 3.95 (dd, J=8.8Hz, 7.4Hz, 1H), 4.09-4.16 (m, 2H), 4.57 (m, 1H), 5.19 (d, J=13.0Hz, 1H), 5.23 (s, 1H), 5.26 (d, J=13.0Hz, 1H), 6.16 (d, J=4.3Hz, 1H), 6.29 (s, 1H), 7.25-7.29 (m, 2H), 7.37-7.41 (m, 2H).

Example 92: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(4-methylbenzyloxycarbonylamino)valerylamino}tetrahydrofuran (Compound No. 566 in Table-2)

Melting point: 161-163°C

IR (KBr, cm^{-1}): 3314, 1691, 1651, 1539.

NMR (CDCl$_3$, δ): 0.91-0.93 (m, 6H) 1.46 (m, 4H), 2.07 (s, 3H), 2.32 (s, 3H), 2.33 (m, 1H), 3.94 (q, J=7.5Hz, 1H), 4.08-4.15 (m, 2H), 4.57 (m, 1H), 5.05 (d, J=12.0Hz, 1H), 5.06 (s, 1H), 5.09 (d, J=12.0Hz, 1H), 6.15 (d, J=4.2Hz, 1H), 6.32 (s, 1H), 7.15 (d, J=7.8Hz, 2H), 7.23 (d, J=7.8Hz, 2H).

Example 93: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-(9-fluorenylmethoxycarbonylamino)-4-methylvalerylamino}tetrahydrofuran (Compound No. 571 in Table-2)

Melting point: 158-159°C

IR (KBr, cm^{-1}): 3308, 1746, 1692, 1657, 1537.

NMR (CDCl$_3$, δ): 0.94 (m, 6H), 1.52-1.74 (m, 3H), 1.64 (m, 1H), 2.06 (s, 3H), 2.36 (m, 1H), 3.96 (ddd, J=7.8Hz, 7.8Hz, 7.8Hz, 1H), 4.11 (m, 2H), 4.19 (t, J=7.5Hz, 1H), 4.43 (m, 2H), 4.56 (m, 1H), 5.15 (s, 1H), 6.06 (d, J=4.2Hz, 1H), 6.22 (s, 1H), 7.31 (dd, J=7.5Hz, 7.5Hz, 2H), 7.41 (dd, J=7.5Hz, 7.5Hz, 2H), 7.58 (d, J=7.5Hz, 2H), 7.77 (d, J=7.5Hz, 2H).

Example 94: Preparation of (2S,3S)-2-acetoxy-3-((S)-2-cyclohexyloxycarbonylamino-4-methylvalerylamino)tetrahydrofuran (Compound No. 580 in Table-2)

Melting point: 135- 137°C

NMR (CDCl$_3$, δ): 0.92-0.95 (m, 6H), 1.26-1.88 (m, 14H), 2.10 (s, 3H), 2.35 (m, 1H), 3.97 (m, 1H), 4.10-4.16 (m, 2H), 4.51-4.64 (m, 2H), 5.03 (s, 1H), 6.16 (d, J=4.5Hz, 1H), 6.40 (s, 1H).

Example 95: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(1-naphtylacetylamino)valerylamino}tetrahydrofuran (Compound No. 605 in Table-2)

Melting point: 182-184°C

IR (KBr, cm^{-1}): 3308, 1745, 1644, 1551.

NMR (CDCl$_3$, δ): 0.72 (d, J=6.4Hz, 3H), 0.74 (d, J=6.4Hz, 3H), 1.10-1.35 (m, 2H), 1.47 (m, 1H), 1.67 (m, 1H), 2.13 (s, 3H), 2.20 (m, 1H), 3.93 (ddd, J=8.9Hz, 8.9Hz, 8.9Hz, 1H), 3.98-4.15 (m, 3H), 4.31 (m, 1H), 4.46 (m, 1H), 5.66 (d, J=8.1Hz, 1H), 6.12 (d, J=4.6Hz, 1H), 6.53 (d, J=8.6Hz, 1H), 7.05-7.57 (m, 4H), 7.80-7.95 (m, 3H).

Example 96: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-(2-fluorobenzoylamino)-4-methylvalerylamino}tetrahydrofuran (Compound No. 633 in Table-2)

Melting point: 165-166°C.

IR (KBr, cm^{-1}): 3343, 3304, 1748, 1694, 1640, 1551.

NMR (CDCl$_3$, δ): 0.96 (d, J=5.7Hz, 3H), 0.98 (d, J=5.7Hz, 3H), 1.58-1.95 (m, 4H), 2.10 (s, 3H), 2.39 (m, 1H), 3.95 (ddd, J=9.0Hz, 9.0Hz, 7.4Hz, 1H), 4.14 (ddd, J=9.0Hz, 9.0Hz, 2.9Hz, 1H), 4.55-4.75 (m, 2H), 6.19 (d, J=4.8Hz, 1H), 6.61 (d, J=8.2Hz, 1H), 7.00 (d, J=7.4Hz, 0.5H), 7.09 (d, J=7.4Hz, 0.5H), 7.15 (dd, J=8.2Hz, 7.5Hz, 1H), 7.29 (ddd, J=6.8Hz, 6.8Hz, 0.9Hz, 1H), 7.55 (m, 1H), 8.06 (ddd, J=7.9Hz, 7.9Hz, 1.9Hz, 1H).

Example 97: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-(4-chlorobenzoylamino)-4-methylvalerylamino}tetrahydrofuran (Compound No. 641 in Table-2)

Melting point: 204-205°C

IR (KBr, cm^{-1}): 3304, 1746, 1674, 1635.

NMR (CDCl$_3$, δ): 0.96 (d, J=6.1Hz, 3H), 0.97 (d, J=6.0Hz, 3H), 1.58-1.78 (m, 3H), 1.85 (m, 1H), 2.13 (s, 3H), 2.31 (m, 1H), 3.95 (m, 1H), 4.14 (ddd, J=8.6Hz, 8.6Hz, 2.9Hz, 1H), 4.50-4.70 (m, 2H), 6.20 (d, J=4.6Hz, 1H), 6.62 (d, J=8.3Hz, 1H), 6.81 (d, J=7.8Hz, 1H), 7.41 (d, J=6.7Hz, 2H), 7.73 (d, J=6.7Hz, 2H).

Example 98: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(2-methylbenzoylamino)valerylamino}tetrahydrofuran (Compound No. 645 in Table-2)

Melting point: 185-186°C

IR (KBr, cm^{-1}): 3264, 1753, 1674, 1626.

NMR (CDCl$_3$, δ): 0.99 (d, J=5.6Hz, 6H), 1.55-1.98 (m, 4H), 2.11 (s, 3H), 2.30 (m, 1H), 2.49 (s, 3H), 3.98 (m, 1H), 4.15 (ddd, J=6.3Hz, 6.3Hz, 2.9Hz, 1H), 4.47-4.70 (m, 2H), 6.19 (d, J=4.6Hz, 1H), 6.28 (d, J=8.2Hz, 1H), 6.81 (d, J=8.4Hz, 1H), 7.16-7.29 (m, 2H), 7.31-7.40 (m, 2H).

Example 99: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(4-methylbenzoylamino)valerylamino}tetrahydrofuran (Compound No. 647 in Table-2)

Melting point: 199-200°C

IR (KBr, cm^{-1}): 3318, 1746, 1663, 1630, 1534.

NMR (CDCl$_3$, δ): 0.96 (d, J=6.0Hz, 3H), 0.97 (d, J=6.0Hz, 3H), 1.59-1.95 (m, 4H), 2.12 (s, 3H), 2.29 (m, 1H), 2.40 (s, 3H), 3.95 (m, 1H), 4.13 (ddd, J=6.3Hz, 6.3Hz, 2.9Hz, 1H), 4.49-4.70 (m, 2H), 6.19 (d, J=4.6Hz, 1H), 6.58 (d, J=7.6Hz, 1H), 6.70 (d, J=7.6Hz, 1H), 7.24 (d, J=7.8Hz, 2H), 7.69 (d, J=7.8Hz, 2H).

Example 100: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(2,4,6-trimethylbenzoylamino)valerylamino}tetrahydrofuran (Compound No. 651 in Table-2)

NMR (CDCl$_3$, δ): 0.97 (d, J=5.0Hz, 6H), 1.59-1.79 (m, 3H), 1.89 (m, 1H), 2.11 (s, 3H), 2.27 (s, 6H), 2.32 (s, 3H), 2.35 (m, 1H), 3.95 (m, 1H), 4.14 (ddd, J=9.0Hz, 9.0Hz, 2.8Hz, 1H), 4.55-4.70 (m, 2H), 5.98 (d, J=8.1Hz, 1H), 6.09 (d, J=4.6Hz, 1H), 6.84 (s, 2H), 6.86 (d, J=7.5Hz, 1H).

Example 101: Preparation of (2S,3S)-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-poropionyloxytetrahydrofuran (Compound No. 720 in Table-2)

Melting point: 154-156°C
IR (KBr, cm^{-1}): 3355, 3274, 1711, 1678.
NMR (CDCl$_3$, δ): 0.56 (d, J=6.3Hz, 3H), 0.80 (d, J=6.6Hz, 3H), 1.17 (t, J=7.5Hz, 3H), 1.39 (m, 2H), 1.57 (m, 1H), 1.79 (m, 1H), 2.23 (m, 1H), 2.40 (qd, J=7.5Hz, 16.8Hz, 1H), 2.49 (qd, J=7.5Hz, 16.8Hz, 1H), 3.62 (m, 1H), 3.94 (ddd, J=9.0Hz, 9.0Hz, 9.0Hz, 1H), 4.13 (ddd, J=9.3Hz, 9.0Hz, 3.0Hz, 1H), 4.92 (d, J=6.9Hz, 1H), 6.17 (d, J=4.8Hz, 1H), 6.49 (d, J=8.7Hz, 1H), 7.53 (m, 2H), 7.62 (m, 1H), 7.86 (d, J=6.0Hz, 2H).

Example 102: Preparation of (2S,3S)-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)-2-pivaloyloxytetrahydrofuran (Compound No. 725 in Table-2)

Melting point: 165-166°C
IR (KBr, cm^{-1}): 3293, 1640.
NMR (CDCl$_3$, δ): 0.64 (d, J=5.9Hz, 3H), 0.81 (d, J=6.1Hz, 3H), 1.26 (s, 9H), 1.48 (m, 3H), 1.72 (m, 1H), 2.23 (m, 1H), 3.64 (m, 1H), 3.94 (ddd, J=9.1Hz, 9.1Hz, 9.1Hz, 1H), 4.10 (ddd, J=9.1Hz, 9.1Hz, 3.1Hz, 1H), 4.45 (m, 1H), 5.07 (d, J=7.7Hz, 1H), 6.10 (d, J=4.5Hz, 1H), 6.21 (d, J=8.4Hz, 1H), 7.51 (m, 2H), 7.61 (m, 1H), 7.86 (d, J=7.1Hz, 2H).

Example 103: Preparation of (2S,3S)-2-benzoyloxy-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)tetrahydrofuran (Compound No. 728 in Table-2)

Melting point: 184-185°C
IR (KBr, cm^{-1}): 3353, 3260, 1698, 1678.
NMR (CDCl$_3$, δ): 0.55 (d, J=6.0Hz, 3H), 0.69 (d, J=6.0Hz, 3H), 1.30-1.48 (m, 3H), 1.90 (m, 1H), 2.30 (m, 1H), 3.64 (m, 1H), 4.00 (ddd, J=9.3Hz, 9.0Hz, 7.5Hz, 1H), 4.19 (ddd, J=9.3Hz, 9.3Hz, 3.0Hz, 1H), 4.51 (m, 1H), 5.13 (d, J=8.1Hz, 1H), 6.37 (d, J=6.6Hz, 1H), 6.39 (d, J=4.2Hz, 1H), 7.26-7.50 (m, 4H), 7.58 (m, 2H), 7.80 (dd, J=7.5Hz, 1.8Hz, 2H), 8.06 (dd, J=7.8Hz, 0.9Hz, 2H).

Example 104: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-(4-chlorophenylsulfonylamino)-4-methylvalerylamino}tetrahydrofuran (Compound No. 754 in Table-2)

Melting point: 136-137°C
IR (KBr, cm^{-1}): 3335, 3258, 1744, 1651, 1535.
NMR (CDCl$_3$, δ): 0.61 (d, J=6.2Hz, 3H), 0.83 (d, J=6.3Hz, 3H), 1.35-1.62 (m, 3H), 1.80 (m, 1H), 2.15 (s, 3H), 2.21 (m, 1H), 3.61 (m, 1H), 3.97 (ddd, J=9.1Hz, 9.1Hz, 9.1Hz, 1H), 4.14 (ddd, J=9.1Hz, 9.1Hz, 2.9Hz, 1H), 4.50 (m, 1H), 5.09 (d, J=7.3Hz, 1H), 6.15 (d, J=4.6Hz, 1H), 6.42 (d, J=8.8Hz, 1H), 7.51 (d, J=8.6Hz, 2H), 7.80 (d, J=8.6Hz, 2H).

Example 105: Preparation of (2S,3S)-3-{(S)-2-(4-chlorophenylsulfonylamino)-4-methylvalerylamino}-2-pivaloyloxytetrahydrofuran (Compound No. 755 in Table-2)

Melting point: 155-156°C
IR (KBr, cm^{-1}): 3366, 3229, 1726, 1684, 1664, 1543.
NMR (CDCl$_3$, δ): 0.70 (d, J=6.3Hz, 3H), 0.83 (d, J=6.4Hz, 3H), 1.25 (s, 9H), 1.40-1.82 (m, 4H), 2.23 (m, 1H), 3.62 (dd, J=12.5Hz, 6.4Hz, 1H), 3.96 (ddd, J=8.9Hz, 8.9Hz, 8.9Hz, 1H), 4.11 (ddd, J=8.9Hz, 8.9Hz, 2.9Hz, 1H), 4.45 (m, 1H), 5.21 (d, J=8.2Hz, 1H), 6.04 (d, J=8.6Hz, 1H), 6.10 (d, J=4.5Hz, 1H), 7.45-7.55 (m, 2H), 7.75-7.85 (m, 2H).

Example 106: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(4-methylphenylsulfonylamino)valerylamino}tetrahydrofuran (Compound No. 761 in Table-2)

Melting point: 159-160°C

IR (KBr, cm^{-1}): 3372, 1721, 1674, 1535.

NMR (CDCl$_3$, δ): 0.52 (d, J=6.2Hz, 3H), 0.80 (d, J=6.3Hz, 3H), 1.36 (m, 2H), 1.56 (m, 1H), 1.83 (m, 1H), 2.16 (s, 3H), 2.22 (m, 1H), 2.44 (s, 3H), 3.59 (m, 1H), 3.94 (m, 1H), 4.14 (m, 1H), 4.56 (m, 1H), 4.81 (d, J=6.6Hz, 1H), 6.15 (d, J=4.6Hz, 1H), 6.35 (d, J=7.8Hz, 1H), 7.33 (d, J=8.0Hz, 2H), 7.74 (d, J=8.0Hz, 2H).

Example 107: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(2,4,6-trimethylphenylsulfonylamino)valerylamino}tetrahydrofuran (Compound No. 765 in Table-2)

Melting point: 158-159°C

IR (KBr, cm^{-1}): 3416, 3191, 1755, 1661, 1605, 1535.

NMR (CDCl$_3$, δ): 0.56 (d, J=6.3Hz, 3H), 0.79 (d, J=6.3Hz, 3H), 1.39 (m, 2H), 1.58 (m, 1H), 1.81 (m, 1H), 2.16 (s, 3H), 2.24 (m, 1H), 2.31 (s, 3H), 2.62 (s, 6H), 3.56 (m, 1H), 3.94 (m, 1H), 4.14 (m, 1H), 4.52(m, 1H), 5.00 (d, J=7.1Hz, 1H), 6.15 (d, J=4.6Hz, 1H), 6.61 (d, J=8.7Hz, 1H), 6.97(s, 2H).

Example 108: Preparation of (2S,3S)-2-acetyl-3-{(S)2-(4-tert-butylphenylsulfonylamino)-4-methylvalerylamino)tetrahydrofuran (Compound No. 768 in Table-2)

NMR (CDCl$_3$, δ): 0.44 (d, J=6.2Hz, 3H), 0.77 (d, J=6.2Hz, 3H), 1.23-1.43 (m, 2H), 1.34 (s, 9H), 1.55 (m, 1H), 1.85 (m, 1H), 2.17 (s, 3H), 2.20 (m, 1H), 3.59 (m, 1H), 3.95 (m, 1H), 4.13 (ddd, J=9.0Hz, 9.0Hz, 2.9Hz, 1H), 4.57 (m, 1H), 4.95 (d, J=6.4Hz, 1H), 6.17 (d, J=4.6Hz, 1H), 6.77(d, J=8.7Hz, 1H), 7.53 (d, J=8.6Hz, 2H), 7.78 (d, J=8.6Hz, 2H).

Example 109: Preparation of (2S,3S)-2-acetyl-3-{(S)-2-(4-methoxyphenylsulfonylamino)-4-methylvalerylamino)tetrahydrofuran (Compound No. 771 in Table-2)

Melting point: 157-158°C

IR (KBr, cm^{-1}): 3329, 3273, 1746, 1659, 1597, 1544, 1501.

NMR (CDCl$_3$, δ): 0.54 (d, J=6.1Hz, 3H), 0.81 (d, J=6.3Hz, 3H), 1.38 (m, 2H), 1.56 (m, 1H), 1.84 (m, 1H), 2.16 (s, 3H), 2.22 (m, 1H), 3.56 (m, 1H), 3.88 (s, 3H), 3.95 (m, 1H), 4.15 (m, 1H), 4.56 (m, 1H), 4.81 (d, J=6.4Hz, 1H), 6.16 (d, J=4.5Hz, 1H), 6.68 (d, J=9.2Hz, 1H), 6.99 (d, J=8.6Hz, 2H), 7.79 (d, J=8.6Hz, 2H).

Example 110: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(2-naphtylsulfonylamino)valerylamino}tetrahydrofuran (Compound No. 780 in Table-2)

Melting point: 158-159°C

IR (KBr, cm^{-1}): 3337, 3275, 1723, 1676, 1543.

NMR (CDCl$_3$, δ): 0.48 (d, J=6.1Hz, 3H), 0.76 (d, J=6.2Hz, 3H), 1.33-1.75 (m, 4H), 1.97 (m, 1H), 2.15 (s, 3H), 3.68 (m, 1H), 3.87 (m, 1H), 4.05 (m, 1H), 4.42 (m, 1H), 5.19 (d, J=7.1Hz, 1H), 6.12 (d, J=4.6Hz, 1H), 6.53 (d, J=8.7Hz, 1H), 7.58-7.71 (m, 2H), 7.82 (dd, J=8.7Hz, 1.9Hz, 1H), 7.87-8.03 (m, 3H), 8.44 (s, 1H).

Example 111: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(3-pyridylsulfonylamino)valerylamino}tetrahydrofuran (Compound No. 782 in Table-2)

Melting point: 160-161°C

IR (KBr, cm^{-1}): 3410, 3075, 1753, 1657, 1535, 1342, 1172.

NMR (CDCl$_3$, δ): 0.68 (d, J=5.9Hz, 3H), 0.85 (d, J=5.9Hz, 3H), 1.40-1.62 (m, 3H), 1.80 (m, 1H), 2.14 (s, 3H), 2.22 (m, 1H), 3.73 (m, 1H), 3.96 (ddd, J=9.1Hz, 8.9Hz, 8.9Hz, 1H), 4.03 (ddd, J=9.1Hz, 9.1Hz, 3.0Hz, 1H), 4.45 (m, 1H), 5.46 (m, 1H), 6.13 (d, J=4.6Hz, 1H), 6.32 (d, J=8.6Hz, 1H), 7.48 (m, 1H), 8.16 (ddd, J=8.2Hz, 1.9Hz, 1.9Hz, 1H), 8.83 (dd, J=4.9Hz, 1.5Hz, 1H), 9.07 (d, J=2.3Hz, 1H).

Example 112: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-[N-acetyl-N-(4-methylphenylsulfonyl)amino]-4-methylvalerylamino}tetrahydrofuran (Compound No. 1065 in Table-2)

Melting point: 154°C

IR (KBr, cm^{-1}): 3387, 1746, 1674, 1522.

NMR (CDCl$_3$, δ): 0.84 (d, J=6.3Hz, 3H), 0.93 (d, J=6.3Hz, 3H), 1.58 (m, 2H), 1.84 (m, 1H), 2.09 (m, 1H), 2.12 (s, 3H), 2.33 (s, 3H), 2.38 (m, 1H), 2.47 (s, 3H), 3.96 (m, 1H), 4.14 (m, 1H), 4.55 (m, 1H), 4.78 (t, J=6.8Hz, 1H), 6.14 (d, J=4.6Hz, 1H), 6.50 (d, J=8.3Hz, 1H), 7.39 (d, J=8.1Hz, 2H), 7.91 (d, J=8.1Hz, 2H).

5   Example 113: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-[N-acetyl-N-(4-methoxyphenylsulfonyl)amino]-4-methylvaler-ylamino}tetrahydrofuran (Compound No. 1066 in Table-2)

Melting point: 64-66°C
IR (KBr, cm^{-1}): 3397, 1748, 1595, 1530.

10   NMR (CDCl$_3$, δ): 0.85 (d, J=6.2Hz, 3H), 0.94 (d, J=6.3Hz, 3H), 1.59 (m, 2H), 1.85 (m, 1H), 2.08 (m, 1H), 2.12 (s, 3H), 2.32 (s, 1H), 2.37 (m, 1H), 3.90 (s, 3H), 3.96 (m, 1H), 4.15 (m, 1H), 4.56 (m, 1H), 4.79 (t, J=6.8Hz, 1H), 6.15 (d, J=4.7Hz, 1H), 6.51 (d, J=8.4Hz, 1H), 7.04 (d, J=9.0Hz, 2H), 7.97 (d, J=9.0Hz, 2H).

Example 114: Preparation of (2S,3S)-2-acetoxy-3-((S)-2-benzyloxycarbonylamino-4-methylvalerylamino)-2-tetrahydro-
15   furan (Compound No. 983 in Table-2)

IR (KBr, cm^{-1}): 3310, 1693, 1653, 1537.
NMR (CDCl$_3$, δ): 0.90-0.93 (m, 6H), 1.43-1.77 (m, 7H), 2.11 (s, 3H), 3.65-3.74 (m, 2H), 4.08-4.17 (m, 2H), 5.09 (s, 2H), 5.30 (d, J=8.1Hz, 1H), 5.96 (d, J=2.7Hz, 1H), 6.17 (s, 1H), 7.34 (m, 5H).

20

Example 115: Preparation of (2S,3S)-2-acetoxy-3-{(S)-2-(2-fluorobenzoylamino)-4-methylvalerylamino}-2-tetrahydro-pyran (Compound No. 1000 in Table-2)

IR (KBr, cm^{-1}): 3298, 2947, 1633, 1545.
25   NMR (CDCl$_3$, δ): 0.91-0.98 (m, 6H), 1.65-1.77 (m, 7H), 2.13 (s, 3H), 3.64-3.77 (m, 2H), 3.90 (m, 1H), 4.20 (m, 1H), 4.60 (s, 1H), 6.00 (d, J=3.3Hz, 1H), 6.42 (s, 1H), 7.13 (m, 1H), 7.26 (m, 1H), 7.48 (m, 1H), 8.02 (m, 1H).

Example 116: Preparation of (2S,3S)-2-acetoxy-3-{(S)-4-methyl-2-(2,4,6-trimethylphenylsulfonylamino)valerylamino}-2-tetrahydoropyran (Compound No. 1023 in Table-2)
30

IR (KBr, cm^{-1}): 3418, 1658, 1606, 1523. NMR (CDCl$_3$, δ): 0.57 (d, J=6.3Hz, 3H), 0.79 (d, J=6.3Hz, 3H), 1.32-1.78 (m, 7H), 2.16 (s, 3H), 2.29 (s, 3H), 2.60 (s, 6H), 3.49 (m, 1H), 3.61-3.78 (m, 2H), 4.08 (m, 1H), 5.08 (d, J=7.2Hz, 1H), 5.94 (d, J=3.0Hz, 1H), 6.33 (d, J=8.7Hz, 1H), 6.95 (s, 2H).

35   Example 117: Preparation of (2S,3S)-3-{(S)-2-(4-tert-butylphenylsulfonylamino)-4-methylvalerylamino}-2-pivaloy-loxytetrahydrofuran (Compound No. 1397 in Table-2)

Melting point: 166-167°C
IR (KBr, cm^{-1}): 3360, 1728, 1682, 1665, 1547.
40   NMR (CDCl$_3$, δ): 0.55 (d, J=6.0Hz, 3H), 0.78 (d, J=6.0Hz, 3H), 1.25 (s, 9H), 1.32 (s, 9H), 1.38-1.82 (m, 4H), 2.25 (m, 1H), 3.60 (m, 1H), 3.95 (m, 1H), 4.12 (ddd, J=9.0Hz, 9.0Hz, 3.0Hz, 1H), 4.46 (m, 1H), 5.05 (d, J=7.1Hz, 1H), 6.10 (d, J=4.5Hz, 1H), 6.41 (d, J=8.3Hz, 1H), 7.52 (d, J=8.6Hz, 2H), 7.78 (d, J=8.6Hz, 2H).

Example 118: Preparation of (3S)-2-methoxy-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)tetrahydrofuran
45   (Compound No. 741 in Table-2)

43 mg of (2S,3S)-2-acetoxy-3-((S)-4-methyl-2-phenylsulfonylaminovalerylamino)tetrahydrofuran obtained in Example 88 was dissolved in 40 ml of methanol, and the solution was added with 1 ml of ethyl acetate containing 4N hydrochloric. After stirring overnight at room temperature, 7 ml of saturated aqueous solution of sodium hydrogencar-
50   bonate was added to the reaction mixture, and then the solvent was evaporated. The resulting residue was added with a saturated aqueous solution of sodium hydrogencarbonate and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine and dried over magnesium sulfate. The drying agent was removed by filtration and the filtrate was concentrated to obtain a crude product. The crude product was added with hexane and diethyl ether and then stirred. After filtration, the desired compound (29 mg) was obtained.
55

Yield: 65%
Melting point: 84-90°C
IR (KBr, cm^{-1}): 3268, 1647, 1618.
NMR (CDCl$_3$, δ): 0.71 (d, J=6.6Hz, 2.1H), 0.80 (d, J=6.6Hz, 0.9H), 0.84 (d, J=6.6Hz, 2.1H), 0.88 (d, J=6.6Hz, 0.9H),

1.42-1.49 (m, 3H), 1.61 (m, 1H), 2.08 (m, 0.3H), 2.27 (m, 0.7H), 3.30 (s, 2.1H), 3.37 (s, 0.9H), 3.63 (m, 1H), 3.69 (m, 0.3H), 3.81-4.11 (m, 1.7H), 4.13 (m, 1H), 4.66 (s, 0.7H), 4.71 (d, J=4.8Hz, 0.3H), 4.66 (d J=7.8Hz, 0.7H), 5.23 (d, J=8.4Hz, 0.3H), 5.96 (d, J=7.8Hz, 0.7H), 6.17 (d, J=8.7Hz, 0.3H), 7.47-7.62 (m, 3H), 7.86 (d, J=7.5Hz, 2H).

Test Example 1: Measurement of inhibitory activity against cysteine proteases

Inhibitory activity against cathepsin B (Sigma, C-6286) was determined according to a method described in a literature (Biochemical Journal, Vol. 201, p. 189, 1982). The results are shown in Table-5.

m-Calpain was purified from brains of rats according to a method described in a literature (Journal of Biological Chemistry, Vol. 259, p. 3210, 1984), and the inhibitory activity was determined in a manner according to a published method (Journal of Biological Chemistry, Vol. 259, p. 12489, 1984). The results are shown in Table-6.

From the results shown in Table-5 and Table-6, it is apparent that the compounds of the present invention have potent inhibitory activity against cystein proteases such as papain, cathepsin B, cahepsin L, and calpain.

Table-5

Inhibitory activity against cathepsin B			
Example No. (Compound No. in Table-1)	IC_{50} (μM)	Example No. (Compound No. in Table-1)	IC_{50} (μM)
1(196)	0.42	40(113)	1.00
4(20)	0.70	41(114)	0.87
5(21)	0.98	42(115)	0.55
6(22)	1.35	43(121)	0.14
7(23)	0.90	44(125)	1.45
10(27)	0.37	45(126)	0.49
11(28)	2.15	46(127)	0.27
12(29)	1.20	48(130)	0.086
13(31)	0.38	50(134)	0.20
15(37)	0.70	51(137)	0.21
16(38)	1.00	53(140)	0.15
17(40)	0.64	55(142)	1.65
18(44)	0.58	56(144)	0.90
19(46)	0.89	57(148)	0.24
20(47)	1.05	58(152)	0.46
21(49)	1.17	59(156)	0.80
24(53)	2.90	60(157)	0.034
25(54)	2.90	61(161)	1.20
27(60)	1.20	62(170)	0.50
28(61)	1.70	65(187)	0.19
33(85)	0.95	75(229)	2.40
34(90)	1.35	77(232)	0.40
39(112)	0.20	78(246)	0.30
Example No. (Compound No. in Table-3)	IC_{50} (μM)		
87(1126)	1.30		